

REGULAR STRENGTH ASPIRIN 325 MG- aspirin tablet, coated
Kroger

Kroger Aspirin 325 mg Tablets

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscle pain
- toothache
- menstrual pain
- colds
- minor pain of arthritis

temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present

new symptoms occur

ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? Call 1-800-632-6900

Kroger health NDC 41226-011-32
COMPARE TO THE ACTIVE INGREDIENT IN GENUINE BAYER® ASPIRIN
REGULAR STRENGTH Aspirin 325 mg
 FOR ADULTS ACTUAL SIZE
 SAFE PAIN RELIEF PLUS LIFESAVING BENEFITS
PAIN RELIEVER/FEVER REDUCER (NSAID)*
300 COATED TABLETS

Talk to your doctor or other healthcare provider before using this product for your heart. TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Drug Facts
Active ingredient (in each tablet) *Purposes*
 Aspirin 325 mg (NSAID)*Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses
 temporarily relieves minor aches and pains due to:
 ■ headache ■ muscle pain
 ■ toothache ■ menstrual pain
 ■ colds ■ minor pain of arthritis
 ■ temporarily reduces fever

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued under label)
 Bayer® is a registered trademark of Bayer Aktiengesellschaft. Bayer Aktiengesellschaft is not affiliated with The Kroger Co. or this product.
DISTRIBUTED BY THE KROGER CO.
 CINCINNATI, OHIO 45202
 For More Product Information, Scan QR Using Your Kroger App or Call 1-800-632-6900
 YOUR PHARMACIST RECOMMENDED 0 4 1260 02 186 2
 LOT: EXP: **PEEL HERE**
Varnish Omit Area

Drug Facts (continued)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed
Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for
 ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint
 ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
 ■ children under 12 years: consult a doctor

Other Information
 ■ store at 25°C (77°F); excursions permitted between 15°-30° C (59°-86° F)
 ■ use by expiration date on package

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? Call 1-800-632-6900

Adhesive Area

REGULAR STRENGTH ASPIRIN 325 MG			
aspirin tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-011
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
Inactive Ingredients			
	Ingredient Name		Strength
	MINERAL OIL (UNII: T5L8T28FGP)		
	STARCH, CORN (UNII: O8232NY3S)		
	HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
	TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
	MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL011
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-011-32	300 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/17/2024	

Labeler - Kroger (006999528)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
Time Cap Laboratories, Inc.		037052099	manufacture(41226-011)

Revised: 4/2024

Kroger