

MAXIMUM STRENGTH DAYTIME COLD AND FLU AND NIGHTTIME SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, triprolidine hydrochloride
TARGET CORPORATION

Target Daytime Cold & Flu and Nighttime Severe Cold & Flu Liquid

Drug Facts

Daytime Cold & Flu

Daytime Cold & Flu

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

Nighttime Severe Cold and Flu

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Tripolidine HCl 2.5 mg

Purposes

Daytime Cold & Flu

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Nighttime Severe Cold & Flu

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Antihistamine

Uses

Daytime Cold & Flu

temporarily relieves these common cold and flu symptoms:

- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- stuffy nose
- sinus congestion and pressure

temporarily reduces fever

helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Nighttime Severe Cold & Flu

temporarily relieves these common cold and flu symptoms:

- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- sinus congestion and pressure
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes due to hay fever

temporarily reduces fever

controls cough to help you get to sleep

Warnings

Liver warning

Daytime Cold & Flu

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Nighttime Severe Cold & Flu

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

Daytime Cold & Flu

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Nighttime Severe Cold & Flu

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

Daytime Cold & Flu

- taking the blood thinning drug warfarin

Nighttime Severe Cold and Flu

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children (***Nightshift Severe Cold & Flu only***)
- marked drowsiness may occur (***Nightshift Severe Cold & Flu only***)
- alcohol, sedatives, and tranquilizers may increase drowsiness (***Nightshift Severe Cold & Flu only***)
- avoid alcoholic drinks (***Nightshift Severe Cold & Flu only***)
- use caution when driving a motor vehicle or operating machinery (***Nightshift Severe Cold & Flu only***)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

Overdose warning

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Daytime Cold & Flu

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Nighttime Severe Cold and Flu

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

Daytime Cold & Flu

- **each 20 mL contains:** sodium 12 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Nighttime Severe Cold and Flu

- **each 20 mL contains:** sodium 17 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

Daytime Cold & Flu

anhydrous citric acid, edetate disodium, FD&C blue 1, FD&C red 40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol, sucralose, xanthan gum

Nighttime Severe Cold and Flu

anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

Call **1-800-910-6874**

PRINCIPAL DISPLAY PANEL - Kit Carton

Compare to the active ingredients in Maximum Strength Mucinex[®] Fast-Max[™] Daytime Cold & Flu Liquid[®]

Maximum Strength Daytime Cold & Flu Acetaminophen
(Pain Reliever / Fever Reducer)
Dextromethorphan HBr (Cough Suppressant)
Guaifenesin
Phenylephrine HCl (Nasal Decongestant)

ALL IN ONE[™]
Headache, sore throat, chest congestion, sinus pressure, sinus congestion, body pain, fever, cough, nasal congestion

up&up

6 FL OZ (177 mL)
TOTAL 12 FL OZ (354 mL) - 2 x 6 FL OZ

Ages 12+ Years

Compare to the active ingredients in Maximum Strength Mucinex[®] Nightshift Severe Cold & Flu Liquid[®]

Maximum Strength Nighttime Severe Cold & Flu Acetaminophen
(Pain Reliever / Fever Reducer)
Dextromethorphan HBr (Cough Suppressant)
Phenylephrine HCl (Nasal Decongestant)
Triprolidine HCl (Antihistamine)

ALL IN ONE[™]
Fever, cough, nasal congestion, body pain, sore throat, itchy throat, runny nose, sneezing, headache

Ages 12+ Years

6 FL OZ (177 mL)

Print/Varnish Omit Area

LOT: _____ EXP: _____

DAYTIME COLD & FLU		NIGHTTIME SEVERE COLD & FLU	
Drug Facts	Active ingredients (in each 20 mL)	Drug Facts	Active ingredients (in each 20 mL)
<p>Warnings:</p> <p>Do not use if you are taking the blood thinning drug warfarin.</p> <p>When using this product do not use more than directed.</p> <p>Stop use and ask a doctor if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur.</p> <p>Directions:</p> <p>Do not take more than directed (see Directions warning). Do not take more than 2 doses in 24-hour period. Do not use more than 4 doses in 24-hour period. Do not use more than 12 doses in 24-hour period.</p>	<p>Warnings:</p> <p>Do not use if you are taking the blood thinning drug warfarin.</p> <p>When using this product do not use more than directed.</p> <p>Stop use and ask a doctor if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur.</p> <p>Directions:</p> <p>Do not take more than directed (see Directions warning). Do not take more than 2 doses in 24-hour period. Do not use more than 4 doses in 24-hour period. Do not use more than 12 doses in 24-hour period.</p>	<p>Warnings:</p> <p>Do not use if you are taking the blood thinning drug warfarin.</p> <p>When using this product do not use more than directed.</p> <p>Stop use and ask a doctor if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur.</p> <p>Directions:</p> <p>Do not take more than directed (see Directions warning). Do not take more than 2 doses in 24-hour period. Do not use more than 4 doses in 24-hour period. Do not use more than 12 doses in 24-hour period.</p>	<p>Warnings:</p> <p>Do not use if you are taking the blood thinning drug warfarin.</p> <p>When using this product do not use more than directed.</p> <p>Stop use and ask a doctor if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur. Do not use if you experience dizziness, confusion, or sleepiness occur.</p> <p>Directions:</p> <p>Do not take more than directed (see Directions warning). Do not take more than 2 doses in 24-hour period. Do not use more than 4 doses in 24-hour period. Do not use more than 12 doses in 24-hour period.</p>

094101085 801 020282-01-08-0000

1 1673 28601 2

6 FL OZ (177 mL)

Compare to the active ingredients in Maximum Strength Mucinex[®] Fast-Max[™] Daytime Cold & Flu Liquid[®]

Maximum Strength Daytime Cold & Flu Acetaminophen
(Pain Reliever / Fever Reducer)
Dextromethorphan HBr (Cough Suppressant)
Guaifenesin
Phenylephrine HCl (Nasal Decongestant)

ALL IN ONE[™]
Headache, sore throat, chest congestion, sinus pressure, sinus congestion, body pain, fever, cough, nasal congestion

up&up

6 FL OZ (177 mL)

Ages 12+ Years

FEEL BETTER TO READ COMPLETE DRUG FACTS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF BREAKBACK ON BOTTLE CAP IS BROKEN OR MISSING

Maximum Strength per 4-hour dose
*Helps to relieve these symptoms day or night

PARENTS:
Learn about teen medicine about
www.SciMedicineAbout.org

NOT FOR RESALE

094101085 801
C-020282-01-08-0000
Distributed by Target Corporation
Minneapolis, MN 55403
Made in India
TM & ©2024 Target Brands, Inc.
NDC 11673-286-03 768, 0524

LOT: _____ EXP: _____

Print/Varnish Omit Area

Drug Facts

Active ingredients (in each 20 mL)

Purpose

Uses

Warnings

Directions

Other information

Drug Facts (continued)

Directions

Other information

Questions or comments?

Drug Facts (continued)

Directions

Other information

Questions or comments?

Drug Facts (continued)

Directions

Other information

Questions or comments?

Compare to the active ingredients in Maximum Strength Mucinex[®] Nightshift Severe Cold & Flu Liquid[®]

Maximum Strength Nighttime Severe Cold & Flu Acetaminophen
(Pain Reliever / Fever Reducer)
Dextromethorphan HBr (Cough Suppressant)
Phenylephrine HCl (Nasal Decongestant)
Triprolidine HCl (Antihistamine)

ALL IN ONE[™]
Fever, cough, nasal congestion, body pain, sore throat, itchy throat, runny nose, sneezing, headache

up&up

6 FL OZ (177 mL)

Ages 12+ Years

FEEL BETTER TO READ COMPLETE DRUG FACTS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF BREAKBACK ON BOTTLE CAP IS BROKEN OR MISSING

Maximum Strength per 4-hour dose
*Helps to relieve these symptoms at night

PARENTS:
Learn about teen medicine about
www.SciMedicineAbout.org

NOT FOR RESALE

094101085 801
C-020282-01-08-0000
Distributed by Target Corporation
Minneapolis, MN 55403
Made in India
TM & ©2024 Target Brands, Inc.
NDC 11673-286-03 768, 0524

LOT: _____ EXP: _____

Print/Varnish Omit Area

Drug Facts

Active ingredients (in each 20 mL)

Purpose

Uses

Warnings

Directions

Other information

Drug Facts (continued)

Directions

Other information

Questions or comments?

Drug Facts (continued)

Directions

Other information

Questions or comments?

Drug Facts (continued)

Directions

Other information

Questions or comments?

MAXIMUM STRENGTH DAYTIME COLD AND FLU AND NIGHTTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, triprolidine hydrochloride kit

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-286

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-286-01	1 in 1 CARTON	05/01/2024	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	180 mL
Part 2	1 BOTTLE	180 mL

Part 1 of 2

NIGHTTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride solution

Product Information

Item Code (Source)	NDC:11673-872
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-872-03	180 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Part 2 of 2

DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:11673-295
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	blue	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-295-03	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment			
Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(11673-286)