

CURATUSSIN- guaifenesin solution
Wittman Pharma, Inc.

CuraTUSSIN

Dosage & Admin

Directions:

- do not take more than 6 doses in any 24-hour period.

AGE	DOSE
Adults & Children 12 years and over	2-4 teaspoonfuls every 4 hours
Children 6 years to under 12 years	1-2 teaspoonfuls every 4 hours
Children 2 to under 6 years	1/2 - 1 teaspoonfuls every 4 hours
Children Under 2 years	Ask a Doctor

Warnings

Warnings:

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic, such as occurs with smoking, asthma, Chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Inactive Ingredients: Citric Acid, Flavor, Methylparaben, Monoammonium Glycyrrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose

Uses:

- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Purpose: Expectorant

Active Ingredient (in each 5 mL tsp)

Guaifenesin 100 mg

<p>Drug Facts</p> <p>Active Ingredient (in each 5 mL tsp) Purpose Guaifenesin 100 mg Expectorant</p> <p>Uses: ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive</p> <p>Warnings: Ask a doctor before use if you have ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic, such as occurs with smoking, asthma, chronic bronchitis, or emphysema</p> <p>Stop use and ask doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><small>*Robitussin® is a registered trademark of Wyeth Consumer Healthcare. Rev. 3/2024</small></p> 	<p>MADE IN USA  NDC 83335-101-16</p> <p>CuraTUSSIN EXPECTORANT</p> <p>GUAIFENESIN 100 mg / 5 mL</p> <p>LOOSENS AND RELIEVES CHEST CONGESTION</p> <p>Compare to the active ingredient in *Robitussin®</p> <p>✓ SUGAR FREE ✓ ALCOHOL FREE ✓ DYE FREE ✓ GLUTEN FREE</p> <p>Grape Flavor</p> <p> Wittman Pharma</p> <p>Net Content: 16 fl oz (473 mL)</p>	<p>Drug Facts (continued)</p> <p>Directions: ■ do not take more than 6 doses in any 24-hour period</p> <table border="1"> <thead> <tr> <th>Age</th> <th>Dose</th> </tr> </thead> <tbody> <tr> <td>adults and children 12 years and over</td> <td>2-4 teaspoonfuls every 4 hours</td> </tr> <tr> <td>children 6 years to under 12 years</td> <td>1-2 teaspoonfuls every 4 hours</td> </tr> <tr> <td>children 2 years to under 6 years</td> <td>½-1 teaspoonfuls every 4 hours</td> </tr> <tr> <td>under 2 years</td> <td>ask a doctor</td> </tr> </tbody> </table> <p>Other information ■ store at 15°-30°C (59°-86°F) ■ tamper evident, do not use if safety seal under cap is broken or missing</p> <p>Inactive Ingredients: Citric Acid, Flavor, Methylparaben, Monoammonium Glycyrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose</p> <p>Questions? Call 1-352-549-9917 M-F 9am - 5pm EST. Manufactured for: Wittman Pharma, Inc. Brooksville, FL 34604</p>	Age	Dose	adults and children 12 years and over	2-4 teaspoonfuls every 4 hours	children 6 years to under 12 years	1-2 teaspoonfuls every 4 hours	children 2 years to under 6 years	½-1 teaspoonfuls every 4 hours	under 2 years	ask a doctor
Age	Dose											
adults and children 12 years and over	2-4 teaspoonfuls every 4 hours											
children 6 years to under 12 years	1-2 teaspoonfuls every 4 hours											
children 2 years to under 6 years	½-1 teaspoonfuls every 4 hours											
under 2 years	ask a doctor											

CURATUSSIN

guaifenesin solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83335-101
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83335-101-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/01/2024	
2	NDC:83335-101-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/01/2024	
3	NDC:83335-101-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/01/2024	

Drug Facts	
Active Ingredient (in each 5 mL tsp) Guaifenesin 100 mg	Purpose Expectorant
Uses: ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive	
Warnings: Ask a doctor before use if you have ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic, such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
Stop use and ask doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions: ■ do not take more than 6 doses in any 24-hour period.	
Age	Dose
adults and children 12 years and over	2-4 teaspoonfuls every 4 hours

MADE IN THE USA NDC 83335-101-16

CuraTUSSIN

EXPECTORANT

GUAIFENESIN 100 mg / 5 mL

LOOSENS AND RELIEVES CHEST CONGESTION

Compare to the active ingredient in *Robitussin®

✓SUGAR FREE ✓ALCOHOL FREE
✓DYE FREE ✓GLUTEN FREE

Net Content: 16 fl oz (473 mL)
Grape Flavor



Drug Facts (continued)	
Directions: (continued) ■ do not take more than 6 doses in any 24-hour period.	
Age	Dose
children 6 years to under 12 years	1-2 teaspoonfuls every 4 hours
children 2 years to under 6 years	½-1 teaspoonfuls every 4 hours
under 2 years	ask a doctor
Other information ■ store at 15°-30°C (59°-86°F) ■ tamper evident, do not use if imprinted safety seal under cap is broken or missing	
Inactive Ingredients: Citric Acid, Flavor, Methylparaben, Monoammonium Glycyrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose	
Questions? Call 1-352-549-9917 M-F 9am - 5pm EST Manufactured for: Wittman Pharma, Inc. Brooksville, FL 34604	
*Robitussin® is a registered trademark of Wyeth Consumer Healthcare. Rev. 1/2024	
 N 3 83335 10116 2	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2024	

Labeler - Wittman Pharma, Inc. (830980947)

Establishment

Name	Address	ID/FEI	Business Operations
Wittman Pharma, Inc.		830980947	analysis(83335-101) , manufacture(83335-101) , label(83335-101)

Revised: 4/2024

Wittman Pharma, Inc.