

CURIST ACID RELIEF- famotidine 20 mg tablet
Little Pharma, Inc.

Curist Acid Relief (famotidine 20 mg tablets)

curistrelief.com

Important: Read all directions and warnings before use.

Tamper Evident: do not use if imprinted inner safety seal is torn or missing

Drug Facts

Active ingredient (in each tablet)

Famotidine USP, 20 mg

Purpose

Acid reducer

(CONTINUED ON BACK OF LABEL)

REV: 032-23-01

NDC 72559-032-23

Distributed by: Little Pharma, Inc.

New York, NY 10023 | Made in India

*Maximum strength famotidine tablet available over-the-counter (OTC) without a prescription

PEEL HERE

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn

- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warning before use
- store at 20°-25°C (68°-77°F)
- protect from moisture

hydroxypropyl cellulose, hypromellose, macrogol, magnesium stearate, microcrystalline cellulose, pre-gelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin

Questions or comments?

Call **1-844-243-1241** or email hi@curistrelief.com

REV: 032-23-01

STOP PEELING

curist

Acid Relief

Maximum Strength OTC*

Famotidine Tablets USP, 20 mg

Acid Reducer

Just One Tablet Prevents & Relieves Heartburn Due to Acid Indigestion

Maximum Strength Without a Prescription*

300 Tablets

curist

Acid Relief Maximum Strength OTC*

Famotidine Tablets USP, 20 mg
Acid Reducer

Just One Tablet Prevents & Relieves
Heartburn Due to Acid Indigestion
Maximum Strength Without a Prescription*

300 Tablets

curistrelief.com

Important: Read all directions and warnings before use.
Tamper Evident: Do not use if imprinted inner safety seal is torn or missing.

Drug Facts

Active ingredient Purpose (in each tablet)
Famotidine USP, 20 mg...Acid reducer


(CONTINUED ON BACK OF LABEL)

REV: 032-23-01

NDC: 72559-032-23

Distributed by: Little Pharma, Inc.
New York, NY 10023 | Made in India

*Maximum strength famotidine tablet available over-the-counter (OTC) without a prescription.



PEEL HERE 3 172559 03223 10

Min 0.1

Drug Facts (continued)

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

HINGE

Drug Facts (continued)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to prevent symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before eating food or drinking beverages** that cause heartburn
 - do not use more than 2 tablets in 24 hours
 - children under 12 years: ask a doctor

Other information

- read the directions and warning before use
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

hydroxypropyl cellulose, hypromellose, macrogol, magnesium stearate, microcrystalline cellulose, pre-gelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin

Questions or comments?

Call **1-844-243-1241** or email hi@curistrelief.com

REV: 032-23-01

STOP PEELING

CURIST ACID RELIEF

famotidine 20 mg tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:72559-032

Route of Administration

ORAL

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)		FAMOTIDINE	20 mg	
Inactive Ingredients				
Ingredient Name		Strength		
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
TRIACETIN (UNII: XHX3C3X673)				
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HF6D95)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
HYDROXYPROPYL CELLULOSE (110000 WAMW) (UNII: 5Y0974F5PW)				
STARCH, CORN (UNII: O8232NY3SJ)				
Product Characteristics				
Color	white (white to off-white)	Score	no score	
Shape	ROUND	Size	6mm	
Flavor		Imprint Code	V;15	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72559-032-23	300 in 1 BOTTLE; Type 0: Not a Combination Product	01/31/2024	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA215822	01/31/2024		

Labeler - Little Pharma, Inc. (074328189)

Revised: 1/2024

Little Pharma, Inc.