

**ARTHRITIS PAIN RELIEF- acetaminophen tablet, extended release
KROGER COMPANY**

Kroger Acetaminophen Extended Release Tablets 650 mg (Arthritis Pain)

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.

See overdose warning

adults	<ul style="list-style-type: none">• take 2 caplets every 8 hours with water• swallow whole; do not crush, chew, split, or dissolve• do not take more than 6 caplets in 24 hours• do not use for more than 10 days unless directed by a doctor
under 18 years of age	<ul style="list-style-type: none">• ask a doctor

Other information

- store between 20-25°C (68-77°F)
- The FDA approved dissolution methods differ from USP

Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? Call **1-800-632-6900**



NDC 41226-699-01

Arthritis Pain Relief Acetaminophen Extended-Release

Tablets USP, 650 mg

LASTS UP TO 8 HOURS
FOR THE TEMPORARY RELIEF OF
MINOR ARTHRITIS PAIN

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

PAIN RELIEVER/FEVER REDUCER

100 CAPLETS*

*CAPSULE-SHAPED BI-LAYER TABLETS

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER
THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 650 mg Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - premenstrual and menstrual cramps
 - the common cold
 - headache
 - toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- skin itching
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions Do not take more than directed. See overdose warning.
adults:

- take 2 caplets every 8 hours with water
- swallow whole, do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years of age adults: ask a doctor

Questions or comments? Call 1-800-632-6900

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

MADE IN INDIA

YOUR PHARMACIST RECOMMENDED

6991 0324

LOT:

EXP:

Varnish Omit Area

C6011AR3

Varnish/Print Omit Area

NDC 41226-699-01



COMPARE TO THE ACTIVE INGREDIENT
IN TYLENOL® 8HR ARTHRITIS PAIN

Arthritis Pain Relief
Acetaminophen Extended-Release


Tablets USP, 650 mg LASTS UP TO 8 HOURS • FOR THE TEMPORARY RELIEF OF MINOR ARTHRITIS PAIN



**PAIN RELIEVER/
FEVER REDUCER**

100 CAPLETS*

*CAPSULE-SHAPED 2-LAYER TABLETS



ACTUAL SIZE

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

Drug Facts

RETAIN CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION

Active ingredient (in each caplet)
Acetaminophen 650 mg.....Pain reliever/
fever reducer

Purpose

Uses
temporarily relieves minor aches and pains due to:
■ minor pain of arthritis ■ muscular aches
■ backache ■ premenstrual and menstrual cramps
■ the common cold ■ headache ■ toothache
temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen.
Severe liver damage may occur if you take:
■ more than 6 caplets in 24 hours, which is the
maximum daily amount

Drug Facts (continued)

■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using
this product
Allergy alert: Acetaminophen may cause severe skin
reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical
help right away.
Do not use
■ with any other drug containing acetaminophen
(prescription or nonprescription). If you are not sure
whether a drug contains acetaminophen, ask a
doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the
inactive ingredients in this product
Ask a doctor before use if you have liver disease

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin
Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health
professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical
help or contact a Poison Control Center right away
(1-800-222-1222). Quick medical attention is critical
for adults as well as for children even if you do not
notice any signs or symptoms.

Drug Facts (continued)

Directions
Do not take more than directed.
See overdose warning
adults
■ take 2 caplets every 8 hours with
water
■ swallow whole; do not crush, chew,
split, or dissolve
■ do not take more than 6 caplets in
24 hours
■ do not use for more than 10 days
unless directed by a doctor
**under 18
years of age**
ask a doctor
Other information
■ store between 20-25°C (68-77°F)

Drug Facts (continued)

■ The FDA approved dissolution methods differ from
USP
Inactive Ingredients
carnauba wax, hydroxyethyl cellulose, hypromellose,
magnesium stearate, microcrystalline cellulose,
povidone, pregelatinized starch, sodium starch
glycolate, titanium dioxide, triacetin
Questions or comments?
Call 1-800-632-6900

Contains No Aspirin
Tylenol® is a registered trademark of Kenvue, Inc.
Kenvue, Inc. is not affiliated with The Kroger Co.
or this product.
DISTRIBUTED BY THE KROGER CO.
CINCINNATI, OHIO 45202
MADE IN INDIA
For More Product Information, Scan UPC Using
Your Kroger App or Call 800-632-6900
OUR PHARMACIST RECOMMENDED
699R 0324

Varnish Omit Area

Varnish Omit Area

LOT:
EXP:
Varnish/Print
Omit Area



Our Brands. Our Guarantee.
Love It or Your Money Back.
www.kroger.com

THANKS TO OUR PARTNERS AND THE SUPPORT OF THE
COMMUNITY, WE'VE MADE THE KROGER STORE THE
BEST PLACE TO SHOP FOR THE BEST VALUE AND SERVICE.

ARTHRITIS PAIN RELIEF
acetaminophen tablet, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-699
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY35J)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
HYDROXYETHYL CELLULOSE (140 MPA.S AT 5%) (UNII: 8136Y38GY5)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white (White to off white)	Score	no score
Shape	CAPSULE (Capsule-Shaped Tablet)	Size	19mm
Flavor		Imprint Code	71
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-699-01	1 in 1 CARTON	04/15/2024	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215486	04/15/2024	

Labeler - KROGER COMPANY (006999528)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(41226-699)

Revised: 4/2024

KROGER COMPANY