

**CHILDRENS PAIN RELIEVER AND FEVER REDUCER GRAPE-
acetaminophen suspension
DeMoulas Market Basket**

Children's Pain & Fever, Grape

Drug Facts

Active ingredient

(in each 5 mL = 1 teaspoonful)
Acetaminophen 160 mg

Purpose

Acetaminophen Pain reliever/fever reducer

Uses temporarily • reduces fever

- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning

This product contains acetaminophen.

Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

Directions

- this product does not contain directions or complete warnings for adult use.
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

| Weight (lb) | Age (yr) | Dose (mL or tsp) * |
|-------------|---------------|--------------------|
| under 24 | under 2 years | ask a doctor |
| 24-35 | 2-3 years | 5 mL (1 tsp) |
| 36-47 | 4-5 years | 7.5 mL (1½ tsp) |
| 48-59 | 6-8 years | 10 mL (2 tsp) |
| 60-71 | 9-10 years | 12.5 mL (2½ tsp) |
| 72-95 | 11 years | 15 mL (3 tsp) |

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL (1 tsp) contains: sodium 3 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- **Keep carton for full directions for use.**

Inactive ingredients

anhydrous citric acid, butylparaben, D&C Red # 33, FD&C Blue #1, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**

CHILDRENS PAIN RELIEVER & FEVER REDUCER, Grape - Product Labeling

MARKET BASKET®
"MORE FOR YOUR DOLLAR"

**Children's
Pain &
Fever**

Compare to the Active
Ingredients in Children's
Tylenol® Oral Suspension

**Acetaminophen
Oral Suspension**

Grape Flavor

4 FL OZ (118 mL)

**DISTRIBUTED BY
DEMOULAS SUPERMARKETS INC.
875 East Street
Tewksbury, MA 01876**

DO NOT USE IF PRINTED NECK BAND IS BROKEN OR MISSING

This product is not manufactured or distributed by McNeil Consumer Healthcare,
distributor of Children's Tylenol®
Oral Suspension.

Carton

DO NOT USE IF PRINTED NECK BAND IS BROKEN OR MISSING

Drug Facts

Active Ingredient Purpose
 Acetaminophen 160 mg Pain reliever/fever reducer

Uses Temporarily relieves fever
 relieves minor aches and pains due to:
 the common cold flu headache sore throat
 toothache

Warnings

Liver warning
 This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount

with other drugs containing acetaminophen
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash
 If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning
 If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use
 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease
 Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if
 pain gets worse or lasts more than 3 days
 fever gets worse or lasts more than 3 days
 new symptoms occur
 redness or swelling is present
 These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning
 Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Outok medical attention is

Drug Facts (continued)

critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 This product does not contain directions or complete warnings for adult use.

- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- If needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor.

| Weight (lb) | Age (yr) | Dose (mL or tsp)* |
|-------------|---------------|-------------------|
| under 24 | under 2 years | ask a doctor |
| 24-35 | 2-3 years | 5 mL (1 tsp) |
| 36-47 | 4-5 years | 7.5 mL (1½ tsp) |
| 48-59 | 6-8 years | 10 mL (2 tsp) |
| 60-71 | 9-10 years | 12.5 mL (2½ tsp) |
| 72-85 | 11 years | 15 mL (3 tsp) |

*or as directed by a doctor
 Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information
 each 5 mL (1 tsp) contains: sodium 3 mg
 store between 20-25°C (68-77°F)
 do not refrigerate
 Keep carton for full directions for use.

Inactive ingredients anhydrous citric acid, butylparaben, D&C Red # 33, FD&C Blue # 1, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-6944

*This product is not manufactured or distributed by Medical Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

DISTRIBUTED BY: DEMIGULAS SUPERMARKETS INC. 875 EAST STREET, TOWNSHIRE, MA 01876

@marketbasket
 Please visit us at shopmarketbasket.com



Grape Flavor
 4 FL OZ (118 mL)
 160 mg per 5 mL



Grape Flavor
 4 FL OZ (118 mL)
 160 mg per 5 mL



0 49705 73958 7

Bottle



DO NOT USE IF PRINTED NECK BAND IS BROKEN OR MISSING
 Active Ingredient (in each 5 mL = 1 teaspoonful) Purpose
 Acetaminophen 160 mg Pain reliever/fever reducer

Uses Temporarily relieves fever relieves minor aches and pains due to:
 the common cold flu headache sore throat toothache
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen. Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash. If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If your child is allergic to acetaminophen or any of the inactive ingredients in this product, do not use. If your child has liver disease, ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product do not exceed recommended dose (see overdose warning). Stop use and ask a doctor if pain gets worse or lasts more than 3 days, fever gets worse or lasts more than 3 days, new symptoms occur, redness or swelling is present. These could be signs of a serious condition. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Outok medical help or contact a Poison Control Center right away. If you do not notice any signs or symptoms directed by a doctor. Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device. Other information: each 5 mL (1 tsp) contains: sodium 3 mg. Store between 20-25°C (68-77°F). Do not refrigerate. Keep carton for full directions for use. Inactive ingredients: anhydrous citric acid, butylparaben, D&C Red # 33, FD&C Blue # 1, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

Directions This product does not contain directions or complete warnings for adult use. Shake well before using. mL = milliliter; tsp = teaspoonful. Find right dose on chart below. If possible, use weight to dose; otherwise, use age. If needed, repeat dose every 4 hours while symptoms last. Do not give more than 5 times in 24 hours. Do not give for more than 5 days unless directed by a doctor.

| Weight (lb) | Age (yr) | Dose (mL or tsp)* |
|-------------|---------------|-------------------|
| under 24 | under 2 years | ask a doctor |
| 24-35 | 2-3 years | 5 mL (1 tsp) |
| 36-47 | 4-5 years | 7.5 mL (1½ tsp) |
| 48-59 | 6-8 years | 10 mL (2 tsp) |
| 60-71 | 9-10 years | 12.5 mL (2½ tsp) |
| 72-85 | 11 years | 15 mL (3 tsp) |

*or as directed by a doctor
 Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device. Other information: each 5 mL (1 tsp) contains: sodium 3 mg. Store between 20-25°C (68-77°F). Do not refrigerate. Keep carton for full directions for use. Inactive ingredients: anhydrous citric acid, butylparaben, D&C Red # 33, FD&C Blue # 1, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

LOT: 0649BMP7 CB1-877-798-6944
 DISTRIBUTED BY: DEMIGULAS SUPERMARKETS INC. 875 EAST STREET, TOWNSHIRE, MA 01876
 EXP: LR-281

CHILDRENS PAIN RELIEVER AND FEVER REDUCER GRAPE

acetaminophen suspension

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:53942-526 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|--------------------------|-----------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 160 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|-----------------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| BUTYLPARABEN (UNII: 3QPI1U3FV8) | |
| D&C RED NO. 33 (UNII: 9DBA0SBB0L) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SORBITOL SOLUTION (UNII: 8KW3E207O2) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|--|
| Color | purple | Score | |
| Shape | | Size | |
| Flavor | GRAPE | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|---|-----------------------------|---------------------------|
| 1 | NDC:53942-526-24 | 1 in 1 CARTON | 01/01/2026 | |
| 1 | | 118 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013 | 01/01/2026 | |

Labeler - DeMoulas Market Basket (007869647)

Registrant - AptaPharma Inc. (790523323)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------|---------|-----------|------------------------|
| AptaPharma Inc. | | 790523323 | manufacture(53942-526) |

Revised: 5/2026

DeMoulas Market Basket