

DR SABHARWALS- lidocaine 4% gel patch patch
Dr. Sabharwal's Wound Care

Lidocaine 4%

Temporarily Relieves Minor Pain

Topical Anesthetic

For external use only

- More than one patch on your body at a time
- On cut, irritated swollen skin
- On puncture wounds
- For more than one week without consulting a doctor
- At the same time as other topical analgesics

- Use only as directed.
- Read and follow all directions and warnings on this label.
- Rare case of serious burns have been reported with products of this type.
- Do not apply to wounds or damaged, broken or irritated skin.
- Do not allow contact with the eyes and mucous membranes.
- Do not bandage tightly or apply local heat (such as heating pads) to the area of use.
- Do not use at the same time as other topical analgesic.
- Dispose of used patch in manner that always keeps product away from children and pets.
- Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch
- A transient burning sensation may occur upon application but generally disappears in several days.
- If a severe burning sensation occurs, discontinue use immediately.
- Discontinue use at least 1 hour before a bath or shower.
- Do not use immediately after a bath or shower.

Condition worsens

Redness is present.

Irritation develops.

Symptoms persist for more than 7 days or clear up and occur again within a few days.

You experience signs of skin injury, such as pain, swelling or blistering where the product was applied.

ask a health professional before use.

If swallowed, get medical help or contact a Poison Control Center right away.

Adult and children over 12 years

Clean and dry affected area.

Remove backing from patch by firmly grasping both ends and gently pulling until back separates in middle.

Carefully remove smaller portion of backing from the patch and apply exposed portion of patch to affected area.

Use 1 patch for up to 12 hours.

Children 12 years or younger:
ask a doctor

Store at room temperature 20-25 °C (68-77 °F)

Alpha Tocopherol Acetate, Aluminium Glycinate, Aluminium Hydroxide, Aluminium Sulphate, Borax, Carbomer, Colloidal Silicon Dioxide, DMDM Hydantoin, Glycerine, Kaolin, Polyacrylic Acid, Polyvinyl Alcohol, Polyvinylpyrrolidone, Propylene Glycol, Purified Water, Sodium Carboxymethyl Cellulose, Sodium Ethylenediaminetetraacetic Acid, Sodium Polyacrylate, Sorbitol, Sorbitan Monooleate, Tartaric Acid & Titanium Dioxide.

Questions? Call +91-718-739-5680


Topical Patch for External Use Only

Dr Sabharwal's
Lidocaine 4% Gel Patch

NDC 71318-114-15

Dr Sabharwal's®
Lidocaine 4% Gel Patch

Dr Sabharwal's®
Lidocaine 4% Gel Patch



MAXIMUM STRENGTH

for temporary relief of pain:

- Back
- Neck
- Shoulders
- Knee & Elbow

Pain Relieving Patch

Dr Sabharwal's®
Lidocaine 4% Gel Patch

Drug Facts	
Active Ingredient Lidocaine 4%.....	Purpose Topical anesthetic.
Use temporarily relieves minor pain.	
Warnings: For external use only.	
Do not use: More than one patch on your body at a time. On cut, irritated, swollen skin. With puncture wounds. For more than one week without consulting a doctor. At the same time as other topical anesthetics.	
When using this product: Use only as directed. Read and follow all directions and warnings on this label. Where case of serious burns have been reported with products of this type. Do not apply to wounds or damaged, broken or irritated skin. Do not allow contact with the eyes and mucous membranes. Do not bandage tightly or apply heat (such as heating pads) to the area of use. Do not use at the same time as other topical anesthetics. Dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.	
Stop use and ask a doctor if: Condition worsens. Itchiness is present. Irritation develops. Symptoms persist for more than 7 days or come up and occur again within a few days. You experience signs of skin injury, such as pain, swelling or blistering where the product was applied.	
<small>*This product is not manufactured or distributed by Chiltern, Inc., a division of Allergan® (Pain-Free Strength Lidocaine Patch).</small>	

Adult and children over 12 years

■ Clean and dry affected area. ■ Remove backing from patch by firmly grasping both ends and gently pulling until back separates in middle. ■ Carefully remove smaller portion of backing from the patch and apply exposed portion of patch to affected area. ■ Use 1 patch for up to 12 hours.


Children 12 years or younger: ask a doctor.

Other Information: Store at room temperature 20-25°C (68-77°F).

Inactive Ingredients: Alpha Tocopherol Acetate, Aluminium Glycinate, Aluminium Hydroxide, Aluminium Sulphate, Borax, Carbomer, Colloidal Silicon Dioxide, DMDM Hydantoin, Glycerine, Kaolin, Polyacrylic Acid, Polyvinyl Alcohol, Polyvinylpyrrolidone, Propylene Glycol, Purified Water, Sodium Carboxymethyl Cellulose, Sodium Ethylenediaminetetraacetic Acid, Sodium Polyacrylate, Sorbitol, Sorbitan Monooleate, Tartaric Acid & Titanium Dioxide.

Questions? Call +91-905-677-0088

Batch No.:
Mfg. Date:
Exp. Date:



01 681923 72242023
085-4232 Rev. 03

MAXIMUM STRENGTH

TOPICAL ANESTHETIC

FAST ACTING

Quantity: 15 N Patch of Size 10cm x 14cm, Individually Packed

Product Code: 4202

Dr. Sabharwal's Wound Care

Mfg. In India by:
Dr. Sabharwal's Wound Care

137 Baramwala, Baroliwala (H.P.), India - 174 103
Manager QA Customer Care Cell
Telefax: +91 1722640018 Cell: +91 90567 70088
E-mail: buyers@drsabharwal.in
Mfg. Lic. No.: MNB/06/348

DR SABHARWALS

lidocaine 4% gel patch patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71318-114
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	224 mg

Inactive Ingredients

Ingredient Name	Strength
CARBOMER 940 (UNII: 4Q93RCW27E)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
DMDM HYDANTOIN (UNII: BYR0546TOW)	
SODIUM POLYACRYLATE (8000 MW) (UNII: 285CYO341L)	
POLYVINYL ALCOHOL (100000 MW) (UNII: 949E52Z6MY)	
SORBITOL (UNII: 506T60A25R)	
POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALPHA-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
TARTARIC ACID (UNII: W4888I119H)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
KAOLIN (UNII: 24H4NWX5CO)	

Product Characteristics

Color	white	Score	
Shape	RECTANGLE	Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71318-114-15	15 in 1 CARTON	01/30/2024	
1		1 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	01/30/2024	

Labeler - Dr. Sabharwal's Wound Care (862184668)

Registrant - Dr. Sabharwal's Wound Care (862184668)

Establishment

Name	Address	ID/FEI	Business Operations
Dr. Sabharwal's Wound Care		862184668	manufacture(71318-114)

Revised: 1/2024

Dr. Sabharwal's Wound Care