

**TYLENOL EXTRA STRENGTH- acetaminophen tablet, film coated**  
**Kenvue Brands LLC**

-----  
**Tylenol<sup>®</sup> Extra Strength**

***Drug Facts***

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 caplets every 6 hours while symptoms last</li><li>▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor</li><li>▪ do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

**Other information**

- store between 20-25°C (68-77°F)
- **do not use if carton is opened. Do not use if foil inner seal imprinted with "TYLENOL" is broken or missing**

**Inactive ingredients**

carnauba wax <sup>1</sup>, corn starch <sup>1</sup>, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, modified starch <sup>1</sup>, polyethylene glycol <sup>1</sup>, powdered cellulose,

pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide

---

1 contains one or more of these ingredients

**Questions or comments?**

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

**PRINCIPAL DISPLAY PANEL**

NDC 50580-937-07

TYLENOL<sup>®</sup>  
FOR ADULTS

Acetaminophen  
Pain Reliever  
Fever Reducer

Extra Strength

Actual Size

100 Caplets  
500 mg each



Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	white (RED PRINT)	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	TYLENOL;500
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-937-01	12 in 1 PACKAGE	08/19/2019	
1		10 in 1 VIAL; Type 0: Not a Combination Product		
2	NDC:50580-937-02	12 in 1 PACKAGE	09/16/2019	
2		10 in 1 VIAL; Type 0: Not a Combination Product		
3	NDC:50580-937-03	2 in 1 POUCH; Type 0: Not a Combination Product	07/31/2020	10/31/2022
4	NDC:50580-937-04	3 in 1 CARTON	07/31/2020	11/08/2022
4		2 in 1 POUCH; Type 0: Not a Combination Product		
5	NDC:50580-937-05	50 in 1 CARTON	07/31/2020	10/31/2022
5		2 in 1 POUCH; Type 0: Not a Combination Product		

6	NDC:50580-937-20	50 in 1 CARTON	07/31/2020	10/31/2022
6		2 in 1 POUCH; Type 0: Not a Combination Product		
7	NDC:50580-937-06	1 in 1 CARTON	08/31/2020	
7		24 in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:50580-937-10	1 in 1 CARTON	08/31/2020	
8		50 in 1 BOTTLE; Type 0: Not a Combination Product		
9	NDC:50580-937-07	1 in 1 CARTON	08/31/2020	
9		100 in 1 BOTTLE; Type 0: Not a Combination Product		
10	NDC:50580-937-15	1 in 1 CARTON	08/31/2020	
10		225 in 1 BOTTLE; Type 0: Not a Combination Product		
11	NDC:50580-937-19	325 in 1 BOTTLE; Type 0: Not a Combination Product	08/31/2020	
12	NDC:50580-937-66	2 in 1 CARTON	05/01/2024	
12		100 in 1 BOTTLE; Type 0: Not a Combination Product		
13	NDC:50580-937-23	3 in 1 CARTON	01/15/2026	
13		2 in 1 POUCH; Type 0: Not a Combination Product		
14	NDC:50580-937-25	2500 in 1 CASE	12/28/2025	
14		2 in 1 POUCH; Type 0: Not a Combination Product		
15	NDC:50580-937-50	50 in 1 CARTON	12/28/2025	
15		2 in 1 POUCH; Type 0: Not a Combination Product		
16	NDC:50580-937-51	50 in 1 CARTON	12/28/2025	
16		2 in 1 POUCH; Type 0: Not a Combination Product		
17	NDC:50580-937-37	320 in 1 BOTTLE; Type 0: Not a Combination Product	03/31/2026	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/19/2019	

**Labeler** - Kenvue Brands LLC (118772437)