

NAPROXEN SODIUM- naproxen sodium tablet
CVS

Naproxen Sodium TABLETS, 220 mg (NSAID)**
Pain reliever/fever reducer
GLUTEN FREE
Not for households with young children
200 TABLETS

Active ingredient (in each tablet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)**

**nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis muscular aches backache menstrual cramps
- headache toothache the common cold
- temporarily reduces fever

Allergy alert:

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives facial swelling asthma (wheezing)
- shock skin reddening rash blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, livercirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke: chest pain trouble breathing
- weakness in one part or side of body
- slurred speech leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing

- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years

- ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive Ingredients

FD&C blue#2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone k-30, sodium starch glycolate, stearic acid, titanium dioxide

Questions or Comments?

call **1-877-770-3183** Mon - Fri 8:00 AM EST to 5:00 PM PST

Principal Display Panel

EASY OPEN ARTHRITIS CAP



STRENGTH TO LAST 12 HOURS

actual size

NAPROXEN SODIUM
Tablets USP, 220 mg
Pain Reliever/
Fever Reducer (NSAID)**

Compare to Aleve® active ingredient!

200 TABLETS PACKAGE NOT CHILD-RESISTANT

NDC 51316-761-400

IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE.
TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Drug Facts
Active ingredient (in each tablet)
Naproxen sodium 220 mg. Pain reliever/fever reducer (naproxen 200 mg) (NSAID)**
**nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:
 minor pain of arthritis muscular aches
 backache menstrual cramps
 headache toothache
 the common cold
 temporarily reduces fever

Warnings
Alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 hives facial swelling rash blisters
 shock skin redness asthma (wheezing)
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleedings. The chance is higher if you:
 are age 60 or older
 have had stomach ulcers or bleeding problems
 take a blood thinning (anticoagulant) or steroid drug
 take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 have 3 or more alcoholic drinks every day while using this product
 take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
 if you have ever had an allergic reaction to any other pain reliever/fever reducer
 right before or after heart surgery

Contains an ingredient made from a grain containing gluten (wheat, barley, or rye)
 This product is not manufactured or distributed by Bayer HealthCare, LLC, distributor of Aleve®, Aleve® is a registered trademark of Bayer HealthCare, LLC.
 Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2025 CVS/Pharmacy. CVS.com
 Made in India V-38638

Lot # 20250205774
 700205020318
 Exp. 08/2026

(CONTINUED ON BACK OF LABEL)
 #879445
 1504282028513

label,
re.

Inside (adhesive side)

Drug Facts (continued)

Ask a doctor before use if:
 you have a history of stomach problems, such as heartburn
 you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 you are taking a diuretic
 you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are:
 under a doctor's care for any serious condition
 taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
 taking any other drug

When using this product:
 take with food or milk if stomach upset occurs

Stop use and ask a doctor if:
 you experience any of the following signs of stomach bleeding:
 red or black stools
 vomit blood or have bloody or black stools
 you have stomach pain that does not get better
 you have symptoms of heart problems or stroke:
 chest pain trouble breathing weakness in one part or side of body dazed speech leg swelling
 pain gets worse or lasts more than 10 days
 fever gets worse or lasts more than 3 days
 you have difficulty swallowing it feels like the pill is stuck in your throat
 redness or swelling is present in the painful area
 any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 do not take more than directed the smallest effective dose should be used drink a full glass of water with each dose
 adults and children 12 years and older:
 take 1 tablet every 8 to 12 hours with symptoms but do not exceed 2 tablets within the first hour
 do not exceed 3 tablets in a 24-hour period
 children under 12 years:
 ask a doctor

Other information
 each tablet contains: sodium 20 mg
 store at 20° - 25°C (68° - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C Blue #2 aluminum lake, hypromellose 2910, maleic anhydride, microcrystalline cellulose, polyethylene glycol, polydioxane, hydroxypropyl methylcellulose, stearic acid, titanium dioxide

Questions or comments?
 call 1-877-776-3183 Mon - Fri 8:00 AM EST to 5:00 PM PST

NAPROXEN SODIUM

naproxen sodium tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-761
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII: 57Y76R9ATQ)	NAPROXEN SODIUM	220 mg	

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K30 (UNII: U725QWY32X)	

Product Characteristics

Color	blue ((Light Blue))	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	220
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-761-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	05/25/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091353	05/25/2024	

Labeler - CVS (062312574)

Revised: 12/2025

CVS