# BUSCAPINA- acetaminophen, caffeine, pyrilamine maleate tablet, coated OPMX LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### **Bus capina**

### **Drug Facts**

Buscapina

### **Active Ingredients (in each caplet)**

Acetaminophen 500mg, Caffeine 60mg, and Pyrilamine maleate 15mg

### **Purpose**

Pain Reliever, Diuretic, Antihistamine, Multi-Symptom Relief

#### Uses

- For the temporary relief of these symptoms associated with menstrual periods
- Cramps
- Bloating
- Water- weight gain
- Headache
- Backache
- Muscle aches
- Fatigue

### **Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 tablets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy warning: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- Liver disease
- Glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

### Ask a doctor or pharmacist before use if you are

- Taking the blood thinning drug warfarin
- Taking sedatives or tranquilizers

### When using this product

- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- limit the use of caffeine-containing medications, foods or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

### Stop use and ask a doctor if

- new symptoms appear
- redness or swelling is present
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days

#### If pregnant or breast-feeding

Ask a health professional before use.

#### Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- Do not exceed recommended dosage.
- Adults and children 12 years and over: Take 2 tablets with water. Repeat every 6 hours, as needed. Do not exceed 6 tablets per day
- Children under 12 years: Consult a doctor

### Other Information

• Store at room temperature

• TAMPER EVIDENT; DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

### **Inactive ingredients**

- Colloidal silicon dioxide
- Hypromellose
- Magnesium stearate
- Microcrystalline cellulose
- Polyethylene glycol
- Polyvinyl pyrrolidone
- Pregelatinized starch
- Titanium dioxide
- Talc

### Questions or comments?

Call toll free 1-844-832-1138 Monday through Friday 9AM - 5PM or www.healthlifeofusa.com

Bus capina Acetaminophen 500mg Caffeine 60mg Pyrilamine maleate 15mg 24 Caplets Distributed By: OPMX San Diego, CA, 92154



### **BUSCAPINA**

acetaminophen, caffeine, pyrilamine maleate tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69729-130
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	500 mg		
CAFFEINE (UNII: 3G6 A5W338E) (CAFFEINE - UNII:3G6 A5W338E)	CAFFEINE	60 mg		
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg		

Inactive Ingredients		
Ingredient Name	Strength	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)		
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)		
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)		
TALC (UNII: 7SEV7J4R1U)		
HYPROMELLOSES (UNII: 3NXW29V3WO)		
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)		
PO VIDO NE (UNII: FZ989GH94E)		

Product Characteristics			
Color	white	Score	no score
Shape	capsule	Size	17mm
Flavor		Imprint Code	
Contains			

ı	Packaging			
ı	# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ı	1 NDC:69729-130-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	0 1/0 5/20 18	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	0 1/0 5/20 18	

## **Labeler -** OPMX LLC (029918743)

Establishment				
Name	Address	ID/FEI	Business Operations	
Elysium Pharmaceuticals Limited		915664486	manufacture(69729-130)	

Revised: 1/2018 OPMX LLC