

AUREOMYCIN- chlortetracycline hydrochloride granule
Phibro Animal Health

Aureomycin® 90 Granular

Aureomycin® 90 Granular

Chlortetracycline
Type A Medicated Article

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Net Wt 50 LB (22.68 kg)

Active drug ingredient

Chlortetracycline calcium complex equivalent to 90 g chlortetracycline hydrochloride per lb.

Ingredients

Dried *Streptomyces aureofaciens* Fermentation Product and Calcium Sulfate.

For use in the manufacture of medicated animal feeds.
For use in dry feed only. Not for use in liquid medicated feeds.

Use directions

Mix sufficient Aureomycin 90 Granular Type A Medicated Article to supply desired concentration of chlortetracycline per ton with part of the feed ingredients to make a preblend. Add the remainder of the ingredients and mix thoroughly. For specific use levels, see **Indications for use**.

Mixing directions

Level desired grams per ton	Amount of medicated article per ton †
50	9 oz
100	1 lb 2 oz
200	2 lb 4 oz
400	4 lb 8 oz
500	5 lb 9 oz

† It is recommended that 1 pound 2 ounces of Aureomycin 90 Granular Type A Medicated Article be diluted with 2 pounds 14 ounces of one of the feed ingredients to form a 4 pound working premix. Use 2 pounds of the working premix to make a preblend (see Use directions) for a Type C feed containing 50 g chlortetracycline / ton of feed

Indications for use

Indications for use	Chlortetracycline mg per lb body wt per day
<p>Cattle Beef Cattle (over 700 lb): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.</p>	0.5
<p>Beef and Non-Lactating Dairy Cattle: As an aid in the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. For use in free-choice feeds. A medicated feed mill license is required when the free-choice feed is manufactured using a proprietary formula and/or specifications. Free-choice feed formulations must be FDA-approved.</p>	0.5-2.0
<p>Calves, Beef and Non-Lactating Dairy Cattle: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to chlortetracycline. Feed for not more than 5 days. The appropriate amount of Aureomycin-containing feed supplement may be mixed in the cattle's daily ration or administered as a top-dress. If the Aureomycin-containing feed supplement is administered as a top-dress, it must be spread uniformly on top of the ration and sufficient space must be provided so that all cattle can eat at the same time.</p>	10
<p>Swine Control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to chlortetracycline. Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Note: this drug level is equivalent to approximately 400 grams per ton, depending on feed consumption and body weight.) Feed for not more than 14 days.</p>	10
<p>Turkeys Control of complicating bacterial organisms associated with</p>	25

bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

Indications for use

Cattle

Growing Cattle (over 400 lb): For the reduction of the incidence of liver abscesses.

Beef Cattle and Dairy Replacement Heifers: Control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline.

Beef Cattle (under 700 lb): Control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline.

Sheep

Breeding Sheep: Reduction in the incidence of (vibriotic) abortions caused by *Campylobacter fetus* infection susceptible to chlortetracycline.

Indications for use

Swine

Reduction in the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E *Streptococci* susceptible to chlortetracycline.

Breeding Swine: Control of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by *Leptospira pomona* susceptible to chlortetracycline. Feed continuously for not more than 14 days.

Ducks

Control and treatment of fowl cholera caused by *Pasteurella multocida* susceptible to chlortetracycline. Feed in complete ration to provide from 8 to 28 mg per pound of body weight per day depending upon age and severity of disease. Feed for not more than 21 days.

Chickens

Control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

**mg per head
per day**

70

350

350

80

**In complete feed
Chlortetracycline g
per ton**

50-100

400

200-400

100-200

Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	200-400
Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. Feed for 5 days.	500
Turkeys Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	200
Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	400
Turkey Poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.	400
Indications for use Psittacine birds Warning: Psittacosis, avian chlamydiosis, or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals and man. Contact appropriate public health and regulatory officials. Caution: Aspergilliosis may occur following prolonged treatment.	mg per g feed
Treatment of psittacine birds (parrots, macaws, cockatoos) suspected or known to be infected with psittacosis caused by <i>Chlamydia psittaci</i> sensitive to chlortetracycline. Feed continuously for 45 days. Each bird should consume an amount of medicated feed equal to one-fifth of its body weight daily. During treatment, parrots, macaws, and cockatoos should be kept individually or in pairs in clean cages.	10

Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling. This drug is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to ducks or turkeys producing eggs for human consumption.

Storage

Store below 25°C (77°F), excursions permitted to 37°C (99°F)

Restricted Drug (California) - Use only as directed. Not for use in humans.

Keep out of reach of children.

Approved by FDA under NADA # 048-761

Questions/Comments? For technical assistance or to obtain a Safety Data Sheet(s), contact Phibro Animal Health Corporation at 1-888-403-0074 or ww.pahc.com. To report side effects, contact Phibro Animal Health Corporation at 1-888-475-7355. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Phibro, Phibro logo, Aureomycin and the A logo are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

Distributed by:

Phibro Animal Health Corporation

Teaneck, NJ 07666

176-9050-02


TAKE TIME



**OBSERVE LABEL
DIRECTIONS**

Take Time Statement

PRINCIPAL DISPLAY PANEL



Aureomycin 90 G Label

AUREOMYCIN

chlortetracycline hydrochloride granule

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66104-1005
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORTETRACYCLINE HYDROCHLORIDE (UNII: O1GX33ON8R) (CHLORTETRACYCLINE - UNII:WCK1KIQ23Q)	CHLORTETRACYCLINE HYDROCHLORIDE	90 g in 0.45 kg

Reference ID	Reference ID	Reference ID	Reference ID
244-04011	244-04011	244-04011	244-04011
244-04011	244-04011	244-04011	244-04011
244-04011	244-04011	244-04011	244-04011

Product Characteristics

Color	gray (gray to brown)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-1005-0	22.68 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA048761	01/01/2009	

Labeler - Phibro Animal Health (006989008)

Revised: 2/2026

Phibro Animal Health