

**BACSTOP- iodine solution
IBA**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

BAC-STOP SANITIZING TEAT DIP A HIGHLY EFFECTIVE PRE-MILKING AND POST-MILKING IODINE TEAT DIP WITH TRIPLE EMOLLIENTS CONTAINS GLYCERIN, LANOLIN AND ALOE VERA AS SKIN CONDITIONERS

ACTIVE INGREDIENT:

Nonylphenoxypolyethoxyethanol Iodine Complex.....2.50%
(Provides .5% minimum Titratable Iodine)

INERT INGREDIENTS

Skin Conditioners.....2.00%
Other Inerts.....95.50%
(Includes Buffering Agents and Surfactants)
Total.....100%
pH of 5.5 at time of manufacture

INDICATIONS AND USAGE PROVIDES

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash Teat Dip cups after each milking.

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. REMOVE VISIBLE SOILS FROM THE TEATS.
2. OBSERVE FOREMILK BY STRIPPING TWO OR THREE STREAMS OF MILK INTO A STRIP CUP.
3. DIP OR SPRAY THE COW'S TEATS FULL LENGTH WITH BAC-STOP TEAT DIP.
4. WAIT APPROXIMATELY 30 SECONDS.
5. REMOVE ALL TEAT DIP BY THOROUGHLY DRYING THE TEATS AND TEAT ORIFICES WITH A CLEAN, SINGLE SERVICE PAPER TOWEL(S).
6. ATTACH MILKING UNIT.

B. POST-MILKING (POST -DIPPING)

Immediately after milking, dip or spray all teats full length with BAC-STOP Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.

UNIVERSITY TESTED AND PASSED USING NATIONAL MASTITIS COUNCIL PROTOCOLS.

Ã,Ã Ã,Ã Ã,Ã Ã,Ã Ã,Ã Ã,Ã PROVEN EFFECTIVE IN REDUCING NEW

INTRAMAMMARY INFECTIONS.

PART OF THE COMPLETE IBA SANITATION PROGRAM

WARNING

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES . TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR CONTACT WITH CONCENTRATED PRODUCTS, FLUSH THOROUGHLY WITH FRESH,TEPID WATER, DISCARD CONTAMINATED CLOTHING AND FOOTWEAR OR WASH BEFORE REUSE. FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION

FOR MEDICAL EMERGENCIES CALL 1-800-424-9300



BAC-STOP

SANITIZING TEAT DIP

**A Highly Effective Pre-Milking and Post-Milking Iodine Teat Dip With Triple Emollients
Contains Glycerin, Lanolin and Aloe Vera as Skin Conditioners**

ACTIVE INGREDIENT:

Nonylphenoxypolyethoxyethanol-Iodine Complex..... 2.50%
(Provides .5% minimum Titratable Iodine)

INERT INGREDIENTS:

Skin Conditioners..... 2.00%
Other Inerts 95.50%
(Includes Buffering Agents and Surfactants)

TOTAL: 100.00%

pH of 5.5 at time of manufacture

This product contains highly effective levels of free iodine for fast kill of mastitis causing organisms.

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.**

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**



RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray the cow's teats full length with BAC-STOP Teat Dip.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats and teat orifices with a clean, single service paper towel(s).
6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

1. Immediately after milking, dip or spray all teats full length with BAC-STOP Teat Dip. **Ensure good coverage of all sides of the teats and teat orifices.**

**UNIVERSITY TESTED AND PASSED USING NATIONAL MASTITIS COUNCIL PROTOCOLS.
PROVEN EFFECTIVE IN REDUCING NEW INTRAMAMMARY INFECTIONS.**

WARNING

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING.

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE: LOT NO.: NET CONTENTS: 5 GALLONS IBA STOCK # 160959 600E

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 27 PROVIDENCE ROAD, MILLBURY, MA 01527

BACSTOP

iodine solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-0003
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	5 g in 1 L

Inactive Ingredients

Ingredient Name	Strength
NONOXYNOL-30 (UNII: JJX07DG188)	

NONOXYNOL-9 (UNII: 48Q180SH9T)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
POTASSIUM IODIDE (UNII: 1C4QK22F9J)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-0003-1	3.78 L in 1 DRUM		
2	NDC:29983-0003-4	18.9 L in 1 DRUM		
3	NDC:29983-0003-6	56.7 L in 1 DRUM		
4	NDC:29983-0003-7	113.4 L in 1 DRUM		
5	NDC:29983-0003-8	207.9 L in 1 DRUM		
6	NDC:29983-0003-9	1039.5 L in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/23/2010	

Labeler - IBA (019494160)