

**PERMAVAN EXTERNAL PATCH- permavan external patch patch**  
**Home Aide Diagnostics, Inc.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----

**PERMAVAN EXTERNAL PATCH**

This product may cause allergic reaction in some individuals.

**DRUG FACTS**

# PERMAVAN

EXTERNAL PATCH

## Drug Facts

### Active Ingredients

	Purpose
Trolamine salicylate 10% -----	Topical analgesic
Dextromethorphan Hydrobromide 4% -----	Sigma-1 Agonist
Lidocaine 4% -----	Topical analgesic

**Uses** For relief of muscle aches and pains as well as:

- ♦ Arthritis ♦ Backaches ♦ Strains ♦ Sprains

### WARNINGS

#### FOR EXTERNAL USE ONLY.

- ♦ Use only as directed
- ♦ Avoid contact with mucous membranes
- ♦ Do not apply to wounds, damaged or irritated skin
- ♦ Do not bandage or cover with any type of wrap except clothing
- ♦ Do not use with a heating pad or apply external heat
- ♦ Do not use in large quantities, particularly over raw surfaces or blistered areas

#### STOP USE AND ASK A DOCTOR IF:

- ♦ Condition worsens
- ♦ Severe skin irritation occurs
- ♦ Pain persists for more than seven days

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**If pregnant or breast-feeding,** ask a healthcare professional before use

- ♦ This product may cause allergic reaction in some individuals.

### DIRECTIONS

Store below 25 °C (77 °F). Avoid direct sunlight.

Clean and dry affected area

Remove patch from backing and apply to affected area

Use only one patch at a time and maximum of four patches daily

Leave patch on affected area for up to 8-hours

Do not use patches for longer than 5 consecutive days unless directed by a physician.

Children under 12 years of age should consult a physician prior to use.

### INACTIVE INGREDIENTS

Acrylic acid ,aluminum hydroxide .carmellose sodium  
2-ethylhexyl acrylate .glycerin . isopropyl myristate .methyl acrylate  
nonoxynol-30 .polyacrylate.polyacrylic acid .polysorbate 80  
sorbitan sesquioleate starch .talc .tartaric acid . titanium dioxide .water

PERMAVAN  
EXTERNAL PATCH

**PERMAVAN**  
EXTERNAL PATCH

# PERMAVAN

## EXTERNAL PATCH

Trolamine salicylate 10% and Lidocaine 4% Dextromethorphan Hydrobromide 4%

NDC#: 69379-0010-15

Qty 15 Patches

Lot#:  
Exp Date:



**Distributed by:**  
Home Aide Diagnostics, Inc.  
1072 S. Powerline Road  
Deerfield Beach, FL 33442  
1 (800) 915-0116  
www.homeaide.us  
Made in China



**Distributed by:**  
Home Aide Diagnostics, Inc.  
1072 S. Powerline Road  
Deerfield Beach, FL 33442  
1 (800) 915-0116  
www.homeaide.us



**PERMAVAN**  
EXTERNAL PATCH

**Drug Facts**

**Active ingredients**  
Trolamine salicylate 10%  
Dextromethorphan Hydrobromide 4%  
Lidocaine 4%

**Warnings**  
• Avoid contact with mucous membranes  
• Use only as directed  
• Do not apply to wounds, damaged or irritated skin  
• Do not bandage or cover with any type of wrap except clothing  
• Do not use with a heating pad or apply external heat  
• Do not use in large quantities, particularly over raw surfaces or irritated areas

**FOR EXTERNAL USE ONLY.**

**STOP USE AND ASK A DOCTOR IF:**  
• Severe skin irritation occurs  
• Condition worsens  
• Pain persists for more than seven days

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.  
If pregnant or breastfeeding, ask a healthcare professional before use

**Directions**  
• This product may cause allergic reaction in some individuals.  
Clean and dry affected area  
Shave below 25 °C (77 °F). Avoid direct sunlight  
Remove patch from backing and apply to affected area  
Use only one patch at a time and maximum of four patches daily  
Leave patch on affected area for up to 8 hours  
Do not use patches for longer than 5 consecutive days unless directed by a physician.  
Children under 12 years of age should consult a physician prior to use.

**INACTIVE INGREDIENTS**  
Acrylic acid, aluminum hydroxide, camphore, sodium  
Zinc-ethylacrylate glycerin, isopropyl myristate, methyl acrylate  
monomethyl polyacrylate, polyethylene glycol, polyethylene glycol  
sorbitan sesquiscorbate stearate, talc, tartaric acid, titanium dioxide, water

**Purpose**  
Topical Analgesic  
Topical Anesthetist  
Topical Antipruritic

**PERMAVAN**  
EXTERNAL PATCH

### PERMAVAN EXTERNAL PATCH

permavan external patch patch

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:69379-010
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>TROLAMINE SALICYLATE</b> (UNII: H8O404BHD) (SALICYLIC ACID - UNII:O414PZ4LPZ)	TROLAMINE SALICYLATE	10 g in 100 g
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	4 g in 100 g

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:69379-010-15	5 in 1 BOX	03/09/2015	
1		3 in 1 PACKAGE		
1		100 g in 1 PATCH; Type 3: Prefilled Biologic Delivery Device/System (syringe, patch, etc.)		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
unapproved drug other		03/09/2015	

**Labeler** - Home Aide Diagnostics, Inc. (783518983)

Revised: 3/2015

Home Aide Diagnostics, Inc.