

**REMAXAZON EXTERNAL PATCH- remaxazon external patch patch**  
**Home Aide Diagnostics, Inc.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

**REMAXAZON EXTERNAL PATCH**

This product may cause allergic reaction in some individuals.

**DRUG FACTS**

# REMAXAZON

EXTERNAL PATCH

## Drug Facts

Active Ingredients	Purpose
Glucosamine 5% -----	Hexosamine
Chondroitin Sulfate Sodium 3% -----	Joint supplement
Capsacin 0.0285% -----	External Analgesic
Lidocaine 4% -----	Topical analgesic

**Uses** For relief of muscle aches and pains as well as:

♦ Arthritis ♦ Backaches ♦ Strains ♦ Sprains

### WARNINGS

#### FOR EXTERNAL USE ONLY.

- ♦ Use only as directed
- ♦ Avoid contact with mucous membranes
- ♦ Do not apply to wounds, damaged or irritated skin
- ♦ Do not bandage or cover with any type of wrap except clothing
- ♦ Do not use with a heating pad or apply external heat
- ♦ Do not use in large quantities, particularly over raw surfaces or blistered areas

#### STOP USE AND ASK A DOCTOR IF:

- ♦ Condition worsens
- ♦ Severe skin irritation occurs
- ♦ Pain persists for more than seven days

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**If pregnant or breast-feeding,** ask a healthcare professional before use

- ♦ This product may cause allergic reaction in some individuals.

### DIRECTIONS

Store below 25 °C (77 °F). Avoid direct sunlight.

Clean and dry affected area

Remove patch from backing and apply to affected area

Use only one patch at a time and maximum of four patches daily

Leave patch on affected area for up to 8-hours

Do not use patches for longer than 5 consecutive days unless directed by a physician.

Children under 12 years of age should consult a physician prior to use.

### INACTIVE INGREDIENTS

Acrylic acid .aluminum hydroxide .carmellose sodium

2-ethylhexyl acrylate .glycerin . isopropyl myristate .methyl acrylate

nonoxynol-30 .polyacrylate.polyacrylic acid .polysorbate 80

rorbitan sesquioleate starch .talc .tartaric acid . titanium dioxide .water

# REMAXAZON

EXTERNAL PATCH

**REMAXAZON**  
EXTERNAL PATCH

# REMAXAZON

## EXTERNAL PATCH

Glucosamine 5% and Chondroitin Sulfate Sodium 3% Capsacin 0.0285 Lidocaine 4%

NDC#: 69379-0285-15

Qty 15 Patches



**Distributed by:**  
Home Aide Diagnostics, Inc.  
1072 S. Powerline Road  
Deerfield Beach, FL 33442  
1 (800) 915-0116  
www.homeaide.us  
Made in China

Lot#:  
Exp Date:

**Distributed by:**  
Home Aide Diagnostics, Inc.  
1072 S. Powerline Road  
Deerfield Beach, FL 33442  
1 (800) 915-0116  
www.homeaide.us



**REMAXAZON**  
EXTERNAL PATCH

**Drug Facts**

**Active Ingredients**

Glucosamine 5%  
Chondroitin Sulfate Sodium 3%  
Lidocaine 4%  
Capsacin 0.0285%  
Topical analgesic

**Warnings**

Use for relief of muscle aches and pains as well as:

- Arthritis
- Osteoarthritis
- Sprains
- Burns

**FOR EXTERNAL USE ONLY**

Use only as directed

- Avoid contact with mucous membranes
- Do not apply to wounds, damaged or irritated skin
- Do not bandage or cover with any type of wrap except clothing
- Do not use with a heating pad or apply external heat
- Do not use in large quantities, particularly over raw surfaces or blistered areas

**STOP USE AND ASK A DOCTOR IF:**

- Contact with mucous membranes occurs
- Severe skin irritation occurs
- Skin persists for more than seven days

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Prevent of breastfeeding.** Ask a healthcare professional before use

**DIRECTIONS**

Store below 25°C (77°F). Avoid direct sunlight

Clean and dry affected area

Remove patch from backing and apply to affected area

Use only one patch at a time and maximum of four patches daily

Leave patch on affected area for up to 8 hours

Do not use patches for longer than 3 consecutive days unless directed by a physician

Children under 12 years of age should consult a physician prior to use

**ACTIVE INGREDIENTS**

Active and inactive ingredients contained sodium

2-hydroxyethyl acrylate glycerin, styrene/acrylate methyl methacrylate

nonylphenol polyoxyethylamine polyethylene glycol dimethylsiloxane

octylphenol polyoxyethylamine and polyethylene glycol dimethylsiloxane

**REMAXAZON**  
EXTERNAL PATCH

### REMAXAZON EXTERNAL PATCH

remaxazon external patch patch

#### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:69379-285
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>CHONDROITIN SULFATE SODIUM (BOVINE)</b> (UNII: 8QTV3DTT8W) (CHONDROITIN SULFATE (BOVINE) - UNII:6IC1M3OG5Z)	CHONDROITIN SULFATE (BOVINE)	3 g in 100 g
<b>CAPSAICIN</b> (UNII: S07O44R1ZM) (CAPSAICIN - UNII:S07O44R1ZM)	CAPSAICIN	.0285 g in 100 g
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g
<b>GLUCOSAMINE</b> (UNII: N08U5BOQ1K) (GLUCOSAMINE - UNII:N08U5BOQ1K)	GLUCOSAMINE	5 g in 100 g

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69379-285-15	5 in 1 BOX	03/09/2015	
1		3 in 1 PACKAGE		
1		100 g in 1 PATCH; Type 3: Prefilled Biologic Delivery Device/System (syringe, patch, etc.)		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/09/2015	

**Labeler** - Home Aide Diagnostics, Inc. (783518983)