

EXTRA STRENGTH PAIN RELIEVER- acetaminophen tablet
INGLES MARKETS, INC.

II 201

Active Ingredients

Acetaminophen 500 mg

Purpose

Pain Reliever/Fever Reducer

Uses

- temporarily relieves minor aches and pains
- temporarily reduces fever

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets (4,000 mg of acetaminophen) in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptom occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

Adults and children 12 years and over	Take 1-2 tablets every 4-6 hours, as needed. Do not take more than 6 tablets in 24 hours, or as directed by a doctor
Children under 12 years	Ask a doctor

Other Information

- **TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.**
- store at 20°C-25°C (68°F-77°F)

Inactive Ingredients

povidone, sodium starch glycolate, starch, stearic acid.

Drug Facts (continued)

Directions • do not take more than directed

Adults and children 12 years of age and over	Take 1-2 tablets every 4-6 hours, as needed. Do not take more than 6 tablets in 24 hours, unless directed by a doctor.
Children under 12 years	Ask a doctor.

Other information

- **TAMPER EVIDENT:** Do not use if imprinted seal under cap is missing or broken.
- store at 20°C - 25°C (68°F-77°F)

Inactive ingredients: povidone, sodium starch glycolate, starch, stearic acid.

*This product is not manufactured or distributed by the owner of the registered trademark **TYLENOL®**, REV 201-0619

NON-ASPIRIN

Pain Reliever

EXTRA STRENGTH

ACETAMINOPHEN 500 mg

Laura Lynn™



Compare to Active Ingredient in Extra Strength Tylenol®*

EXTRA STRENGTH Pain Reliever

ACETAMINOPHEN 500 mg

NON-ASPIRIN

- pain reliever
- fever reducer



60 TABLETS



31
000000
00/00

Drug Facts

SAVE CARTON FOR COMPLETE DRUG FACTS

Active ingredient (in each tablet)

Purposes

Acetaminophen 500 mg Pain Reliever / Fever Reducer

Uses • temporarily relieves minor aches and pains • temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 6 tablets in 24 hours. Severe liver damage may occur if you take:

- more than 8 tablets (4,000 mg of acetaminophen) in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur • redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

DISTRIBUTED BY: Ingles, P.O. Box 6676, Asheville, NC 28816



EXTRA STRENGTH PAIN RELIEVER

acetaminophen tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:62936-0201

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white (WHITE)	Score	no score
Shape	ROUND (Round)	Size	12mm
Flavor		Imprint Code	M2A457344
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62936-0201-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2019	
2	NDC:62936-0201-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/01/2006	

Labeler - INGLES MARKETS, INC. (024418584)

Registrant - Geri-Care Pharmaceutical Corp (611196254)