

HURRICAINA TOPICAL ANESTHETIC- benzocaine gel
Beutlich Pharmaceuticals, LLC

HurriCaine Gel W/C Tubes

Active ingredient

Benzocaine 20%

Purpose

Oral Anesthetic

Uses

for the temporary relief of occasional minor irritation and pain, associated with:

- canker sores
- sore mouth and throat
- minor injury of the mouth and gums
- minor dental procedures
- minor irritation of the mouth and gums caused by dentures or orthodontic appliances

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or irritation, pain, or redness persists or worsens, see your dentist or doctor promptly.

Do not use

Do not use:

- if the tube has been punctured or previously opened
- for teething
- in children under 2 years of age

When using this product

- avoid contact with eyes

Keep out of reach of children.

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

Directions**Directions:**

Do not exceed recommended dosage.

- adults and children 2 years of age and older: apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.
- children under 12 years of age: should be supervised in the use of the product
- children under 2 years of age: do not use

Other information

- store at 20°-25° C (68°-77° F)

Inactive ingredients

- flavor, polyethylene glycol, sodium saccharin

Questions or comments?

1-800-238-8542

M-F: 8:00 a.m. - 4:30 p.m. ET

Principal Display Panel



TAMPER EVIDENT FEATURE: For your protection, this product is supplied in a sealed tube. Remove cap and strip off the end of the tube to open. Do not use if the tube has been punctured or previously opened. Please retain Drug Facts for future reference.

Drug Facts		Drug Facts (continued)	
Active ingredient Benzocaine 20%.....	PurposeOral Anesthetic	Warnings (continued)	
Uses for the temporary relief of occasional minor irritation and pain, associated with <ul style="list-style-type: none"> • canker sores • sore mouth and throat • minor injury of the mouth and gums • minor dental procedures • minor irritation of the mouth and gums caused by dentures or orthodontic appliances 		Do not use <ul style="list-style-type: none"> • for teething • in children under 2 years of age 	
Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: <ul style="list-style-type: none"> • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.		When using this product avoid contact with eyes. Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.		Directions do not exceed recommended dosage adults and children 2 years of age and older children under 12 years of age children under 2 years of age apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor. should be supervised in the use of the product do not use	
Other information store at 20-25 °C (68-77 °F)		Inactive ingredients flavor, polyethylene glycol, sodium saccharin	
Questions or comments? 1-800-238-8842 M-F: 8:00 a.m. - 4:30 p.m. ET			

WHAT YOUR DENTIST USES

NDC 0283-0871-75

HURRICAINÉ®

Topical Anesthetic GEL

20% Benzocaine Oral Anesthetic

ORIGINAL WILD CHERRY

Manufactured for:

Beutlich® Pharmaceuticals, LLC

Bunnell, FL

Made in the USA

1 Tube - NET WT. 0.18 oz. (5.25 g)

HURRICAINÉ TOPICAL ANESTHETIC

benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0283-0871
Route of Administration	ORAL, DENTAL, PERIODONTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	brown (amber)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0283-0871-12	12 in 1 CELLO PACK	08/15/1971	
1	NDC:0283-0871-75	1 in 1 BOX		
1	NDC:0283-0871-01	5.25 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	08/15/1971	

Labeler - Beutlich Pharmaceuticals, LLC (005209325)

Registrant - Beutlich Pharmaceuticals, LLC (005209325)

Establishment

Name	Address	ID/FEI	Business Operations
Dental Technologies, Inc.		148312838	manufacture(0283-0871)

