

**WELLY FIRST AID KIT- hydrocortisone, bacitracin zinc, neomycin sulfate,
polymyxin b sulfate, ibuprofen
Welly Health PBC**

Welly First Aid Kit

1% Hydrocortisone Cream

Drug Facts

Active ingredient

Hydrocortisone 1.0%

Purpose

Anti-itch

Uses

- For temporary relief of itching associated with minor skin irritations, inflammation, or rashes.
- Other uses of product should be only under the advice and supervision of a doctor.

Warnings

For external use only

Do not use

- in eyes
- for treatment of diaper rash
- for feminine itching

Stop use, ask a doctor

- if condition worsens or lasts more than 7 days, or clears up and occurs again within a few days
- with use of other hydrocortisone products

Keep out of reach of children. If ingested, contact a Poison Control Center right away

Directions

- apply to affected area not more than 3 to 4 times daily
- children under 2: ask a doctor

Inactive ingredients

-
emulsifying wax, ethanol, methylparaben, mineral oil, paraffin, petrolatum, propylparaben, purified water, white wax

Triple Antibiotic Ointment

Drug Facts

Active Ingredient (in each gram)

Bacitracin zinc 400 units

Neomycin sulfate (3.5mg Neomycin)

Polymyxin B sulfate 5000 units

Purpose

First Aid Antibiotics

Uses

First aid to help prevent infection in minor cuts, scrapes and burns

Warnings

For external use only

Do not use

- internally
- in eyes
- over large areas of the body or on puncture wounds, animal bites or serious burns
- for more than 1 week unless directed by a doctor
- if you are allergic to any of the ingredients

Stop use and ask a doctor if

- a rash or allergic reactions develops
- condition worsens or persists

Keep out of reach of children. If ingested, contact a Poison Control Center right away.

Directions

- clean affected area
- apply a small amount 1 to 3 times daily
- may cover with a sterile bandage

Inactive Ingredients

petrolatum

Pain Relief and Fever Reducer

Drug Facts

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- muscular aches
- backache
- toothache
- menstrual cramps
- headache
- the common cold
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- shock
- facial swelling
- asthma (wheezing)
- rash
- skin reddening
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn.
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding.
- Feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- slurred speech
- leg swelling
- trouble breathing
- weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breastfeeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- If pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NO USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see label for expiration date and lot number
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

1-833-BE-WELLY

First Aid Kit

LARGE FIRST AID KIT TO COVER ANY BUMP, BRUISE OR SCRAPE.

Everything you need for when the unbeaten path gets the best of you.

Flexible fabric bandages that can move with your every move.

Durable metal case keeps it all together.

Cleansing wipes, tape and non-stick pads to help you get back out there.

Triple Antibiotic ointment helps prevent infection.

KIT INCLUDES

48 COUNT - STANDARD FABRIC BANDAGES

30 COUNT - SMALL FABRIC BANDAGES

3 COUNT - THICK STRIP FLEX FABRIC BANDAGES (2"x4")

12 COUNT - WATERPROOF BANDAGES

1 COUNT - ROLL OF TAPE

5 COUNT - NON STICK PADS

10 COUNT - CLEANSING WIPES

10 COUNT - BUTTERFLY STRIPS

5 COUNT - TRIPLE ANTIBIOTIC PACKETS

5 COUNT - HYDROCORTISONE PACKETS

1 SET PACKETS (16 TABLETS) - IBUPROFEN

Manufactured for: Welly Health PBC

Minneapolis, MN 55402

1-833-BE-WELLY

Recycle me!

Welly™ www.GetWelly.com

Products of USA and China with globally sourced materials

Product not made with natural rubber latex

Sterile unless individual wrapper opened or damaged.

Patent Pending

Packaging

First Aid Kit Label

FIRST AID KIT

48 COUNT - STANDARD FABRIC BANDAGES
 30 COUNT - SMALL FABRIC BANDAGES
 12 COUNT - STANDARD WATERPROOF BANDAGES
 10 COUNT - BUTTERFLY STRIPS
 5 COUNT - NON-STICK PADS
 3 COUNT - LARGE FABRIC ADHESIVE PADS
 1 COUNT - ROLL OF TAPE
 1 SET OF PACKETS (16 TABLETS) - IBUPROFEN
 5 COUNT - TRIPLE ANTIBIOTIC FIRST AID OINTMENT PACKETS
 5 COUNT - 1% HYDROCORTISONE CREAM PACKETS
 10 COUNT - CLEANSING WIPES

SEE BACK PANEL FOR SIZES

130
COUNT

welly[®]
First Aid Kit

130 COUNT

Manufactured for: Welly Health PBC
 Minneapolis, MN 55402
 1-833-8E-WELLY
 Welly[®] www.GetWelly.com
 Products of USA (with globally sourced materials) and China.
 Please recycle paper band, tin and inserts.

Product not made with natural rubber latex.
 Bandages are sterile unless individual wrapper is opened or damaged.
 Patent: www.getwelly.com/Patents/851043.07

WLY1043

3 72563 00005 6

Everything you need for when the unbeaten path gets the best of you.

Flexible Fabric bandages that can move with your every move.

Durable metal case keeps it all together.

Cleansing wipes, tape and non-stick pads to help you get back out there.

Triple Antibiotic First Aid Ointment helps prevent infection in minor injuries.

KIT INCLUDES

48 CT. STANDARD FLEX FABRIC BANDAGES 1/2 IN (13 x 75 cm)	16 CT. IBUPROFEN PACKETS (2.5 mg/25 mg/25 mg) 16 TABLETS (25 mg) 20.0 mg (NSAID)
20 CT. SMALL FLEX FABRIC BANDAGES 1/2 IN (13 x 57 cm)	5 CT. TRIPLE ANTIBIOTIC FIRST AID OINTMENT PACKETS (NET WT. 1.97 oz (55 g) PER PACKET)
12 CT. STANDARD WATERPROOF BANDAGES 1 1/2 IN (3.8 x 61 cm)	5 CT. 1% HYDROCORTISONE CREAM PACKETS (NET WT. 1.92 oz (54 g) PER PACKET)
10 CT. BUTTERFLY STRIPS 2 x 2 in (5.1 x 7.6 cm)	10 CT. CLEANSING WIPES
5 CT. NON-STICK PADS 2 x 2 in (5.1 x 7.6 cm)	
3 CT. LARGE FABRIC ADHESIVE PADS 2 3/4 IN (7.0 x 10.2 cm)	
1 CT. ROLL OF TAPE NET WT. 5.0 oz (141 g) x 4.5 in	

BANDAGE DIRECTIONS
 For use on minor cuts and scrapes. Apply bandage to clean, dry skin.
 Water proof bandage: remove bandage after applied. Change daily or when pad becomes wet. DO NOT USE IF SEAL OR OTHER PACKAGING OPENED.

ACTIVE INGREDIENTS
 Triple Antibiotic First Aid Ointment: Bacitracin Zinc, Neomycin Sulfate, Polymyxin B Sulfate
 Ibuprofen: Supina 100 SF, 200mg (NSAID)
 1% Hydrocortisone Cream: 1% Hydrocortisone

Full drug facts include the usual directions for packaging and available at www.getwelly.com

Certified
 Cooperative

Kit Components Labels

welly 

1% Hydrocortisone Cream
0.9g / 1/32 oz

Drug Facts
Active ingredient Purpose
Hydrocortisone 1.0%... Anti-itch

Uses For temporary relief of itching associated with minor skin irritations, inflammation, or rashes. Other uses of product should be only under the advice and supervision of a doctor.

Warnings
For external use only
Do not use • in eyes • for treatment of diaper rash • for feminine itching

Drug Facts (continued)
Warnings (continued)
Stop use, ask a doctor
• if condition worsens or lasts more than 7 days, or clears up and occurs again within a few days • with use of other hydrocortisone products

Keep out of reach of children. If ingested, contact a Poison Control Center right away.

Directions • apply to affected area not more than 3 to 4 times daily • children under 2: ask a doctor

Inactive ingredients
emulsifying wax, ethanol, methylparaben, mineral oil, paraffin, petrolatum, propylparaben, purified water, white wax

Welly Health PBC, Minn., MN 55402
1-833-BE-WELLY

welly 

Triple Antibiotic Ointment
0.5g / 1/32 oz

Drug Facts
Active ingredient Purpose
(in each gram)
Bacitracin zinc, 400 units
Neomycin sulfate, 3.5mg/Neomycin Polymyxin B sulfate First Aid 5000 units Antibiotics

Uses First aid to help prevent infection in minor cuts, scrapes, and burns

Warnings
For external use only

Drug Facts (continued)
Warnings (continued)
Do not use • internally • in eyes • over large areas of the body or on puncture wounds, animal bites or serious burns • for more than 1 week unless directed by a doctor if you are allergic to any of the ingredients

Stop use and ask a doctor if
• rash or allergic reaction develops • condition worsens or persists

Keep out of reach of children. If ingested, contact a Poison Control Center right away.

Directions • clean affected area • apply a small amount 1 to 3 times daily • may cover with a sterile bandage

Inactive Ingredients
petrolatum

Welly Health PBC, Minn., MN 55402
1-833-BE-WELLY

2 Tablets **welly** 

Ibuprofen Tablets

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID)¹... Pain reliever/fever reducer
¹nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:
• headache • toothache • backache • menstrual cramp
• common cold • muscular aches • minor arthritis pain
temporarily reduces fever **Warnings Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • skin redness • asthma (wheezing) • facial swelling • rash • shock • blisters If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions do not use more than directed. The smallest effective dose should be used.**
• do not take longer than 10 days, unless directed by a doctor (see Warnings) **Adults and children: (12 years and older)** • take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not exceed 6 tablets in 24 hours, unless directed by a doctor **Children under 12 years:** ask a doctor. **Other information** • store at 68-77°F (20-25°C) • avoid excessive heat, above 104°F (40°C) • **TAMPER EVIDENT** child resistant packets • do not use any opened or torn packets **Questions or comments?** 1-833-BE-WELLY

LOT AND EXPIRE DATE

© Ltd. for WELLY-HEALTH-PBC • Minneapolis, MN 55402
PKT-IBU-2ct.01

stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions do not use more than directed. The smallest effective dose should be used.**
• do not take longer than 10 days, unless directed by a doctor (see Warnings) **Adults and children: (12 years and older)** • take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not exceed 6 tablets in 24 hours, unless directed by a doctor **Children under 12 years:** ask a doctor. **Other information** • store at 68-77°F (20-25°C) • avoid excessive heat, above 104°F (40°C) • **TAMPER EVIDENT** child resistant packets • do not use any opened or torn packets **Questions or comments?** 1-833-BE-WELLY

LOT AND EXPIRE DATE

© Ltd. for WELLY-HEALTH-PBC • Minneapolis, MN 55402
PKT-IBU-2ct.01

Drug Facts **Pain Reliever & Fever Reducer**

Active ingredient (in each brown tablet) Purpose
Ibuprofen USP, 200 mg (NSAID)¹... Pain reliever/fever reducer
¹nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:
• muscular aches • backache • toothache • menstrual cramps
• headache • the common cold • minor pain of arthritis • temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• shock • facial swelling • asthma (wheezing) • rash • skin redness • blisters • hives
If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• take more or for a longer time than directed

Drug Facts (continued)

• take a blood thinning (anticoagulant) or steroid drug • are age 60 or older
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have had stomach ulcers or bleeding problems
• have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer
• right before or after heart surgery

Ask a doctor before use if
• stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke • you are taking a diuretic
• you have problems or serious side effects from taking pain relievers or fever reducers

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are
• under a doctor's care for any serious condition • taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin • taking any other drug

When using this product
• take with food or milk if stomach upset occurs

Stop use and ask a doctor if
• you experience any of the following signs of stomach bleeding: • Foul faint • have bloody or black stools • vomit blood • have stomach pain that does not get better • you have symptoms of heart problems or stroke
• chest pain • slurred speech • leg swelling • trouble breathing
• weakness in one part or side of body • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area • any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a

Drug Facts (continued)

doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
• do not take more than directed • the smallest effective dose should be used • adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not exceed 6 tablets in 24 hours, unless directed by a doctor • children under 12 years: ask a doctor

Other information
TAMPER EVIDENT. DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
• store between 20°-25°C (68°-77°F) • avoid excessive heat 40°C (104°F) • see label for expiration date and lot number • use by expiration date on package

Drug Facts (continued)

Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, hydroxypropylcellulose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
1-833-BE-WELLY

hydrocortisone, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, ibuprofen kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72663-152
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72663-152-05	1 in 1 KIT; Type 0: Not a Combination Product	04/06/2020	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	5 POUCH	4.5 mL
Part 2	10 POUCH	5 mL in .5
Part 3	8 BLISTER PACK	16 in 8
Part 4	1 POUCH	10

Part 1 of 4

ANTI-ITCH

hydrocortisone cream

Product Information

Item Code (Source)	NDC:72663-580
---------------------------	---------------

Route of Administration	TOPICAL
--------------------------------	---------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE (UNII: W4X0X7BPJ) (HYDROCORTISONE - UNII:W4X0X7BPJ)	HYDROCORTISONE	1 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
MINERAL OIL (UNII: T5L8T28FGP)	
PARAFFIN (UNII: I9O0E3H2ZE)	
PETROLATUM (UNII: 4T6H12BN9U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
WHITE WAX (UNII: 7G1J5DA97F)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72663-580-02	0.9 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/25/2019	

Part 2 of 4

ANTIBIOTIC

bacitracin zinc, neomycin sulfate, polymyxin b sulfate ointment

Product Information

Item Code (Source)	NDC:72663-560
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	6 mg in 1 mL
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 mL
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	0.77 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72663-560-34	0.5 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M004	02/25/2019	

Part 3 of 4

PAIN RELIEF AND FEVER REDUCER

ibuprofen tablet

Product Information

Item Code (Source)	NDC:72663-428
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44291
Contains			

Packaging

Marketing Start	Marketing End
-----------------	---------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72663-428-08	2 in 1 BLISTER PACK; Type 0: Not a Combination Product		



Marketing Information

Marketing

Application Number or Monograph

Marketing Start

Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M013	04/06/2020	

Part 4 of 4

WELLY CLEANSING WIPE

disposable wipes [personal cleanliness] cloth

Product Information

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		10 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Cosmetic		01/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	04/06/2020	

Labeler - Welly Health PBC (116766884)

Revised: 5/2025

Welly Health PBC