

GLUCAMINOLYTE FORTE- glucaminolyte solution
MWI Veterinary Supply, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Glucaminolyte Forte

Drench

Supplemental Nutritive Source

For Animal Use Only

Keep out of reach of children

INDICATIONS

For use as a supplemental nutritive source of electrolytes, dextrose and amino acids in cattle.

DOSAGE AND ADMINISTRATION

Administer orally as a drench or by use of a stomach tube. The usual recommended dose in adult cattle is 500 to 1000 mL, depending on size and condition.

SHAKE WELL BEFORE USING.

STORAGE

Store at controlled room temperature between 15° and 30°C (59-86°F)

TAKE TIME OBSERVE LABEL DIRECTIONS

CONTAINS

Each 1000 mL of aqueous solution contains:

- Dextrose • H₂O 300 g
- Sodium Chloride 5.4 g
- Sodium Acetate, Anhydrous 3.0 g
- Potassium Acetate 1.0 g
- Calcium Chloride • 2H₂O 330.0 mg
- Magnesium Chloride • 5H₂O ... 260.0 mg

Comprised of: L-Glutamic Acid, L-Arginine, Hydrochloride, L-Proline, L-Lysine Hydrochloride, L-Leucine, L-Phnylalanine, L-Valine, L-Threonine, L-Isoleucine, L-Histidine Hydrochloride, L-Methionine, L-Tyosine, with Methylparaben (preservative) 0.18% and Propylparaben (preservative) 0.02%.

NDC 13985-402-52

1000 mL



Glucaminolyte Forte

Drench

INDICATIONS: For use as a supplemental nutritive source of electrolytes, dextrose and amino acids in cattle.

DOSAGE AND ADMINISTRATION: Administer orally as a drench or by use of a stomach tube. The usual recommended dose in adult cattle is 500 to 1000 mL, depending on size and condition.

SHAKE WELL BEFORE USING.

STORAGE: Store at controlled room temperature between 15°C and 30°C (59°F–86°F).

TAKE TIME



OBSERVE LABEL DIRECTIONS

A-0436-07
Iss. 11-11

Distributed by: MWI
Boise, ID 83705
(888) 694-8381
www.vetone.net



Supplemental Nutritive Source

For animal use only.
Keep out of reach
of children.



CONTAINS: Each 1000 mL of aqueous solution contains:

Dextrose • H ₂ O	300.0 g
Sodium Chloride	5.4 g
Sodium Acetate, Anhydrous	3.0 g
Potassium Acetate	1.0 g
Calcium Chloride • 2H ₂ O	330.0 mg
Magnesium Chloride • 6H ₂ O	260.0 mg
L-Glutamic Acid	560.0 mg
L-Arginine Hydrochloride	450.0 mg
L-Proline	400.0 mg
L-Lysine Hydrochloride	240.0 mg
L-Leucine	160.0 mg
L-Phenylalanine	110.0 mg
L-Valine	110.0 mg
L-Threonine	100.0 mg
L-Isoleucine	70.0 mg
L-Histidine Hydrochloride	40.0 mg
L-Methionine	40.0 mg
L-Tyrosine	24.0 mg
with methylparaben 0.18% and propylparaben 0.02% (preservatives).	

LOT NO.:

EXP. DATE:

V1 510204

Net Contents: 1000 mL (33.8 fl oz)

GLUCAMINOLYTE FORTE

glucaminolyte solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:13985-402
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM CHLORIDE (UNII: M410D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	330 mg in 1000 mL
DEXTROSE MONOHYDRATE (UNII: LX22YL083G) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE MONOHYDRATE	300 g in 1000 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	260 mg in 1000 mL
POTASSIUM ACETATE (UNII: M911911U02) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM ACETATE	1 g in 1000 mL
SODIUM ACETATE ANHYDROUS (UNII: NVG71ZZ7P0) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:L4YR4M0NH37)	SODIUM ACETATE ANHYDROUS	3 g in 1000 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698, SODIUM CATION - UNII:L4YR4M0NH37)	SODIUM CHLORIDE	5.4 g in 1000 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-402-52	1000 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		11/02/2011	

Labeler - MWI Veterinary Supply, Inc. (019926120)

Revised: 6/2017

MWI Veterinary Supply, Inc.