# EXTRA STRENGTH PAIN RELIEVER- acetaminophen tablet INGLES MARKETS, INC.

-----

**LL222** 

# **Active ingredient**

Acetaminophen 500 mg

# **Purpose**

Pain Reliever/Fever Reducer

### Uses

- temporarily relieves minor aches and pains
- temporarily reduces fever

# Warnings

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 6

caplets in 24 hours. Severe liver damage may occur if you take:

- more than 8 caplets (4,000 mg of acetaminophen) in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### Do not use

 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

# Ask a doctor before use if you have

liver disease.

# Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

# Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- new symptom occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

### do not take more than directed

12 years of age and	take 1-2 caplets every 4-6 hours, as needed. Do not take more than 6 caplets in 24 hours.
Children under 12 years	Ask a doctor

### Other Information

- TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.
- store at 20°C-25°C (68°F-77°F)

# **Inactive ingredients**

povidone, sodium starch glycolate, starch, stearic acid

### Package label



# NON-ASPIRIN

- pain reliever
- fever reducer



100 CAPLETS

Compare to Active Ingredient in Extra Strength Tylenol®\*

Active Ingredient in Extra Strength Tylenol®









NON-ASPIRIN

- pain reliever
- fever reducer



100 CAPLETS

### Drug Facts

SAVE CARTON FOR COMPLETE DRUG FACTS

Active ingredient (in each caplet)

**Purposes** 

**Uses** • temporarily relieves minor aches and pains • temporarily reduces fever

### Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 6 caplets in 24 hours. Severe liver damage may occur if you take:

more than 8 caplets (4,000 mg of acetaminophen) in 24 hours

with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

### Drug Facts (continued)

Stop use and ask a doctor if . pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions • do not take more than directed

Adults and children 12 Take 1-2 caplets every 4-6 hours, as needed. Do not take more than 6 caplets in 24 hours, unless directed by a doctor. ears of age and over Children under 12 years Ask a doctor.

Other information • TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.

Inactive ingredients: povidone, sodium starch glycolate, starch, stearic acid.

\* This product is not manufactured or distributed by the owner of the registered trademark TYLENOL®.

REV 221-0919



### **EXTRA STRENGTH PAIN RELIEVER**

acetaminophen tablet

### **Product Information**

HUMAN OTC DRUG Item Code (Source) NDC:62936-0222

**Product Type** 

**Route of Administration** 

ORAL

# **Active Ingredient/Active Moiety**

Ingredient Name	<b>Basis of Strength</b>	Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients			
Ingredient Name	Strength		
POVIDONE (UNII: FZ989GH94E)			
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)			
STARCH, CORN (UNII: O8232NY3SJ)			
STEARIC ACID (UNII: 4ELV7Z65AP)			

Product Characteristics			
Color	white (WHITE)	Score	no score
Shape	CAPSULE (Capsule)	Size	17mm
Flavor		Imprint Code	AZ 328
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62936- 0222-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2019	
2	NDC:62936- 0222-5	50 in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2019	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/01/2018	

# Labeler - INGLES MARKETS, INC. (024418584)

# **Registrant -** Geri-Care Pharmaceutical Corp (611196254)

Revised: 2/2024 INGLES MARKETS, INC.