

PAIN RELIEVER- acetaminophen tablet
L.N.K. International, Inc.

Quality Plus 44-104

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - toothache
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has

liver disease.

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- new symptoms occur
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

adults and children 12 years and over	<ul style="list-style-type: none">■ take 2 tablets every 4 to 6 hours while symptoms last■ do not take more than 10 tablets in 24 hours■ do not take for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">■ take 1 tablet every 4 to 6 hours while symptoms last■ do not take more than 5 tablets in 24 hours■ do not take for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, povidone, sodium starch glycolate, stearic acid

Questions or comments?

1-800-426-9391

Principal display panel

QUALITY

+PLUS

NDC 50844-104-16

***Compare to
active ingredient in
Tylenol® Regular Strength**

**REGULAR STRENGTH
PAIN RELIEVER
Acetaminophen 325 mg
PAIN RELIEVER/FEVER REDUCER**

ACTUAL SIZE

1000 Tablets

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

*This product is not manufactured or distributed by Johnson & Johnson Corporation,
owner of the registered
trademark Tylenol® Regular Strength. 50844 ORG042110416

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA



NDC 50844-104-16

*Compare to active ingredient in Tylenol® Regular Strength

REGULAR STRENGTH

PAIN RELIEVER

Acetaminophen 325 mg

PAIN RELIEVER/FEVER REDUCER

1000 Tablets

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Acetaminophen 325 mg

Purpose
Pain reliever/fever reducer

Uses

- Temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - muscular aches
 - minor pain of arthritis
 - the common cold
 - toothache
 - menstrual and menstrual cramps
- Temporarily reduces fever

Warnings

Use as directed. This product contains acetaminophen. Severe liver damage may occur if:

- you take more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- itching
- swelling of the face, lips, tongue, or throat
- difficulty breathing
- fever
- skin rash
- itching
- swelling of the face, lips, tongue, or throat
- difficulty breathing

 If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if the user has liver disease.

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- you feel dizzy or lightheaded
- you get a rash or hives
- you get a fever or sore throat
- you get a stomach ache or diarrhea
- you get a headache or dizziness
- you get a sore throat or difficulty swallowing
- you get a rash or hives
- you get a fever or sore throat
- you get a stomach ache or diarrhea
- you get a headache or dizziness
- you get a sore throat or difficulty swallowing

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children: take 2 tablets every 4 to 6 hours while symptoms last. Do not take more than 10 tablets in 24 hours.
- children 6-11 years: take 1 tablet every 4 to 6 hours while symptoms last. Do not take more than 5 tablets in 24 hours.
- children under 6 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients corn starch, povidone, sodium starch glycolate, stearic acid

Questions or comments? 1-800-468-9381



3 50844 110416 8

Quality Plus 44-104

PAIN RELIEVER

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-104
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;104
Contains			

Packaging

Marketing Start Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-104-16	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/25/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/25/2023	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-104)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-104)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-104)

Revised: 5/2023

L.N.K. International, Inc.