


DANTROLENE SODIUM- dantrolene sodium capsule Par Pharmaceutical, Inc.

Dantrolene Sodium Capsules

Dantrolene sodium has a potential for hepatotoxicity, and should not be used in conditions other than those recommended. Symptomatic hepatitis (fatal and non-fatal) has been reported at various dose levels of the drug. The incidence reported in patients taking up to 400 mg/day is much lower than in those taking doses of 800 mg or more per day. Even sporadic short courses of these higher dose levels within a treatment regimen markedly increased the risk of serious hepatic injury. Liver dysfunction as evidenced by blood chemical abnormalities alone (liver enzyme elevations) has been observed in patients exposed to **dantrolene sodium** for varying periods of time. Overt hepatitis has occurred at varying intervals after initiation of therapy, but has been most frequently observed between the third and twelfth month of therapy. The risk of hepatic injury appears to be greater in females, in patients over 35 years of age, and in patients taking other medication(s) in addition to **dantrolene sodium**. Spontaneous reports suggest a higher proportion of hepatic events with fatal outcome in elderly patients receiving **dantrolene sodium**. However, the majority of these cases were complicated with confounding factors such as intercurrent illnesses and/or concomitant potentially hepatotoxic medications (see Geriatric Use subsection). **Dantrolene sodium** should be used only in conjunction with appropriate monitoring of hepatic function including frequent determination of SGOT or SGPT. If no observable benefit is derived from the administration of **dantrolene sodium** after a total of 45 days, therapy should be discontinued. The lowest possible effective dose for the individual patient should be prescribed.

DESCRIPTION

The chemical formula of **dantrolene sodium** is hydrated 1-[[[5-(4-nitrophenyl)-2-furanyl]methylene]amino]-2, 4-imidazolidinedione sodium salt. It is an orange powder, slightly soluble in water, but due to its slightly acidic nature the solubility increases somewhat in alkaline solution. The anhydrous salt has a molecular weight of 336. The hydrated salt contains approximately 15% water (3-1/2 moles) and has a molecular weight of 399. The structural formula for the hydrated salt is:



Dantrolene sodium is supplied in capsules of 25 mg, 50 mg and 100 mg.

Inactive Ingredients: Each capsule contains Edible black ink, FD&C Yellow No. 6, Gelatin, Lactose, Magnesium stearate, Starch, Synthetic iron oxide red, Synthetic iron oxide yellow, Talc, and Titanium dioxide.

CLINICAL PHARMACOLOGY

In isolated nerve-muscle preparation, **dantrolene sodium** has been shown to produce relaxation by affecting the contractile response of the skeletal muscle at a site beyond the myoneural junction, directly on the muscle itself. In skeletal muscle, **dantrolene sodium** dissociates the excitation-contraction coupling, probably by interfering with the release of Ca^{++} from the sarcoplasmic reticulum. This effect

appears to be more pronounced in fast muscle fibers as compared to slow ones, but generally affects both. A central nervous system effect occurs, with drowsiness, dizziness, and generalized weakness occasionally present. Although **dantrolene sodium** does not appear to directly affect the CNS, the extent of its indirect effect is unknown. The absorption of **dantrolene sodium** after oral administration in humans is incomplete and slow but consistent, and dose-related blood levels are obtained. The duration and intensity of skeletal muscle relaxation is related to the dosage and blood levels. The mean biologic half-life of **dantrolene sodium** in adults is 8.7 hours after a 100-mg dose. Specific metabolic pathways in the degradation and elimination of **dantrolene sodium** in human subjects have been established. Metabolic patterns are similar in adults and pediatric patients. In addition to the parent compound, dantrolene, which is found in measurable amounts in blood and urine, the major metabolites noted in body fluids are the 5-hydroxy analog and the acetamido analog. Since **dantrolene sodium** is probably metabolized by hepatic microsomal enzymes, enhancement of its metabolism by other drugs is possible. However, neither phenobarbital nor diazepam appears to affect **dantrolene sodium** metabolism.

Clinical experience in the management of fulminant human malignant hyperthermia, as well as experiments conducted in malignant hyperthermia susceptible swine, have revealed that the administration of intravenous dantrolene, combined with indicated supportive measures, is effective in reversing the hypermetabolic process of malignant hyperthermia. Known differences between human and swine malignant hyperthermia are minor. The prophylactic administration of oral or intravenous dantrolene to malignant hyperthermia susceptible swine will attenuate or prevent the development of signs of malignant hyperthermia in a manner dependent upon the dosage of dantrolene administered and the intensity of the malignant hyperthermia triggering stimulus. Limited clinical experience with the administration of oral dantrolene to patients judged malignant hyperthermia susceptible, when combined with clinical experience in the use of intravenous dantrolene for the treatment of malignant hyperthermia and data derived from the above cited animal model experiments, suggests that oral dantrolene will also attenuate or prevent the development of signs of human malignant hyperthermia, provided that currently accepted practices in the management of such patients are adhered to (see **INDICATIONS AND USAGE**); intravenous dantrolene should also be available for use should the signs of malignant hyperthermia appear.

INDICATIONS AND USAGE

In Chronic Spasticity

Dantrolene sodium is indicated in controlling the manifestations of clinical spasticity resulting from upper motor neuron disorders (e.g., spinal cord injury, stroke, cerebral palsy, or multiple sclerosis). It is of particular benefit to the patient whose functional rehabilitation has been retarded by the sequelae of spasticity. Such patients must have presumably reversible spasticity where relief of spasticity will aid in restoring residual function. **Dantrolene sodium** is not indicated in the treatment of skeletal muscle spasm resulting from rheumatic disorders.

If improvement occurs, it will ordinarily occur within the dosage titration (see **DOSAGE AND ADMINISTRATION**), and will be manifested by a decrease in the severity of spasticity and the ability to resume a daily function not quite attainable without **dantrolene sodium**.

Occasionally, subtle but meaningful improvement in spasticity may occur with **dantrolene sodium** therapy. In such instances, information regarding improvement should be solicited from the patient and those who are in constant daily contact and attendance with him. Brief withdrawal of **dantrolene sodium** for a period of 2 to 4 days will frequently demonstrate exacerbation of the manifestations of spasticity and may serve to confirm a clinical impression.

A decision to continue the administration of **dantrolene sodium** on a long-term basis is justified if introduction of the drug into the patient's regimen:

- produces a significant reduction in painful and/or disabling spasticity such as clonus, or

- permits a significant reduction in the intensity and/or degree of nursing care required, or
- rids the patient of any annoying manifestation of spasticity considered important by the patient himself.

In Malignant Hyperthermia

Oral **dantrolene sodium** is also indicated preoperatively to prevent or attenuate the development of signs of malignant hyperthermia in known, or strongly suspect, malignant hyperthermia susceptible patients who require anesthesia and/or surgery. Currently accepted clinical practices in the management of such patients must still be adhered to (careful monitoring for early signs of malignant hyperthermia, minimizing exposure to triggering mechanisms and prompt use of intravenous Dantrolene Sodium and indicated supportive measures should signs of malignant hyperthermia appear); see also the package insert for **Dantrium**[®] (dantrolene sodium) **Intravenous**.

Oral **dantrolene sodium** should be administered following a malignant hyperthermic crisis to prevent recurrence of the signs of malignant hyperthermia.

CONTRAINDICATIONS

Active hepatic disease, such as hepatitis and cirrhosis, is a contraindication for use of **dantrolene sodium**. **Dantrolene sodium** is contraindicated where spasticity is utilized to sustain upright posture and balance in locomotion or whenever spasticity is utilized to obtain or maintain increased function.

WARNINGS

It is important to recognize that fatal and non-fatal liver disorders of an idiosyncratic or hypersensitivity type may occur with **dantrolene sodium** therapy.

At the start of **dantrolene sodium** therapy, it is desirable to do liver function studies (SGOT, SGPT, alkaline phosphatase, total bilirubin) for a baseline or to establish whether there is pre-existing liver disease. If baseline liver abnormalities exist and are confirmed, there is a clear possibility that the potential for **dantrolene sodium** hepatotoxicity could be enhanced, although such a possibility has not yet been established.

Liver function studies (e.g., SGOT or SGPT) should be performed at appropriate intervals during **dantrolene sodium** therapy. If such studies reveal abnormal values, therapy should generally be discontinued. Only where benefits of the drug have been of major importance to the patient, should reinitiation or continuation of therapy be considered. Some patients have revealed a return to normal laboratory values in the face of continued therapy while others have not.

If symptoms compatible with hepatitis, accompanied by abnormalities in liver function tests or jaundice appear, **dantrolene sodium** should be discontinued. If caused by **dantrolene sodium** and detected early, the abnormalities in liver function characteristically have reverted to normal when the drug was discontinued.

Dantrolene sodium therapy has been reinstated in a few patients who have developed clinical and/or laboratory evidence of hepatocellular injury. If such reinstatement of therapy is done, it should be attempted only in patients who clearly need **dantrolene sodium** and only after previous symptoms and laboratory abnormalities have cleared. The patient should be hospitalized and the drug should be restarted in very small and gradually increasing doses. Laboratory monitoring should be frequent and the drug should be withdrawn immediately if there is any indication of recurrent liver involvement. Some patients have reacted with unmistakable signs of liver abnormality upon administration of a challenge dose, while others have not.

Dantrolene sodium should be used with particular caution in females and in patients over 35 years of age in view of apparent greater likelihood of drug-induced, potentially fatal, hepatocellular disease in these groups. Spontaneous reports suggest a higher proportion of hepatic events with fatal outcome in

elderly patients receiving **dantrolene sodium**. However, the majority of these cases were complicated with confounding factors such as intercurrent illnesses and/or concomitant potentially hepatotoxic medications (see Geriatric Use subsection).

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term safety of **dantrolene sodium** in humans has not been established. Chronic studies in rats, dogs, and monkeys at dosages greater than 30 mg/kg/day showed growth or weight depression and signs of hepatopathy and possible occlusion nephropathy, all of which were reversible upon cessation of treatment. Sprague-Dawley female rats fed dantrolene sodium for 18 months at dosage levels of 15, 30, and 60 mg/kg/day showed an increased incidence of benign and malignant mammary tumors compared with concurrent controls. At the highest dose level, there was an increase in the incidence of benign lymphatic neoplasms. In a 30-month study at the same dose levels also in Sprague-Dawley rats, Dantrolene Sodium produced a decrease in the time of onset of mammary neoplasms. Female rats at the highest dose level showed an increased incidence of hepatic lymphangiomas and hepatic angiosarcomas.

The only drug-related effect seen in a 30-month study in Fischer-344 rats was a dose-related reduction in the time of onset of mammary and testicular tumors. A 24-month study in HaM/ICR mice revealed no evidence of carcinogenic activity. Carcinogenicity in humans cannot be fully excluded, so that this possible risk of chronic administration must be weighed against the benefits of the drug (i.e., after a brief trial) for the individual patient.

Dantrolene Sodium has produced positive results in the Ames S. Typhimurium bacterial mutagenesis assay in the presence and absence of a liver activating system.

Pregnancy

Pregnancy Category C

Adequate animal reproduction studies have not been conducted with **dantrolene sodium**. It is also not known whether **dantrolene sodium** can cause fatal harm when administered to a pregnant woman or can affect reproduction capacity. **Dantronlene sodium** should be given to a pregnant woman only if clearly needed.

Labor and Delivery

In one non-randomized open-label study, 21 term pregnant patients received prophylactic oral **dantrolene sodium** 100 mg per day for 2 to 10 days prior to delivery. Dantrolene readily crossed the placenta with maternal and fetal whole blood levels approximately equal at delivery; neonatal levels then fell approximately 50% per day for 2 days before declining sharply. No neonatal respiratory and neuromuscular side effects were detected at low dose. More data, at higher doses, are needed before more definitive conclusions can be made.

Nursing Mothers

Dantrolene sodium should not be used in nursing mothers.

Usage in Pediatric Patients

The long-term safety of **dantrolene sodium** in pediatric patients under the age of 5 years has not been established. Because of the possibility that adverse effects of the drug could become apparent only after many years, a benefit-risk consideration of the long-term use of **dantrolene sodium** is particularly important in pediatric patients.

Geriatric Use

Clinical studies of **dantrolene sodium** did not include sufficient numbers of subjects aged 65 and over

to determine whether they respond differently from younger subjects. Other reported clinical experience in the literature has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy. As with all patients receiving **dantrolene sodium**, it is recommended that elderly patients receive the lowest dose compatible with the optimal response. Spontaneous reports suggest a higher proportion of hepatic events with fatal outcome in elderly patients receiving **dantrolene sodium**. However, the majority of these cases were complicated with confounding factors such as intercurrent illnesses and/or concomitant potentially hepatotoxic medications (for hepatotoxicity details and its management see **Black Box** and **Warnings Sections**).

Drug Interactions

Drowsiness may occur with **dantrolene sodium** therapy, and the concomitant administration of CNS depressants such as sedatives and tranquilizing agents may result in further drowsiness.

While a definite drug interaction with estrogen therapy has not yet been established, caution should be observed if the two drugs are to be given concomitantly. Hepatotoxicity has occurred more often in women over 35 years of age receiving concomitant estrogen therapy.

Cardiovascular collapse in patients treated simultaneously with verapamil and Dantrolene Sodium is rare. The combination of therapeutic doses of intravenous Dantrolene Sodium and verapamil in halothane/ α -chloralose anesthetized swine has resulted in ventricular fibrillation and cardiovascular collapse in association with marked hyperkalemia. Until the relevance of these findings to humans is established, the combination of Dantrolene Sodium and calcium channel blockers is not recommended during the management of malignant hyperthermia.

Administration of **dantrolene sodium** may potentiate vecuronium-induced neuromuscular block.

PRECAUTIONS

Dantrolene sodium should be used with caution in patients with impaired pulmonary function, particularly those with obstructive pulmonary disease, and in patients with severely impaired cardiac function due to myocardial disease. **Dantrolene sodium** is associated with pleural effusion with associated eosinophilia. It should be used with caution in patients with a history of previous liver disease or dysfunction (see **WARNINGS**).

Information for Patients

Patients should be cautioned against driving a motor vehicle or participating in hazardous occupations while taking **dantrolene sodium**. Caution should be exercised in the concomitant administration of tranquilizing agents.

Dantrolene sodium might possibly evoke a photosensitivity reaction; patients should be cautioned about exposure to sunlight while taking it.

ADVERSE REACTIONS

The most frequently occurring side effects of **dantrolene sodium** have been drowsiness, dizziness, weakness, general malaise, fatigue, and diarrhea. These are generally transient, occurring early in treatment, and can often be obviated by beginning with a low dose and increasing dosage gradually until an optimal regimen is established. Diarrhea may be severe and may necessitate temporary withdrawal of **dantrolene sodium** therapy. If diarrhea recurs upon readministration of **dantrolene sodium**, therapy should probably be withdrawn permanently.

Other less frequent side effects, listed according to system, are:

Gastrointestinal: Constipation, rarely progressing to signs of intestinal obstruction, GI bleeding,

anorexia, swallowing difficulty, gastric irritation, abdominal cramps, nausea and/or vomiting.

Hepatobiliary: Hepatitis (see **WARNINGS**).

Neurologic: Speech disturbance, seizure, headache, light-headedness, visual disturbance, diplopia, alteration of taste, insomnia, drooling.

Cardiovascular: Tachycardia, erratic blood pressure, phlebitis, heart failure.

Hematologic: Aplastic anemia, anemia, leukopenia, lymphocytic lymphoma, thrombocytopenia.

Psychiatric: Mental depression, mental confusion, increased nervousness.

Urogenital: Increased urinary frequency, crystalluria, hematuria, difficult erection, urinary incontinence and/or nocturia, difficult urination and/or urinary retention.

Integumentary: Abnormal hair growth, acne-like rash, pruritus, urticaria, eczematoid eruption, sweating.

Musculoskeletal: Myalgia, backache.

Respiratory: Feeling of suffocation, respiratory depression.

Special Senses: Excessive tearing.

Hypersensitivity: Pleural effusion with pericarditis, pleural effusion with associated eosinophilia, anaphylaxis.

Other: Chills and fever.

The published literature has included some reports of **dantrolene sodium** use in patients with Neuroleptic Malignant Syndrome (NMS). **Dantrolene sodium** capsules are not indicated for the treatment of NMS and patients may expire despite treatment with **dantrolene sodium** capsules.

For medical advice about adverse reactions contact your medical professional. To report SUSPECTED ADVERSE REACTIONS, contact Par Pharmaceutical at 1-800-828-9393 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG ABUSE AND DEPENDENCE

Drug abuse and dependency potential has not been evaluated in human or animal studies.

OVERDOSAGE

Symptoms which may occur in case of overdose include, but are not limited to, muscular weakness and alterations in the state of consciousness (e.g. lethargy, coma), vomiting, diarrhea, and crystalluria. For acute overdose, general supportive measures should be employed along with immediate gastric lavage.

Intravenous fluids should be administered in fairly large quantities to avert the possibility of crystalluria. An adequate airway should be maintained and artificial resuscitation equipment should be at hand. Electrocardiographic monitoring should be instituted, and the patient carefully observed. To date, no experience has been reported with dialysis and its value in **dantrolene sodium** overdose is not known.

DOSAGE AND ADMINISTRATION

For Use in Chronic Spasticity

Prior to the administration of **dantrolene sodium**, consideration should be given to the potential response to treatment. A decrease in spasticity sufficient to allow a daily function not otherwise attainable should be the therapeutic goal of treatment with **dantrolene sodium**. Refer to

INDICATIONS AND USAGE section for description of response to be anticipated.

It is important to establish a therapeutic goal (regain and maintain a specific function such as therapeutic exercise program, utilization of braces, transfer maneuvers, etc.) before beginning **dantrolene sodium** therapy. Dosage should be increased until the maximum performance compatible with the dysfunction due to underlying disease is achieved. No further increase in dosage is then indicated.

Usual Dosage

It is important that the dosage be titrated and individualized for maximum effect. The lowest dose compatible with optimal response is recommended.

*In view of the potential for liver damage in long-term **dantrolene sodium** use, therapy should be stopped if benefits are not evident within 45 days.*

Adults

The following gradual titration schedule is suggested. Some patients will not respond until higher daily dosage is achieved. Each dosage level should be maintained for seven days to determine the patient's response. If no further benefit is observed at the next higher dose, dosage should be decreased to the previous lower dose.

- 25 mg once daily for seven days, then
- 25 mg t.i.d. for seven days
- 50 mg t.i.d. for seven days
- 100 mg t.i.d.

Therapy with a dose four times daily may be necessary for some individuals. Doses higher than 100 mg four times daily should not be used. (See **Box Warning**.)

Pediatric Patients

The following gradual titration schedule is suggested. Some patients will not respond until higher daily dosage is achieved. Each dosage level should be maintained for seven days to determine the patient's response. If no further benefit is observed at the next higher dose, dosage should be decreased to the previous lower dose.

- 0.5 mg/kg once daily for seven days, then
- 0.5 mg/kg t.i.d. for seven days
- 1 mg/kg t.i.d. for seven days
- 2 mg/kg t.i.d.

Therapy with a dose four times daily may be necessary for some individuals. Doses higher than 100 mg four times daily should not be used. (See **Box Warning**.)

For Malignant Hyperthermia

Preoperatively

Administer 4 to 8 mg/kg/day of oral **dantrolene sodium** in 3 or 4 divided doses for one or two days prior to surgery, with the last dose being given approximately 3 to 4 hours before scheduled surgery with a minimum of water.

This dosage will usually be associated with skeletal muscle weakness and sedation (sleepiness or drowsiness); adjustment can usually be made within the recommended dosage range to avoid incapacitation or excessive gastrointestinal irritation (including nausea and/or vomiting).

Post Crisis Follow-up

Oral **dantrolene sodium** should also be administered following a malignant hyperthermia crisis, in

doses of 4 to 8 mg/kg per day in four divided doses, for a one to three day period to prevent recurrence of the manifestations of malignant hyperthermia.

HOW SUPPLIED

Dantrolene sodium is available in:

25-mg opaque, orange and tan capsules imprinted with **DANTRIUM** 25 mg on the cap and 0149 0030 with a single bar on the body.

NDC 49884-362-01 bottle of 100

50-mg opaque, orange and tan capsules imprinted with **DANTRIUM** 50 mg on the cap and 0149 0031 with a double bar on the body.

NDC 49884-363-01 bottle of 100

100-mg opaque, orange and tan capsules imprinted with **DANTRIUM** 100 mg on the cap and 0149 0033 with a triple bar on the body.

NDC 49884-364-01 bottle of 100

Store between 20° to 25°C (68° to 77°F)[See USP Controlled Room Temperature].

Rx Only.

Dist. by:

Par Pharmaceutical

Chestnut Ridge, NY 10977 U.S.A.

Mfg. by:

Par Formulations Private Limited,

9/215, Pudupakkam, Kelambakkam – 603 103.

Made in India

OS362-01-74-01

Revised: 07/2017

PRINCIPAL DISPLAY PANEL - 25 mg Capsule Bottle Label

NDC 49884-362-01

Dantrolene

Sodium Capsules

25 mg

For oral use only

PERIPHERAL SKELETAL MUSCLE RELAXANT

Rx only

100 Capsules

PAR PHARMACEUTICAL

NDC 49884-362-01

Dantrolene Sodium Capsules

25 mg

For oral use only

PERIPHERAL SKELETAL MUSCLE RELAXANT

Rx only

100 Capsules



THIS IS A BULK CONTAINER AND NOT INTENDED FOR DISPENSING.

Dosage: Dosage must be titrated and individualized for maximum effect. See package insert for full prescribing information.

Dantrolene Sodium contains about 15% water of hydration.

Dispense in a tight container.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Store between 20° to 25°C (68° to 77°F). (See USP Controlled Room Temperature).

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.

Mfg. by: Par Formulations Private Limited,
9/215, Pudupakkam, Kelambakkam - 603 103.

Made in India

Mfg. Lic. No.: TN00002121



R07/17
LA362-01-74-01
0
N 3 4 9 8 8 4 1 3 6 2 0 1

Unvarnished Area
W17 X H45 mm

PRINCIPAL DISPLAY PANEL - 50 mg Capsule Bottle Label

NDC 49884-363-01

Dantrolene Sodium Capsules

50 mg

For oral use only

PERIPHERAL SKELETAL MUSCLE RELAXANT

Rx only

100 Capsules

PAR PHARMACEUTICAL

NDC 49884-363-01

Dantrolene Sodium Capsules

50 mg

For oral use only

PERIPHERAL SKELETAL MUSCLE RELAXANT

Rx only

100 Capsules



THIS IS A BULK CONTAINER AND NOT INTENDED FOR DISPENSING.

Dosage: Dosage must be titrated and individualized for maximum effect. See package insert for full prescribing information.

Dantrolene Sodium contains about 15% water of hydration.

Dispense in a tight container.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Store between 20° to 25°C (68° to 77°F). (See USP Controlled Room Temperature).

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.

Mfg. by: Par Formulations Private Limited,
9/215, Pudupakkam, Kelambakkam - 603 103.

Made in India

Mfg. Lic. No.: TN00002121



R07/17
LA363-01-74-01
7
N 3 4 9 8 8 4 1 3 6 3 0 1

Unvarnished Area
W17 X H45 mm

PRINCIPAL DISPLAY PANEL – 100 mg Capsules Bottle Label

NDC 49884-364-01

Dantrolene Sodium Capsules

100 mg

For oral use only
PERIPHERAL SKELETAL MUSCLE RELAXANT
 Rx only
100 Capsules
PAR PHARMACEUTICAL

NDC 49884-364-01

Dantrolene Sodium Capsules

100 mg

For oral use only
 PERIPHERAL SKELETAL MUSCLE RELAXANT
 Rx only
100 Capsules



THIS IS A BULK CONTAINER AND NOT INTENDED FOR DISPENSING.

Dosage: Dosage must be titrated and individualized for maximum effect. See package outsert for full prescribing information.

Dantrolene Sodium contains about 15% water of hydration.
 Dispense in a tight container.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Store between 20° to 25°C (68° to 77°F).
 (See USP Controlled Room Temperature).

Dist. by: **Par Pharmaceutical**
 Chestnut Ridge, NY 10977 U.S.A.

Mfg. by: **Par Formulations Private Limited,**
 9/215, Pudupakkam, Kelambakkam - 603 103.

Made in India

Mfg. Lic. No.: TN00002121



R07/17

LA364-01-74-01

Unvarnished Area
 W17 X H50 mm

DANTROLENE SODIUM

dantrolene sodium capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:49884-362
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DANTROLENE SODIUM (UNII: 287M0347EV) (DANTROLENE - UNII:F64QU97QCR)	DANTROLENE SODIUM	25 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	ORANGE, BROWN (tan)	Score	no score
Shape	CAPSULE	Size	16mm
Flavor		Imprint Code	Dantrium;25;mg;0 149;0030
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49884-362-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/28/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC	NDA017443	03/28/2016	

DANTROLENE SODIUM

dantrolene sodium capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:49884-363
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DANTROLENE SODIUM (UNII: 287M0347EV) (DANTROLENE - UNII:F64QU97QCR)	DANTROLENE SODIUM	50 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	ORANGE, BROWN (tan)	Score	no score
Shape	CAPSULE	Size	18mm

Flavor		Imprint Code	Dantrium;50 ;mg;0 149 ;0031	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49884-363-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/28/2016	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA AUTHORIZED GENERIC	NDA017443	03/28/2016		

DANTROLENE SODIUM				
dantrolene sodium capsule				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:49884-364	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	DANTROLENE SODIUM (UNII: 287M0347EV) (DANTROLENE - UNII:F64QU97QCR)	DANTROLENE SODIUM	100 mg	
Inactive Ingredients				
	Ingredient Name	Strength		
	FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
	GELATIN (UNII: 2G86QN327L)			
	LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
	MAGNESIUM STEARATE (UNII: 70097M6I30)			
	TALC (UNII: 7SEV7J4R1U)			
	TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
	STARCH, CORN (UNII: O8232NY3SJ)			
	FERRIC OXIDE YELLOW (UNII: EX438O2MRT)			
	FERRIC OXIDE RED (UNII: 1K09F3G675)			
Product Characteristics				
Color	ORANGE, BROWN	Score	no score	
Shape	CAPSULE	Size	20mm	
Flavor		Imprint Code	Dantrium;100 ;mg;0 149 ;0033	
Contains				

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49884-364-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/28/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC	NDA017443	03/28/2016	

Labeler - Par Pharmaceutical, Inc. (092733690)

Establishment

Name	Address	ID/FEI	Business Operations
Par Formulations Private Limited		676159161	ANALYSIS(49884-362, 49884-363, 49884-364) , MANUFACTURE(49884-362, 49884-363, 49884-364) , PACK(49884-362, 49884-363, 49884-364)

Revised: 12/2017

Par Pharmaceutical, Inc.