

**COLD AND FLU PLUS CONGESTION NIGHTTIME- acetaminophen,
dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution
WALMART INC.**

Equate 44-060

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 20 mg
Doxylamine succinate 12.5 mg
Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - runny nose and sneezing
 - nasal congestion
 - sore throat
 - headache
 - minor aches and pains
 - cough to help you sleep
 - sinus congestion and pressure
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters

- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- glaucoma
- heart disease
- high blood pressure
- liver disease
- thyroid disease
- diabetes

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- mL = milliliter
- only use the dose cup provided
- do not exceed 4 doses per 24 hours
- adults and children 12 years and over: 30 mL every 4 hours
- children under 12 years: do not use

Other information

- **each 30 mL contains:** sodium 11 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, D&C yellow #10, FD&C blue #1, flavors, glycerin, polyethylene glycol 400, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sorbitol, sucralose

Questions or comments?

1-888-287-1915

Principal display panel

equate[™]

NDC 79903-191-12

Compare to
Vicks® NyQuil®
VapoCOOL® Severe
Cold & Flu + Congestion
active ingredients*

NIGHTTIME

SEVERE
VAPOR CHILLING

**Cold & Flu
+ Congestion**

- **Acetaminophen** - Pain Reliever/Fever Reducer
- Dextromethorphan HBr - Cough Suppressant
- Doxylamine Succinate - Antihistamine
- Phenylephrine HCl - Nasal Decongestant

Relieves:

- Headache, fever, sore throat, minor aches & pains
- Nasal congestion, sinus pressure
- Sneezing, runny nose
- Cough

For Ages 12+

12 FL OZ (355 mL)

**F-060
ORG**

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® VapoCOOL® SEVERE COLD & FLU + CONGESTION.

PARENTS:

**Learn about teen medicine abuse
www.StopMedicineAbuse.org**

50844 ORG062306002
B-060 ORG



Cold & Flu + Congestion

- **Acetaminophen** – Pain Reliever/Fever Reducer
- **Dextromethorphan HBr** – Cough Suppressant
- **Doxylamine Succinate** – Antihistamine
- **Phenylephrine HCl** – Nasal Decongestant

Relieves:

- Headache, fever, sore throat, minor aches & pains
- Nasal congestion, sinus pressure
- Sneezing, runny nose
- Cough

For Ages 12+

12 FL OZ (355 mL)

F-060
ORG

- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin reddening. If a skin reaction occurs, stop use and seek medical help right away. ▶

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® VapoCOOL® SEVERE COLD & FLU + CONGESTION.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

60844 ORG062306002

B-060 ORG

1 94346 11210 4

NO PRINT / NO VARNISH AREA
LOT NO. & EXP. DATE

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- glaucoma
- heart disease
- high blood pressure
- liver disease
- thyroid disease
- diabetes

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery ▶

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- mL = milliliter
- only use the dose cup provided
- do not exceed 4 doses per 24 hours
- adults and children 12 years and over: 30 mL every 4 hours
- children under 12 years: do not use

Other information

- each 30 mL contains: sodium 11 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous citric acid, D&C yellow #10, FD&C blue #1, flavors, glycerin, polyethylene glycol 400, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sorbitol, sucralose

Questions or comments?

1-888-287-1915

COLD AND FLU PLUS CONGESTION NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79903-191
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	green (bluish)	Score	
Shape		Size	
Flavor	MINT (eucalyptus)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a	08/16/2022	

191-12	Combination Product	08/16/2023	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/16/2023	

Labeler - WALMART INC. (051957769)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(79903-191) , pack(79903-191)

Revised: 8/2025

WALMART INC.