

NAPROXEN SODIUM- naproxen sodium tablet, coated
VALU MERCHANDISERS COMPANY

1145-BST-2024-0613

Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) *

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-822-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

adults and children 12 years and older	<ul style="list-style-type: none"> ▪ take 1 caplet every 8 to 12 hours while symptoms last ▪ for the first dose you may take 2 caplets within the first hour ▪ do not exceed 2 caplets in any 8- to 12-hour period ▪ do not exceed 3 caplets in a 24-hour period
children under 12 years	<ul style="list-style-type: none"> ▪ ask a doctor

Other information

- **each caplet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- retain carton for complete product information and warnings

Inactive ingredients

croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

Questions or comments?

1-844-292-1112

PRINCIPAL DISPLAY PANEL

Best Choice®

†Compare to the active ingredient in Aleve® Caplets

Naproxen Sodium

TABLETS, 220mg

PAIN RELIEVER/FEVER REDUCER

(NSAID)

Strength to last 12 hours

Actual Size

50 CAPLETS**

**CAPSULE-SHAPED TABLETS

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING

F114502BST_R2



This product is not manufactured or distributed or distributed by Bayer Consumer Healthcare, LLC, distributor of Aleve® Caplets.

**DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING**

Drug Facts

Active ingredient (in each caplet) Purpose
Naproxen sodium 220 mg Pain reliever/
Naproxen 200 mg (NSAID)fever reducer
nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - headache
 - the common cold
 - temporarily reduces fever
- menstrual cramps
- toothache

Warnings

- Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
 - facial swelling
 - asthma (wheezing)
 - shock
 - skin redness/ rash
 - dizziness
- If an allergic reaction occurs, stop use and seek medical help right away.
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 - you are taking a diuretic
 - you have problems or serious side effects from taking pain relievers or fever reducers
- Ask a doctor or pharmacist before use if you are**
- under a doctor's care for any serious condition
 - taking any other drug

When using this product

- take with food or milk, if stomach upset occurs
- Stop use and ask a doctor if
 - you experience any of the following signs of stomach bleeding:
 - have bloody or black stools
 - feel faint
 - vomit blood
 - have stomach pain that does not get better
 - you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat

Drug Facts (continued)

- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
 - the smallest effective dose should be used
 - drink a full glass of water with each dose
- | | |
|--|--|
| adults and children 12 years and older | <ul style="list-style-type: none"> ■ take 1 caplet every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 caplets within the first hour ■ do not exceed 2 caplets in any 8- to 12-hour period ■ do not exceed 3 caplets in a 24-hour period |
| children under 12 years | <ul style="list-style-type: none"> ■ ask a doctor |

Other information

- each caplet contains sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- retain carton for complete product information and warnings




PROUDLY DISTRIBUTED BY:
ASSOCIATED WHOLESOME PRODUCTS, INC.
KANSAS CITY, KANSAS 66108
Call 1-844-292-1112 for more product information.
www.bestchoicebrand.com

100% Guaranteed
Best Choice.

Drug Facts (continued)
Inactive ingredients FD&C blue #2, macrogol, croscarmellose sodium, pregelatinized starch, talc, titanium dioxide, magnesium stearate, polyvinyl alcohol, povidone, croscarmellose sodium, FD&C blue #2, macrogol, croscarmellose sodium, pregelatinized starch, talc, titanium dioxide

Questions or comments?



†Compare to the active ingredient in Aleve® Caplets

Naproxen Sodium

TABLETS, 220 mg
PAIN RELIEVER/FEVER REDUCER
(NSAID)

Strength to last 12 hours

50 CAPLETS**
**Capsule-shaped tablets

Actual Size

NAPROXEN SODIUM naproxen sodium tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-445
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL (Biconvex)	Size	12mm
Flavor		Imprint Code	I7
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-445-01	1 in 1 CARTON	05/01/2024	03/31/2028
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63941-445-02	1 in 1 CARTON	05/01/2024	03/31/2028
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:63941-445-03	1 in 1 CARTON	05/01/2024	06/30/2028
3		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079096	05/01/2024	06/30/2028

Revised: 12/2025

VALU MERCHANDISERS COMPANY