

**NAPROXEN SODIUM - naproxen sodium tablet, film coated  
WALMART INC.**

-----

**Naproxen Sodium Tablets, USP 220 mg**

***Drug Facts***

***Active ingredient (in each caplet)***

Naproxen sodium USP 220 mg (naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purposes***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- menstrual cramps
- toothache
- backache
- headache
- the common cold
- temporarily reduces fever

***Warnings***

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> <li>• take 1 caplet every 8 to 12 hours while symptoms last</li> <li>• for the first dose you may take 2 caplets within the first hour</li> <li>• do not exceed 2 caplets in any 8 to 12 hour period</li> <li>• do not exceed 3 caplets in a 24 hour period</li> </ul>
Children under 12 years	<ul style="list-style-type: none"> <li>• ask a doctor</li> </ul>

**Other information**

- **each caplet contains:** sodium 20 mg
- store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients**

colloidal silicon dioxide, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, and titanium dioxide

**Questions or comments?**

**1-888-287-1915**

**DISTRIBUTED BY:  
Walmart Inc.,  
Bentonville, AR 72716**

PRODUCT OF INDIA

\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve® Caplets.

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 220 mg (90 Tablets Bottle)**

**NDC 79903-232-90**

**equate™  
All Day Pain Relief**

**Naproxen Sodium  
Tablets USP, 220 mg  
(NSAID)**

**Pain Reliever/  
Fever Reducer  
Strength to Last 12 Hours**

**12  
HOURS**

**220  
mg**

**90  
CAPLETS†  
†Capsule-Shaped Tablets**

**equate™**

**All Day  
Pain Relief**  
Naproxen Sodium  
Tablets USP, 220 mg  
(NSAID)

NDC 79903-232-90

**Drug Facts**

**Active ingredient (in each caplet)**  
Naproxen sodium USP 220 mg (naproxen 200 mg)  
(NSAID)\*

**Purpose(s)**

Pain reliever/fever reducer

\*nonsteroidal anti-inflammatory drug

**Uses** ■temporarily relieves minor aches and pains due to: ■minor pain of arthritis ■muscular aches ■backache ■headache ■the common cold ■menstrual cramps ■toothache ■temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■hives ■facial swelling ■asthma (wheezing) ■shock ■skin reddening ■rash ■blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■are age 60 or older ■have had stomach ulcers or bleeding problems ■take a blood thinning (anticoagulant) or steroid drug ■take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■have 3 or more alcoholic drinks every day while using this product ■take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use**

**DO NOT USE IF SEAL OVER BOTTLE OPENING IS BROKEN OR MISSING.**

Lift Here

Satisfaction guaranteed -  
For questions or comments  
please call 1-888-287-1915.

**DISTRIBUTED BY: Walmart Inc.,  
Bentonville, AR 72716  
PRODUCT OF INDIA 535063**

P1433825 LM-5511

\*  
**Unvarnished Zone  
(dotted line not for printing)  
30 x 14 mm**

**Pain Reliever/  
Fever Reducer**  
Strength to Last  
12 Hours

12  
HOURS

220  
mg

90  
CAPLETS\*

\*Capulet-Shaped Tablets



\*Lot: XXXXXXXXX  
Exp.: YYYY-MMM  
Prefix, Variables of Lot, Exp and  
Neutral code shall be  
printed online during packing.

**Drug Facts (continued)**

■if you have ever had an allergic reaction to any other pain reliever/fever reducer ■right before or after heart surgery. **Ask a doctor before use if** ■the stomach bleeding warning applies to you ■you have a history of stomach problems, such as heartburn ■you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■you are taking a diuretic ■you have problems or serious side effects from taking pain relievers or fever reducers. **Ask a doctor or pharmacist before use if you are** ■under a doctor's care for any serious condition ■taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ■taking any other drug. **When using this product** ■take with food or milk if stomach upset occurs. **Stop use and ask a doctor if** ■you experience any of the following signs of stomach bleeding: ■feel faint ■vomit blood ■have bloody or black stools ■have stomach pain that does not get better ■you have symptoms of heart problems or stroke: ■chest pain ■trouble breathing ■weakness in one part or side of body ■slurred speech ■leg swelling ■pain gets worse or lasts more than 10 days ■fever gets worse or lasts more than 3 days ■you have difficulty swallowing ■it feels like the pill is stuck in your throat ■redness or swelling is present in the painful area ■any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. **Directions** ■do not take more than directed ■the smallest effective dose should be used ■drink a full glass of water with each dose. ■Adults and children 12 years and older: ■take 1 caplet every 8 to 12 hours while symptoms last ■for the first dose you may take 2 caplets within the first hour ■do not exceed 2 caplets in any 8 to 12 hour period ■do not exceed 3 caplets in a 24 hour period ■Children under 12 years: ask a doctor.

**Other information** ■each caplet contains: sodium 20 mg ■store at 20° to 25°C (68° to 77°F). Avoid high humidity and excessive heat above 40°C (104°F).  
**Inactive ingredients** colloidal silicon dioxide, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, and titanium dioxide. **Questions or comments?** Call 1-888-287-1915 M-F 9 AM-5 PM EST

P1433825 LM-5511

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 220 mg (90 Tablets Carton Label)**

**NDC 79903-232-90**

**equate™**

**Compare**

**to Aleve®  
Caplets  
active  
ingredient\*\***

**All Day  
Pain Relief**

**Naproxen Sodium  
Tablets USP, 220 mg  
(NSAID)**

**Pain Reliever/  
Fever Reducer  
Strength to  
Last 12 Hours**

**12  
HOURS**

**220  
mg  
EACH**

**90  
CAPLETS†**

**Actual Size**

**†Capsule-Shaped Tablets**



# NAPROXEN SODIUM

naproxen sodium tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79903-232
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>NAPROXEN SODIUM</b> (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>HYPROMELLOSE 2910 (5 MPA.S)</b> (UNII: R75537T0T4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 8000</b> (UNII: Q662QK8M3B)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	BLUE (light blue)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	K;45
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-232-90	1 in 1 CARTON	02/14/2024	
1		90 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205497	02/14/2024	

**Labeler** - WALMART INC. (051957769)

**Registrant** - Aurohealth LLC (078728447)

## Establishment

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Limited		650381903	ANALYSIS(79903-232) , MANUFACTURE(79903-232)

Revised: 4/2024

WALMART INC.