TIGER SUPPLY INC APF- sodium fluoride gel Tiger Supply Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Active Ingredient:

Fluoride Ion 1.23%

Available from 2.09% Sodium Fluoride and Hydrofluoric Acid

Purpose:

Fluoride Treatment Gel

Indications and Usage:

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Use Only. This product is not intended for home or unsupervised use.

Warnings:

- Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- Read directions carefully before using.

Dosage and Administration:

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

1. After thorough prophylaxsis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.

2. Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).

3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

Other Information:

Store at controlled room temperature 59° to 86°F (15°-30°C).

Inactive Ingredients:

Citric Acid, FD&C Red #40, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.

| | Active Ingredient: Purpose Fluoride Ion 1.23% Fluoride Treatment G Available from 2.09% Sodium Fluoride and Hydrofluoric Acid |
|--|---|
| SUPPLY INC. | Indications and Usage: • A stable thixotropic fluoride treatment gel used to help prevendental decay. • For Professional Office Use Only. This product is not intender for home or unsupervised consumer use. |
| APF GEL THIXOTROPIC | Warnings: Keep out of reach of children. Do not swallow. If product is accidentally swallowed i quantities greater than would normally occur with a treatmen gel, seek medical help or contact a Poison Control Center right away. Read directions carefully before using. |
| with Xylitol | Dosage and Administration: Shake well before use. This is a one minute or four minut fluoride gel for in-office patient use. It is normally used as preventative caries treatment two times a year. |
| Acidulated Phosphate Fluoride Treatment Gel 1.23% Fluoride Ion | After thorough prophylaxis, fill two single or one dual tra one third full with gel. Air dry teeth and insert trays into th mouth. Instruct patient to bite down lightly but firmly for one minu (80% effectiveness) or four minutes (100% effectiveness). |
| GLUTEN FREE | Remove trays. Instruct patient to expectorate any excergel and not to eat or drink for at least 30 minutes. Other Information: Store at controlled room temperature 59° to 86°F (15°-30°C |
| Ronly | Protect from freezing. Inactive Ingredients: Citric Acid, FD&C Red #40, Flavor, Magnesium Aluminu Silicate, Phosphoric Acid, Polysorbate 20, Sodium Benzoal Sodium Saccharin, Titanium Dioxide, Water, Xantham Gu Xylitol. |
| | Manufactured for: |
| Cherry Re-order#: TIG-24-45177 | 99 Sherwood Avenue Farmingdale, NY 11735 Tel: 631.293.2700 www.tigersupplyinc.co |
| IMPORTANT: READ DIRECTIONS FOR PROPER USE | |
| MADE IN USA Net wt. 16 oz (454 g) | 99092281 Rev 03/20 |

TIGER SUPPLY INC APF

sodium fluoride gel

Product Information

Product Type

HUMAN PRESCRIPTION DRUG

Item Code (Source)

| Route of Administ | ration DEN |
|--------------------------|------------|
|--------------------------|------------|

| Route of Administration | DENTAL | | | |
|---------------------------------|------------------------------------|------------------|--------------------------|----------------|
| | | | | |
| | | | | |
| Active Ingredient/Active | Moiety | | | |
| Ingredient Name Basis of Streng | | | Basis of Strength | Strength |
| SODIUM FLUORIDE (UNII: 8ZYQ1 | 474W7) (FLUORIDE ION - L | JNII:Q80VPU408O) | FLUORIDE ION | 5.6 g in 454 g |
| | | | | |
| | | | | |
| Inactive Ingredients | | | | |
| | Ingredient Name | • | | Strength |
| CITRIC ACID MONOHYDRATE (UI | NII: 2968PHW8QP) | | | |
| MAGNESIUM ALUMINUM SILICAT | re (UNII: 6M3P64V0NC) | | | |
| PHOSPHORIC ACID (UNII: E4GA88 | 84NN) | | | |
| POLYSORBATE 20 (UNII: 7T1F30V | ′5YH) | | | |
| SODIUM BENZOATE (UNII: OJ245 | SODIUM BENZOATE (UNII: OJ245FE5EU) | | | |
| SACCHARIN SODIUM (UNII: SB8Z | UX40TY) | | | |
| WATER (UNII: 059QF0KO0R) | | | | |
| XANTHAN GUM (UNII: TTV12P4NE | Ξ) | | | |
| XYLITOL (UNII: VCQ006KQ1E) | | | | |
| FD&C RED NO. 40 (UNII: WZ B912 | 7XOA) | | | |
| TITANIUM DIOXIDE (UNII: 15FIX9) | /2JP) | | | |
| | | | | |
| | | | | |
| Product Characteristics | | | | |
| Color | red | Score | | |
| Shape | | Size | | |
| Flavor | CHERRY | Imprint Code | | |
| Contains | | | | |
| | | | | |

| | - |
|-------|-------|
| Dack | DAIDA |
| Packa | |
| | - 33 |

| 7 | # Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|----------------------|---|-------------------------|-----------------------|
| : | NDC:69484- 051-15 | 454 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 03/01/2013 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------------|---|-------------------------|-----------------------|
| unapproved drug other | | 03/01/2013 | |
| | | | |

Labeler - Tiger Supply Inc (168720709)