

TRANSCEND- idoine solution IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

TRANSCEND SANITIZING 1% IODINE BARRIER TEAT DIP A PREMIUM, "TRUE" BARRIER TEAT DIP CONTAINING AN ADVANCED TRIPLE INGREDIENT EMOLLIENT SYSTEM. PROTECTS AGAINST BOTH CONTAGIOUS & ENVIRONMENTAL BACTERIA

ACTIVE INGREDIENT

Nonylphenoxyethoxyethanol Iodine Complex.....5%

INERT INGREDIENTS

Emollients.....12%

Other Inerts.....83%

(Includes Buffering Agents and Surfactants)

Total.....100%

pH of 5.5 at time of manufacture

This product contains highly effective levels of free iodine for fast kill of mastitis and killing organisms

INDICATIONS AND USAGE

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment
4. Wash Teat Dip cups after each milking.

RECOMMENDED PROCEDURES FOR TEAT SANITATION POST-MILKING (POST DIPPING)

Immediately after milking dip all teats full length with TRANSCEND Teat Dip Ensure good coverage of all sides of the teats and teat orifices.

WARNING

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES
CALL 1-800-424-9300**

TRANSCEND

- Unique "Drip-less" formulation.
- Forms a full barrier with plug.
- Forms a dry non-tacky film that lasts between milkings.
- Easily removed before milking.
- Compatible with popular pre-dip products.



TRANSCEND
SANITIZING 1% IODINE BARRIER TEAT DIP

A PREMIUM, "TRUE" BARRIER TEAT DIP CONTAINING
AN ADVANCED TRIPLE-INGREDIENT EMOLLIENT SYSTEM.
PROTECTS AGAINST BOTH CONTAGIOUS & ENVIRONMENTAL BACTERIA.

NDC # 29983-0032-4
**LESS DRIP
LESS WASTE**

ACTIVE INGREDIENT:
Nonylphenoxyethoxyethanol-Iodine Complex 5.00 %
(Provides 1.0% minimum Titratable Iodine)

INERT INGREDIENTS:

Emollients	12.00 %
Other Inerts	83.00 %
(Includes Buffering Agents and Surfactants)	
TOTAL:	100.00 %

pH of 5.5 at time of manufacture

This product contains highly effective levels of free iodine for fast kill of mastitis causing organisms.

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

RECOMMENDED PROCEDURES FOR TEAT SANITATION
POST-MILKING (POST-DIPPING)
Immediately after milking, dip all teats full length with TRANSCEND Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.



An Aid in the Prevention
Of New Intra-mammary
Infections

WARNING

DO NOT DILUTE THIS PRODUCT.
DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.
ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.
PROTECT THIS PRODUCT FROM FREEZING.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300**

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE:	LOT NO.:	NET CONTENTS: 5 GALLONS	IBA STOCK # 160121	600E
------------	----------	-------------------------	--------------------	------

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 27 PROVIDENCE ROAD, MILLBURY, MA 01527

TRANSCEND

idoine solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-0032
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	10 g in 1 L

Inactive Ingredients

Ingredient Name	Strength
MONOETHANOLAMINE (UNII: 5KV86114PT)	
WATER (UNII: 059QF0KOOR)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-0032-1	3.78 L in 1 DRUM		
2	NDC:29983-0032-4	18.9 L in 1 DRUM		
3	NDC:29983-0032-6	56.7 L in 1 DRUM		
4	NDC:29983-0032-7	113.4 L in 1 DRUM		
5	NDC:29983-0032-8	207.9 L in 1 DRUM		
6	NDC:29983-0032-9	1039.5 L in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/05/2010	

Labeler - IBA (019494160)

Revised: 3/2022

IBA