**AXIV ALLERGY- diphenhydramine hydrochloride capsule, liquid filled**
**VIVUNT LLC**

---

**AXIV Allergy**

***Drug Facts***

| ***Active Ingredient (in each softgel)*** | ***Purpose*** |
|---|---|
| Diphenhydramine HCl 25 mg | Antihistamine |

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
- runny nose
- sneezing

**Warnings**

**Do not use**

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Keep out of reach of children**

**Keep out of reach of children.**

**If pregnant or breast-feeding**

ask a health professional before use.

**Overdose Warning**

In case of overdose, get medical help or contact a Poison Control Center immediately.

**Directions**

- Take every 4 to 6 hours, or as directed by a doctor
- Do not take more than 12 Softgels in 24 hours for Adults and children 12 years and over. Do not take more than 6 Softgels for children 6 to under 12 years of age in 24 hours.

| | |
|---|---|
| adults and children 12 years and over | 1 or 2 Softgels |
| children 6 to under 12 years | 1 Softgel |
| children under 6 years | do not use |

**Other information**

- Store at 20° - 25 °C (68 °- 77 °F)
- Read all product information before using
- Tamper Evident: Do not use if the foil printed on the blister is torn or ripped.
- Tamper Evident: Do not use if carton or pouch is open

**Inactive ingredients**

Gelatin, Glycerin, Methylparaben, Polyethylene Glycol 400, Propylparaben, Purified Water, Sorbitol, Titanium Dioxide

**PRINCIPAL DISPLAY PANEL 24**

Compare to Benadryl® Allergy Liqui-Gels®

active ingredients*

NDC 82706-019-01

AXIV Antihistamine

- Runny Nose
- Sneezing
- Itchy, Watery Eyes
- Itchy Throat

DYE-FREE

Diphenhydramine HCl

24 SOFTGELS** **Liquid-filled capsules

Sorbitol, Titanium Dioxide.

*This product is not manufactured or distributed by Johnson & Johnson Consumer INC, owner of the registered trademarks Benadryl® Allergy Liqui-Gels®.

VIVUNT™ 30 YEARS

Distributed by: VIVUNT PHARMA LLC
8950 SW 74th Court, Suite 1901
Miami, FL 33156-3178
Made in India
www.vivunt.live

8 50047 78011 9

Compare to Benadryl® Allergy Liqui-Gels® active ingredients*
NDC 82706-019-01

Diphenhydramine HCl

DYE-FREE

24 SOFTGELS**
**Liquid-filled capsules

- Runny Nose
- Sneezing
- Itchy, Watery Eyes
- Itchy Throat

ANTIHISTAMINE

# Allergy Relief

OXIV

Allergy Relief

ANTIHISTAMINE

OXIV

## Drug Facts

**Active ingredient (in each softgel)** — **Purpose**
Diphenhydramine HCl 25 mg ..... Antihistamine

**Uses** Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ■ temporarily relieves these symptoms due to the common cold: ■ runny nose ■ sneezing

**Warnings**

**Do not use**
■ to make a child sleep ■ with any other product containing diphenhydramine, even one used on skin.

**Ask a doctor before use if you have** ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland.

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers.

**When using this product do not exceed recommended dose**
■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children.

**Keep out of reach of children.**

**If pregnant or breast-feeding,** ask a health professional before use.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center immediately.

**Directions**
■ take every 4 to 6 hours, or as directed by a doctor
■ do not take more than 12 softgels in 24 hours for adults and do not take more than 6 softgels for children 6 to 12 years of age in 24 hours.

| | |
|---|---|
| adults and children over 12 years | 1 or 2 softgels |
| children 6 to under 12 years | 1 softgel |
| children under 6 years | do not use |

**Other information**
■ Store at 20°-25°C (68°-77°F)
■ Read all product information before using
■ Tamper evident: Do not use it if the foil printed on the blister is torn or ripped.


**PRINCIPAL DISPLAY PANEL 6**

Compare to Benadryl® Allergy Liqui-Gels®

active ingredients*

NDC 82706-019-02

AXIV Antihistamine

Runny Nose
Sneezing
Itchy, Watery Eyes
Itchy Throat

DYE-FREE

Diphenhydramine HCl

6 SOFTGELS** **Liquid-filled capsules


Drug Facts
Active ingredient (in each caplet)
Purpose
Diphenhydramine HCl 25 mg
Antihistamine
Uses Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ■ temporarily relieves these symptoms due to the common cold: ■ runny nose ■ sneezing
Warnings
Do not use
■ to make a child sleep ■ with any other product containing diphenhydramine, even one used on skin.
Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland.
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.
When using this product do not exceed recommended dose
■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children.
Keep out of reach of children.
If pregnant or breast-feeding, ask a health professional before use.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center immediately.
Directions
■ take every 4 to 6 hours, or as directed by a doctor
■ do not take more than 12 caplets in 24 hours for adults and do not take more than 6 caplets for children 6 to 12 years of age in 24 hours.
■ adults and children over 12 years: 1 or 2 caplets
■ children 6 to under 12 years: 1 caplet
■ children under 6 years: do not use
Other information
■ Store at 20°-25°C (68°-77°F)
■ Read all product information before using
■ Tamper evident: Do not use if carton or pouch is open.
Inactive ingredients
Gelatin, Glycerin, Methylparaben, Polyethylene Glycol 400, Propylparaben, Purified Water, Sorbitol, Titanium Dioxide.
VIVUNT
30 Years
Distributed by:
VIVUNT PHARMA LLC
8950 SW 74th Court, Suite 1901
Miami, FL 33156-3178
Made in India.
www.vivunt.live
8 50047 78022 5
*This product is not manufactured or distributed by Johnson & Johnson Consumer INC, owner of the registered trademarks Benadryl® Allergy Ultratab®.
6 CAPLETS
ACTUAL SIZE
Diphenhydramine HCl
DYE-FREE
ANTIHISTAMINE
Allergy Relief

# AXIV ALLERGY

diphenhydramine hydrochloride capsule, liquid filled

## Product Information

| | | | |
|---|---|---|---|
| **Product Type** | HUMAN OTC DRUG | **Item Code (Source)** | NDC:82706-019 |
| **Route of Administration** | ORAL | | |

## Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|---|
| **DIPHENHYDRAMINE HYDROCHLORIDE** (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M) | DIPHENHYDRAMINE HYDROCHLORIDE | 25 mg |

## Inactive Ingredients

| Ingredient Name | Strength |
|---|---|
| **PROPYLPARABEN** (UNII: Z8IX2SC1OH) | |
| **GELATIN** (UNII: 2G86QN327L) | |
| **METHYLPARABEN** (UNII: A2I8C7HI9T) | |
| **WATER** (UNII: 059QF0KO0R) | |
| **GLYCERIN** (UNII: PDC6A3C0OX) | |
| **SORBITOL** (UNII: 506T60A25R) | |
| **POLYETHYLENE GLYCOL 400** (UNII: B697894SGQ) | |
| **TITANIUM DIOXIDE** (UNII: 15FIX9V2JP) | |

## Product Characteristics

| | | | |
|---|---|---|---|
| **Color** | white (Transparent) | **Score** | no score |
| **Shape** | OVAL | **Size** | 22mm |
| **Flavor** | | **Imprint Code** | AXIV |
| **Contains** | | | |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:82706-019-01 | 2 in 1 CARTON | 03/08/2024 | |
| **1** | | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| **2** | NDC:82706-019-02 | 3 in 1 CARTON | 03/08/2024 | |
| **2** | | 2 in 1 POUCH; Type 0: Not a Combination Product | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC Monograph Drug | M012 | 03/08/2024 | |

## Labeler - VIVUNT LLC (045829437)

Revised: 3/2024 VIVUNT LLC