

IODINE TINCTURE 7%- iodine liquid
ProActive Solutions USA, LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Iodine Tincture 7%

Iodine Tincture 7%

Use Instructions:

Topical Antiseptic

Use full Strength for superficial cuts, wounds, abrasions, insect bites and bruises on the skin

of animals. Apply iodine with a swab.

If necessary, clip hair around the area being treated and clean with soap and water.

Apply Iodine Tincture 7%TM only once daily. dilute product 3 to 1 if repeating application.

Do not apply under bandage.

Irritation may occur if used on tender skin areas. If redness, irritation, or swelling persists

or increases, discontinue use and consult a veterinarian.

Storage:

Store at 2-30 degrees C (36-86 degrees F).

Keep container away from heat and out of sunlight. Rinse empty container thoroughly and discard.

Hazard statement(s):

Toxic if swallowed. May cause eye irritation.

Precautionary statement(s):

Keep away from heat, sparks, open flames. No smoking. Do not eat, drink or smoke while

using this product. Wear protective clothing, and eye and face protection.

Hazards not otherwise classified: Flammable

Not for use on burns, deep cuts, or body cavities.

First Aid:

External: If irritation occurs, flush with plenty of cool, running water for at least 15 minutes

while removing contaminated clothing and shoes. Wash clothing before reuse.

Eyes: Immediately flush with plenty of cool, running water. Remove contact lenses; then continue flushing for at least 15 minutes, holding eyelids apart.

Internal: Call a physician immediately. If swallowed, do not induce vomiting. Rinse mouth, then immediately drink one or two large glasses of water or milk. Never give anything by mouth to an unconscious person.


Inhalation of vapors: Should not occur with proper use of this product.

Most important symptoms/effects, acute and delayed: None known

Immediate medical attention and special treatment needed, if any: If swallowed, give starch paste, milk, bread, egg white, or activated charcoal. A 5% solution of sodium

thiosulfate (photographic ("hypo")) may be administered orally at a rate of 10 ml per kilogram of body weight.

Iodine Tincture 7%



Hazard statement(s):
Toxic if swallowed. May cause eye irritation.
Precautionary statement(s):
Keep away from heat, sparks, open flames. No smoking. Do not eat, drink or smoke while using this product. Wear protective clothing, and eye and face protection.
Hazards not otherwise classified: Flammable

Not for use on burns, deep cuts, or body cavities.

First Aid:
External: If irritation occurs, flush with plenty of cool, running water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.
Eyes: Immediately flush with plenty of cool, running water. Remove contact lenses; then continue flushing for at least 15 minutes, holding eyelids apart.
Internal: Call a physician immediately. If swallowed, do not induce vomiting. Rinse mouth, then immediately drink one or two large glasses of water or milk. Never give anything by mouth to an unconscious person.
Inhalation of vapors: Should not occur with proper use of this product.
Most important symptoms/effects, acute and delayed: None known
Immediate medical attention and special treatment needed, if any: If swallowed, give starch paste, milk, bread, egg white, or activated charcoal. A 5% solution of sodium thiosulfate (photographic ("hypo")) may be administered orally at a rate of 10 ml per kilogram of body weight.

Iodine Tincture™

Active Ingredients: Titratable Iodine.....7.00%
Isopropyl Alcohol.....65.00%

FOR FARM OR INDUSTRIAL USE ONLY -
KEEP OUT OF REACH OF CHILDREN



Use Instructions:
Topical Antiseptic
Use full strength for superficial cuts, wounds, abrasions, insect bites and bruises on the skin of animals. Apply iodine with a swab.
If necessary, clip hair around the area being treated and clean with soap and water.
Apply Iodine Tincture 7%™ only once daily. Dilute product 3 to 1 if repeating application.
Do not apply under bandage.
Irritation may occur if used on tender skin areas. If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian.

Storage:
Store at 2-30 degrees C (36-86 degrees F).

Keep container away from heat and out of sunlight. Rinse empty container thoroughly and discard.

Net Contents: 5 gal. 15 gal. 55 gal. 275 gal. 310 gal. 330 gal.
Case Contents: 4x1 gal (total: 4 gal.)



FOR CHEMICAL EMERGENCIES:
CALL CHEMTREC 1-800-424-9300



ProActive Solutions® USA, LLC • 301 Bridge Street • Green Bay, WI 54303
800.279.7761 www.proactivesolutionsusa.com



UN1293
TINCTURES, MEDICINAL, 3, II
(IODINE, ISOPROPYL ALCOHOL)



REV Date: 2/9/2026



Hazard statement(s):
Toxic if swallowed. May cause eye irritation.
Precautionary statement(s):
Keep away from heat, sparks, open flames. No smoking. Do not eat, drink or smoke while using this product. Wear protective clothing, and eye and face protection.
Hazards not otherwise classified: Flammable

Not for use on burns, deep cuts, or body cavities.

First Aid:
External: If irritation occurs, flush with plenty of cool, running water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.
Eyes: Immediately flush with plenty of cool, running water. Remove contact lenses; then continue flushing for at least 15 minutes, holding eyelids apart.
Internal: Call a physician immediately. If swallowed, do not induce vomiting. Rinse mouth, then immediately drink one or two large glasses of water or milk. Never give anything by mouth to an unconscious person.
Inhalation of vapors: Should not occur with proper use of this product.
Most important symptoms/effects, acute and delayed: None known
Immediate medical attention and special treatment needed, if any: If swallowed, give starch paste, milk, bread, egg white, or activated charcoal. A 5% solution of sodium thiosulfate (photographic ("hypo")) may be administered orally at a rate of 10 ml per kilogram of body weight.

Iodine Tincture™

Active Ingredients: Titratable Iodine.....7.00%
Isopropyl Alcohol.....65.00%

FOR FARM OR INDUSTRIAL USE ONLY -
KEEP OUT OF REACH OF CHILDREN



Use Instructions:
Topical Antiseptic
Use full strength for superficial cuts, wounds, abrasions, insect bites and bruises on the skin of animals. Apply iodine with a swab.
If necessary, clip hair around the area being treated and clean with soap and water.
Apply Iodine Tincture 7%™ only once daily. Dilute product 3 to 1 if repeating application.
Do not apply under bandage.
Irritation may occur if used on tender skin areas. If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian.

Storage:
Store at 2-30 degrees C (36-86 degrees F).

Keep container away from heat and out of sunlight. Rinse empty container thoroughly and discard.

Net Contents: 5 gal. 15 gal. 55 gal. 275 gal. 310 gal. 330 gal.
Case Contents: 4x1 gal (total: 4 gal.)




FOR CHEMICAL EMERGENCIES:
CALL CHEMTREC 1-800-424-9300



ProActive Solutions® USA, LLC • 301 Bridge Street • Green Bay, WI 54303
800.279.7761 www.proactivesolutionsusa.com



UN1293
TINCTURES, MEDICINAL, 3, II
(IODINE, ISOPROPYL ALCOHOL)



REV Date: 2/9/2026



FLAMMABLE
(Toxic if Swallowed)



Hazard statement(s):
Toxic if swallowed. May cause eye irritation.
Precautionary statement(s):
Keep away from heat, sparks, open flames. No smoking. Do not eat, drink or smoke while using this product. Wear protective clothing, and eye and face protection.
Hazards not otherwise classified: Flammable
Not for use on burns, deep cuts, or body cavities.

First Aid:
External: If irritation occurs, flush with plenty of cool, running water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.
Eyes: Immediately flush with plenty of cool, running water. Remove contact lenses; then continue flushing for at least 15 minutes, holding eyelids apart.
Internal: Call a physician immediately. If swallowed, do not induce vomiting. Rinse mouth, then immediately drink one or two large glasses of water or milk. Never give anything by mouth to an unconscious person.
Inhalation of vapors: Should not occur with proper use of this product.
Most important symptoms/effects, acute and delayed: None known
Immediate medical attention and special treatment needed, if any: If swallowed, give starch paste, milk, bread, egg white, or activated charcoal. A 5% solution of sodium thiosulfate [photographic ("hypo")] may be administered orally at a rate of 10 ml per kilogram of body weight.

Use Instructions:
Topical Antiseptic
Use full Strength for superficial cuts, wounds, abrasions, insect bites and bruises on the skin of animals. Apply iodine with a swab.
If necessary, clip hair around the area being treated and clean with soap and water.
Apply Iodine Tincture 7%™ only once daily. dilute product 3 to 1 if repeating application. Do not apply under bandage.
Irritation may occur if used on tender skin areas. If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian.

Storage:
Store at 2-30 degrees C (36-86 degrees F).
Keep container away from heat and out of sunlight. Rinse empty container thoroughly and discard.

Net Contents: 5 gal. 15 gal. 55 gal. 275 gal. 310 gal. 330 gal.
Case Contents: 4x1 gal (total: 4 gal.)




FOR CHEMICAL EMERGENCIES:
CALL CHEMTREC 1-800-424-9300



ProActive Solutions® USA, LLC • 301 Bridge Street • Green Bay, WI 54303
800.279.7761 www.proactivesolutionsusa.com



UN1293
TINCTURES, MEDICINAL, 3. II
(IODINE, ISOPROPYL ALCOHOL)



REV Date: 2/9/2026

IODINE TINCTURE 7%

iodine liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:63927-4240
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	7 kg in 100 kg
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	65 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63927-4240-1	13.94 kg in 1 CASE		
2	NDC:63927-4240-2	17.42 kg in 1 JUG		
3	NDC:63927-4240-3	52.26 kg in 1 DRUM		
4	NDC:63927-4240-4	191.63 kg in 1 DRUM		
5	NDC:63927-4240-5	958.13 kg in 1 CONTAINER, FLEXIBLE INTERMEDIATE BULK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/12/2008	

Labeler - ProActive Solutions USA, LLC (029368225)

Registrant - ProActive Solutions USA, LLC (029368225)

Establishment

Name	Address	ID/FEI	Business Operations
ProActive Solutions USA, LLC		029368225	api manufacture, manufacture, pack, repack, label

Revised: 2/2026

ProActive Solutions USA, LLC