

**IBUPROFEN- ibuprofen capsule, liquid filled  
Family Dollar (FAMILY WELLNESS)**

-----  
**Drug Facts**

**Active ingredient (in each capsule)**

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - slurred speech
  - trouble breathing
  - leg swelling
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

## **Other information**

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F).
- swallow whole, do not crush, chew, or dissolve

## **Inactive ingredients**

FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol solution, white ink

## **Questions or comments?**

**Call 1-877-753-3935** Monday-Friday 9AM-5PM EST

## **Principal Display Panel**

COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS®†

## **Ibuprofen**

### **Capsules, 200 mg**

Pain Reliever/Fever Reducer **(NSAID)**

SOFTGELS\*\*

(\*\* LIQUID FILLED CAPSULES)

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**DISTRIBUTED BY: MIDWOOD BRANDS, LLC**

10611 MONROE RD., MATTHEWS, NC 28105

**Product Label**

**COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS™**

**FAMILY wellness**

NDC 55319-745-16

**Ibuprofen**  
Capsules, 200 mg  
Pain Reliever / Fever Reducer (NSAID)

**160 SOFTGELS\*\***  
(\*\*\*LIQUID-FILLED CAPSULES)

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**Drug Facts**

**Active ingredient (in each capsule)**  
Sublabeled ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt).....

**Purpose**  
Pain reliever/  
fever reducer

**Uses**  
Temporarily relieves minor aches and pains due to:  
■ headache  
■ muscular aches  
■ minor pain of arthritis  
■ toothache  
■ the common cold  
■ menstrual cramps  
Temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives  
■ facial swelling  
■ asthma (wheezing)  
■ shock  
■ skin redness  
■ rash  
■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery

**Ask a doctor before use if**  
■ the stomach bleeding warning applies to you  
■ you have problems or serious side effects from taking pain relievers or fever reducers  
■ you have a history of stomach problems, such as heartburn

**Drug Facts (continued under label)**  
This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

SKU 999373

DISTRIBUTED BY: MIDWOOD BRANDS, LLC, 10611 MONROE RD. MATTHEWS, NC 28105

NOT 100% SATISFIED? Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

PLD-45248 L8006560 Lot No.: Exp. Date:

0 32251 48324 0

PEEL HERE →

**Drug Facts (continued)**

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin  
■ taking any other drug

**When using this product:** Take with food or milk if stomach upset occurs.

**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint  
■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain  
■ slurred speech  
■ trouble breathing  
■ leg swelling  
■ weakness in one part or side of body  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
■ do not take more than directed  
■ the smallest effective dose should be used  
■ adults and children 12 years and over:  
■ take 1 capsule every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 capsule, 2 capsules may be used  
■ do not exceed 6 capsules in 24 hours, unless directed by a doctor  
■ children under 12 years: ask a doctor

**Other information**  
■ each capsule contains potassium 20 mg  
■ read all warnings and directions before use  
■ store between 20° to 25°C (68° to 77°F)  
■ avoid excessive heat above 40°C (104°F)  
■ swallow whole; do not crush, chew, or dissolve

**Inactive ingredients**  
FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution, white ink

**Questions or comments?**  
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**FAMILY WELLNESS Ibuprofen Capsules 200 mg**

**IBUPROFEN**

ibuprofen capsule, liquid filled

**Product Information**

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:55319-745 |
| <b>Route of Administration</b> | ORAL           |                           |               |

**Active Ingredient/Active Moiety**

| Ingredient Name | Basis of Strength | Strength |
|-----------------|-------------------|----------|
|-----------------|-------------------|----------|

|   |   |   |                             |                           |
|---|---|---|-----------------------------|---------------------------|
| <b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) |   | IBUPROFEN   | 200 mg                      |                           |
| <b>Inactive Ingredients</b>                                       |   |   |                             |                           |
| <b>Ingredient Name</b>  |   |   | <b>Strength</b>             |                           |
| FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)                               |   |   |                             |                           |
| GELATIN (UNII: 2G86QN327L)  |   |   |                             |                           |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)               |   |   |                             |                           |
| POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)                            |   |   |                             |                           |
| WATER (UNII: 059QF0KO0R)  |   |   |                             |                           |
| SORBITOL (UNII: 506T60A25R)                                       |   |   |                             |                           |
| SORBITAN (UNII: 6O92ICV9RU)                                       |   |   |                             |                           |
| MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)                     |   |   |                             |                           |
| <b>Product Characteristics</b>                                    |   |   |                             |                           |
| <b>Color</b>  | blue  | <b>Score</b>  | no score                    |                           |
| <b>Shape</b>  | CAPSULE   | <b>Size</b>   | 19mm                        |                           |
| <b>Flavor</b>   |   | <b>Imprint Code</b>   | PC16                        |                           |
| <b>Contains</b>   |   |   |                             |                           |
| <b>Packaging</b>  |   |   |                             |                           |
| <b>#</b>  | <b>Item Code</b>                                | <b>Package Description</b>                                  | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
| 1   | NDC:55319-745-30                                | 1 in 1 BOX  | 01/31/2018                  | 01/31/2026                |
| 1   |   | 30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product  |                             |                           |
| 2   | NDC:55319-745-80                                | 1 in 1 BOX  | 01/31/2018                  | 01/31/2026                |
| 2   |   | 80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product  |                             |                           |
| 3   | NDC:55319-745-16                                | 160 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 01/31/2018                  | 01/31/2026                |
| <b>Marketing Information</b>                                      |   |   |                             |                           |
| <b>Marketing Category</b>   | <b>Application Number or Monograph Citation</b> | <b>Marketing Start Date</b>                                 | <b>Marketing End Date</b>   |                           |
| ANDA  | ANDA206568                                      | 01/31/2018  | 01/31/2026                  |                           |

**Labeler** - Family Dollar (FAMILY WELLNESS) (024472631)