

**TUSSIN COUGH AND CHEST CONGESTION- dextromethorphan hydrobromide,
guaifenesin solution
Rite Aid Corporation**

Rite Aid Corporation Tussin Cough + Chest Congestion Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Purposes

Cough suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- **each 20 mL contains:** sodium 14 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

SEE NEW DOSING

Compare to the active ingredients of Robitussin[®] Cough + Chest Congestion DM

FREE FROM

GLUTEN FREE

ALCOHOL FREE

TUSSIN

COUGH + CHEST CONGESTION

ADULT NON-DROWSY

DEXTROMETHORPHAN HBr

COUGH SUPPRESSANT


GUAIFENESIN/EXPECTORANT

Relieves cough, chest congestion & mucus

For ages 12 & over

DM

8 FL OZ (237 mL)



Drug Facts

Active ingredients Purposes (in each 20 mL)

Dextromethorphan HBr, USP 20 mg.....Cough suppressant
Guaifenesin, USP 200 mg.....Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product

Drug Facts (continued)

- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 14 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Cough + Chest Congestion DM.

OUR PROMISE:



DISTRIBUTED BY:
RITE AID
30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com

SATISFACTION GUARANTEE

If you're not satisfied, we'll happily refund your money.

TUSSIN COUGH + CHEST CONGESTION

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

NDC 11822-1630-1

SEE NEW DOSING

Compare to the active ingredients of Robitussin® Cough + Chest Congestion DM*

FREE FROM
GLUTEN FREE
ALCOHOL FREE

TUSSIN COUGH + CHEST CONGESTION

ADULT
NON-DROWSY

DEXTROMETHORPHAN HBr
COUGH SUPPRESSANT
GUAIFENESIN/EXPECTORANT

Relieves cough,
chest congestion
& mucus
For ages 12 & over

DM



8 FL OZ (237 mL)



63K34 83 C1

NDC 11822-1630-1

SEE NEW DOSING

Compare to the active ingredients of Robitussin® Cough + Chest Congestion DM*

FREE FROM
GLUTEN FREE
ALCOHOL FREE

TUSSIN COUGH + CHEST CONGESTION

ADULT
NON-DROWSY

DEXTROMETHORPHAN HBr
COUGH SUPPRESSANT
GUAIFENESIN/EXPECTORANT

Relieves cough,
chest congestion
& mucus
For ages 12 & over

DM



8 FL OZ (237 mL)

TUSSIN COUGH AND CHEST CONGESTION

dextromethorphan hydrobromide, guaifenesin solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-1630
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	RED	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-1630-0	1 in 1 CARTON	06/23/2022	11/30/2026
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-1630-1	1 in 1 CARTON	06/08/2022	01/31/2026
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/08/2022	11/30/2026

Labeler - Rite Aid Corporation (014578892)

Revised: 1/2026

Rite Aid Corporation