

**DAYTIME NIGHTTIME COLD FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl  
EQUALINE (SuperValu)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**DRUG FACTS**

**Active ingredients for Daytime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Active ingredients for Nighttime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose for Daytime**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Purpose for Nighttime**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

**DAYTIME**

- temporarily relieves common cold and flu symptoms
  - cough due to minor throat and bronchial irritation
  - nasal congestion
  - headache
  - minor aches and pains

- fever
- sore throat

## **NIGHTTIME**

- temporarily relieves common cold and flu symptoms
  - cough due to minor throat and bronchial irritation
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing

## **Warnings**

### **DAYTIME**

**Liver warning:** These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using these products

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **NIGHTTIME**

**Liver warning:** This product contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or following by fever, headache, rash, nausea, vomiting, consult a doctor promptly.

## **Do not use**

### **DAYTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **NIGHTTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## **Ask a doctor before use if you have**

### DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

### NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

## **Ask a doctor or pharmacist before use if you are**

### DAYTIME

taking the blood thinning drug warfarin

### NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

## **When using this product,**

## **DAYTIME**

**do not exceed recommended dosage**

## **NIGHTTIME**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

### DAYTIME

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash headache that lasts.

These could be a signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

### DAYTIME

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over; take 2 softgels with water every 4 hours.

- children under 12 years: do not use

#### NIGHTTIME

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

#### Other information

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

#### Inactive ingredients

##### DAYTIME

FD&C yellow #6, FD&C Red #40, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

##### NIGHTTIME

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

#### Questions or comments?

Call toll free 1-855-423-2630

#### Principal Display Panel

compare to Vicks® Dayquil® & NuQuil® Cold & Flu LiquiCaps® active ingredient†

multi-symptom

daytime

cold & flu relief

acetaminophen 325 mg

(aches/fever reducer)

dextromethorphan HBr 10 mg

(cough suppressant)

phenylephrine HCL 5 mg

(nasal decongestant)

non-drowsy

relieves

- aches, fever, sore throat
- nasal congestion & cough

ALCOHOL FREE

ANTIHISTAMINE FREE

softgels\*\*

\*\*liquid-filled capsules)

multi-symptom

**nighttime**

cold & flu relief

acetaminophen 325 mg

(aches/fever reducer)

dextromethorphan HBr 15 mg

(cough suppressant)

doxylamine succinate 6.25 mg

(antihistamine)

relieves

- Aches, Fever, Sore Throat
- cough sneezing & runny nose

softgels

\*\*liquid-fill capsules

when using Daytime and nighttime product, carefully read the labeling to ensure correct dosing.

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil, NyQuil®, and LiquiCaps® are registered trademarks of the Procter and Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY UNFI, PROVIDENCE, RI 02908 USA

**Product Label**

**Daytime Cold & Flu Softgel**

**Drug Facts (continued)**

**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if** ■ nervousness, dizziness, or sleepiness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children. Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 4 hours ■ children under 12 years: do not use

**Nighttime Cold & Flu Softgel**

**Drug Facts (continued)**

**Ask a doctor before use if you have** ■ liver disease ■ gout ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

**When using this product** ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ do not take more than 4 doses in 24 hours ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use

**Daytime Cold & Flu Softgel**

**Drug Facts**

**Active ingredients (in each softgel)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
 Dextromethorphan HBr 10 mg.....Cough suppressant  
 Pseudoephedrine HCl 5 mg.....Nasal decongestant

**Purposes**

■ temporarily relieves common cold and flu symptoms: ■ headache ■ minor aches and pains ■ fever ■ sore throat ■ cough due to minor throat and bronchial irritation ■ nasal congestion

**Uses**

■ temporarily relieves common cold and flu symptoms: ■ headache ■ minor aches and pains ■ fever ■ sore throat ■ cough due to minor throat and bronchial irritation ■ nasal congestion

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

**If a skin reaction occurs, stop use and seek medical help right away.**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Nighttime Cold & Flu Softgel**

**Drug Facts**

**Active ingredients (in each softgel)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
 Dextromethorphan HBr 15 mg.....Cough suppressant  
 Doxylamine succinate 6.25 mg.....Antihistamine

**Purposes**

■ temporarily relieves common cold and flu symptoms: ■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing ■ cough due to minor throat and bronchial irritation ■ sore throat

**Uses**

■ temporarily relieves common cold and flu symptoms: ■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing ■ cough due to minor throat and bronchial irritation ■ sore throat

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

**If a skin reaction occurs, stop use and seek medical help right away.**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**EQUALINE®**

multi-symptom  
**daytime**  
 cold & flu relief  
 acetaminophen 325 mg  
 (aches/fever reducer)

compare to Vicks®  
 DayQuil® & NyQuil®  
 Cold & Flu LiquiCaps®  
 active ingredients\*

NDP-106-69117 CDG

multi-symptom  
**nighttime**  
 cold & flu relief  
 acetaminophen 325 mg  
 (aches/fever reducer)

<p>dextromethorphan HBr 10 mg (cough suppressant) phenylephrine HCl 5 mg (nasal decongestant) <b>non-drowsy</b> relieves: • aches, fever &amp; sore throat • nasal congestion &amp; cough</p> <p>ALCOHOL FREE ANTIHISTAMINE FREE</p> <p><b>32 daytime softgels**</b> **liquid-filled capsules</p>  <p>actual size</p>	<p>dextromethorphan HBr 15 mg (cough suppressant) doxylamine succinate 6.25 mg (antihistamine) relieves: • aches, fever &amp; sore throat • cough, sneezing &amp; runny nose</p> <p>When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.</p> <p><b>16 nighttime softgels**</b> **liquid-filled capsules</p>  <p>actual size</p>
--	---



**PARENTS:**  
Learn about your medicine here  
[www.StepMedicine.com](http://www.StepMedicine.com)

Like it or let us make it right.  
855-423-2630  
That's our quality promise.



DISTRIBUTED BY UNF, PROVIDENCE, RI 02908 USA

KEEP OUTER CAPTION FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

TAMPER EVIDENT: DO NOT USE IF CAPTION IS OPENED OR IF BULSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks, DayQuil, NyQuil, and LiquiCaps are registered trademarks of The Procter and Gamble Company.

PLD-F5888  
FC007706  
Product of China

<p><b>Daytime Cold &amp; Flu Softgel</b></p> <p><b>Drug Facts (continued)</b></p> <p>Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat</p> <p>Inactive ingredients FD&amp;C red #40, FD&amp;C yellow #6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p>Questions or comments? Call toll free 1-855-423-2630</p>	<p><b>Nighttime Cold &amp; Flu Softgel</b></p> <p><b>Drug Facts (continued)</b></p> <p>Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat</p> <p>Inactive ingredients D&amp;C yellow #10, FD&amp;C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p>Questions or comments? Call toll free 1-855-423-2630</p>
--	--

## EQUALINE Multi-Symptom Daytime Nighttime Cold & Flu Relief

### DAYTIME NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41163-901
---------------------	----------------	---------------------------	---------------

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-901-48	1 in 1 KIT; Type 0: Not a Combination Product	07/30/2021	

#### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
--------	------------------	------------------------



<b>Part 1</b>	16 BLISTER PACK	16
<b>Part 2</b>	32 BLISTER PACK	32

## Part 1 of 2

### NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

#### Product Information

**Item Code (Source)** NDC:41163-899

**Route of Administration** ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

#### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	

#### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC10
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		16 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/30/2021	

## Part 2 of 2

### DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

## Product Information

Item Code (Source)	NDC:41163-998
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O921CV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

LIGHT MINERAL OIL (UNII: N6K5787QVP)

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC9
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		32 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/30/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/30/2021	

**Labeler** - EQUALINE (SuperValu) (006961411)

Revised: 11/2022

EQUALINE (SuperValu)