

# DEFENCATH- taurolidine and heparin solution CorMedix Inc.

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DEFENCATH safely and effectively. See full prescribing information for DEFENCATH.

**DEFENCATH® (taurolidine and heparin) catheter lock solution, for central venous catheter instillation use**

**Initial U.S. Approval: 2023**

**LIMITED POPULATION**

## RECENT MAJOR CHANGES

Dosage and Administration (2)

12/2024

## INDICATIONS AND USAGE

**LIMITED POPULATION:** DEFENCATH is a combination of taurolidine, a thiadiazinane antimicrobial, and heparin, an anti-coagulant, indicated to reduce the incidence of catheter-related bloodstream infections (CRBSI) in adult patients with kidney failure receiving chronic hemodialysis (HD) through a central venous catheter (CVC). This drug is indicated for use in a limited and specific population of patients. (1)

### Limitations of Use

The safety and effectiveness of DEFENCATH have not been established for use in populations other than adult patients with kidney failure receiving chronic HD through a CVC. (1)

## DOSAGE AND ADMINISTRATION

- DEFENCATH is for instillation into CVCs only. (2)
- DEFENCATH is **not** intended for systemic administration. (2)
- Do **not** use DEFENCATH as a catheter lock flush product. (2)
- Withdraw a sufficient volume of DEFENCATH catheter lock solution (CLS) from the vial using a sterile needle and syringe to fill the catheter lumens. (2)
- Use 3 mL or 5 mL single dose vial (depending on the volume of the catheter lumens) to instill DEFENCATH into each catheter lumen at the conclusion of each HD session. (2)
- DEFENCATH should be aspirated from the catheter and discarded prior to the initiation of the next HD session. (2)
- Discard any unused portion of DEFENCATH remaining in the vial. (2)

## DOSAGE FORMS AND STRENGTHS

- DEFENCATH is a sterile catheter lock solution available in single-dose vials in the following strengths:
  - 3 mL containing taurolidine 40.5 mg/3 mL (13.5 mg/mL), and heparin 3,000 USP Units/3 mL (1,000 USP Units/mL) (3)
  - 5 mL containing taurolidine 67.5 mg/5 mL (13.5 mg/mL), and heparin 5,000 USP Units/5 mL (1,000 USP Units/mL) (3)

## CONTRAINDICATIONS

- Known heparin-induced thrombocytopenia. (4)
- Known hypersensitivity to taurolidine, heparin or the citrate excipient (components of DEFENCATH), or pork products. (4)

## WARNINGS AND PRECAUTIONS

- Heparin-Induced Thrombocytopenia (HIT): HIT was reported in Trial 1 at an incidence rate of 0.3% in patients using heparin, a component of DEFENCATH as a CLS. If HIT occurs, discontinue DEFENCATH and institute appropriate supportive measures. (5.1)
- Drug Hypersensitivity Reactions: Drug hypersensitivity reactions were reported in Trial 1 at an incidence rate of 0.5% in patients using heparin, a component of DEFENCATH, as a CLS. If a hypersensitivity reaction occurs, discontinue DEFENCATH and institute appropriate supportive measures. (5.2)

## ADVERSE REACTIONS

The most frequently reported adverse reactions occurring in greater than or equal to 2% of patients in Trial

1 using DEFENCATH as a CLS were hemodialysis catheter malfunction, hemorrhage/bleeding, nausea, vomiting, dizziness, musculoskeletal chest pain, and thrombocytopenia. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact CorMedix Inc at 1-844-424-6345 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**Revised: 12/2024**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

5.1 Heparin-Induced Thrombocytopenia

5.2 Drug Hypersensitivity

### **6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

### **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.4 Microbiology

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **14 CLINICAL STUDIES**

### **15 REFERENCES**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

LIMITED POPULATION: DEFENCATH® is indicated to reduce the incidence of catheter-related bloodstream infections (CRBSI) in adult patients with kidney failure receiving

chronic hemodialysis (HD) through a central venous catheter (CVC). This drug is indicated for use in a limited and specific population of patients [see *Clinical Studies (14)*].

### Limitations of Use

The safety and effectiveness of DEFENCATH have not been established for use in populations other than adult patients with kidney failure receiving chronic HD through a CVC.

## **2 DOSAGE AND ADMINISTRATION**

DEFENCATH is for instillation into CVCs only [see *Indications and Usage (1)*]. DEFENCATH is **not** intended for systemic administration. Do **not** use DEFENCATH as a catheter lock flush product.

Withdraw a sufficient volume of DEFENCATH catheter lock solution (CLS) from the vial using a sterile needle and syringe to fill the catheter lumens. Use 3 mL or 5 mL single-dose vial (depending on the volume of the catheter lumens) to instill DEFENCATH into each catheter lumen at the conclusion of each HD session for patients with kidney failure requiring chronic HD. Prior to initiation of the next HD session, DEFENCATH should be aspirated from the catheter and discarded. If DEFENCATH cannot be aspirated, continue with standard of care CVC preparation and flush with normal saline. If a catheter malfunction is suspected, appropriate standard of care measures should be instituted.

Each DEFENCATH single-dose vial is designed for use with a single patient as a single instillation in the CVC. Discard any unused portion of DEFENCATH remaining in the vial.

## **3 DOSAGE FORMS AND STRENGTHS**

DEFENCATH CLS is available as a sterile, preservative-free, clear, aqueous-based solution in the following strengths:

- 3 mL of catheter lock solution in a single-dose vial containing taurolidine 40.5 mg/3 mL (13.5 mg/mL), and heparin 3,000 USP Units/3 mL (1,000 USP Units/mL)
- 5 mL of catheter lock solution in a single-dose vial containing taurolidine 67.5 mg/5 mL (13.5 mg/mL), and heparin 5,000 USP Units/5 mL (1,000 USP Units/mL)

## **4 CONTRAINDICATIONS**

DEFENCATH is contraindicated in patients with:

- Known heparin-induced thrombocytopenia (HIT) [see *Warnings and Precautions (5.1)*].
- Known hypersensitivity to taurolidine, heparin or the citrate excipient (components of DEFENCATH) or pork products [see *Warnings and Precautions (5.2)*].

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Heparin-Induced Thrombocytopenia**

Heparin-induced thrombocytopenia (HIT) was reported at an incidence rate of 0.3% in

Trial 1 in patients using heparin, a component of DEFENCATH, as a CLS. If HIT occurs, discontinue DEFENCATH and institute appropriate supportive measures [see *Contraindications (4)*].

## **5.2 Drug Hypersensitivity**

Drug hypersensitivity reactions were reported at an incidence rate of 0.5% in Trial 1 in patients using heparin, a component of DEFENCATH, as a CLS. If a hypersensitivity reaction occurs, discontinue DEFENCATH and institute appropriate supportive measures [see *Contraindications (4)*].

## **6 ADVERSE REACTIONS**

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Heparin-Induced Thrombocytopenia [see *Warnings and Precautions (5.1)*]
- Drug Hypersensitivity [see *Warnings and Precautions (5.2)*]

### **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### Overview of the Safety Evaluation of DEFENCATH

The safety of DEFENCATH was evaluated in a single, active-controlled, double-blind, randomized Phase 3 trial, LOCK-IT-100 (referred to as Trial 1), in adult patients with kidney failure receiving chronic HD through a CVC. In Trial 1, the safety population was comprised of 797 patients with 398 patients using DEFENCATH as a CLS and 399 using heparin as a CLS. Patients used DEFENCATH as a CLS for a mean duration of 173 days, and a range of 4 to 863 days.

The mean age of the patients in Trial 1 was 61 years, 58% of patients identified as male (42% patients identified as female), 63% of patients identified as white and 30% identified as black or African-American. Patients with diabetes at baseline accounted for 70% of patients and the mean body mass index (BMI) of patients in the study was 30. The majority of patients (68%) had been receiving HD for 12 months or less. The location of the CVC was the jugular vein in 92% of patients.

#### Serious Adverse Reactions and Adverse Reactions Leading to Discontinuation

In Trial 1, serious adverse reactions occurred in 40% (159/398) of patients receiving DEFENCATH as a CLS and 42% (167/399) of patients receiving heparin as a CLS. Adverse reactions leading to death occurred in 5% (18/398) of patients in the DEFENCATH arm and 5% (21/399) of patients in the heparin arm.

Adverse reactions leading to discontinuation of study drug occurred in 17% (69/398) of patients in the DEFENCATH arm and 18% (72/399) of patients in the heparin arm.

#### Common Adverse Reactions

In Trial 1, adverse reactions were reported in 79% (314/398) of patients using DEFENCATH as a CLS and 79% (315/399) of patients using heparin.

Table 1 lists selected adverse reactions occurring in  $\geq 2\%$  of patients using DEFENCATH or heparin in Trial 1.

**Table 1: Selected Adverse Reactions Occurring in Greater Than or Equal to 2% of Patients Receiving DEFENCATH in Trial 1**

<b>Adverse Reactions</b>	<b>DEFENCATH (N=398) N (%)</b>	<b>Heparin (N=399) N (%)</b>
<b>Product Issues</b>		
Hemodialysis catheter malfunction*	68 (17)	47 (12)
<b>Blood and lymphatic system disorders</b>		
Hemorrhage/bleeding†	27 (7)	34 (9)
Thrombocytopenia	7 (2)	4 (1)
<b>Gastrointestinal disorders</b>		
Nausea‡	28 (7)	44 (11)
Vomiting§	24 (6)	32 (8)
<b>Nervous system disorders</b>		
Dizziness	22 (6)	16 (4)
<b>Musculoskeletal and Connective Tissue Disorders</b>		
Musculoskeletal chest pain¶	11 (3)	7 (2)

\* Grouped term includes device malfunction

† Grouped term includes: hematoma/hemorrhage, intracranial hemorrhage, arteriovenous graft or fistula site hemorrhage, catheter site hematoma/hemorrhage, cerebral/cerebellar hemorrhage, eye/conjunctival hemorrhage, gastrointestinal hemorrhage, hematuria, hemobilia, hemoptysis, renal hematoma, vaginal/uterine hemorrhage

‡ Grouped term includes nausea and procedural nausea

§ Grouped term includes vomiting and procedural vomiting

¶ Grouped term includes non-cardiac chest pain, musculoskeletal chest pain, chest discomfort

Table 2 lists the proportion of patients in Trial 1 that developed loss of catheter patency defined as requiring use of a thrombolytic agent to resolve catheter thrombosis or removal of the catheter due to malfunction/dysfunction.

**Table 2: Loss of Catheter Patency in Trial 1**

	<b>DEFENCATH (N=398) N (%)</b>	<b>Heparin (N=399) N (%)</b>
<b>Loss of Catheter Patency</b>	63 (16)	48 (12)
Use of a Thrombolytic Agent*	46 (12)	33 (8)
Catheter Removal†	28 (7)	26 (7)
Catheter Removal without Thrombolytic Agent Use	17 (4)	15 (4)

- 
- \* 11 subjects in each arm had catheter removal following thrombolytic agent use and are included under both categories of use of a thrombolytic agent and catheter removal.
  - † Catheter removal refers to catheters removed due to loss of catheter patency. Overall, a total of 236 patients in the DEFENCATH arm and 225 patients in the heparin arm had a catheter removal for any reason, including loss of catheter patency.

### Other Adverse Reactions

Clinically significant adverse reactions that occurred in fewer than 1% of patients receiving DEFENCATH in Trial 1 are listed below:

*Metabolism and Nutrition Disorders:* Hypocalcemia

*Nervous System Disorders:* Dysgeusia

## **6.2 Postmarketing Experience**

The following adverse reaction has been identified during the use of a taurolidine and heparin containing CLS outside of the United States. Because this reaction is reported from a population of uncertain size, it is not always possible to reliably estimate its frequency or establish a causal relationship to drug exposure.

*Nervous System Disorders:* Paresthesia

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

DEFENCATH is not intended for systemic administration. It is intended for use as a CLS in patients with kidney failure requiring chronic HD; therefore, maternal use is not expected to result in fetal exposure to the drug [see *Dosage and Administration (2)*]. No animal reproduction study was conducted with DEFENCATH.

### **8.2 Lactation**

#### Risk Summary

DEFENCATH is not intended for systemic administration. It is intended for use as a CLS in patients with kidney failure requiring chronic HD; therefore, breastfeeding is not expected to result in exposure of the infant to DEFENCATH [see *Dosage and Administration (2)*].

### **8.4 Pediatric Use**

The safety and effectiveness of DEFENCATH have not been established in pediatric patients. There are no available data on DEFENCATH use in pediatric patients.

### **8.5 Geriatric Use**

There were 327 patients 65 years of age and older in Trial 1 [see *Clinical Studies (14)*]. Of the total number of patients using DEFENCATH in this study, 162 (41%) were 65 years



as a sterile, non-pyrogenic, preservative-free, clear, aqueous-based solution of taurolidine and heparin with a pH of 5.5-6.5 in the following strengths:

- 3 mL of catheter lock solution in a single-dose vial containing taurolidine 40.5 mg/3 mL (13.5 mg/mL), and heparin 3,000 USP Units/3 mL (1,000 USP Units/mL)
- 5 mL of catheter lock solution in a single-dose vial containing taurolidine 67.5 mg/5 mL (13.5 mg/mL), and heparin 5,000 USP Units/5 mL (1,000 USP Units/mL)

The heparin potency is determined by a biological assay using a USP reference standard based on units of heparin activity per milligram. The inactive ingredients are 26.1 mg/mL of citric acid, anhydrous (which is added to control pH and maintain the solubility of taurolidine), and may include hydrochloric acid and/or sodium hydroxide for pH adjustment.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

#### Heparin

Heparin interacts with the naturally occurring plasma protein, Antithrombin III, to induce a conformational change, which markedly enhances the serine protease activity of Antithrombin III, thereby inhibiting the activated coagulation factors involved in the clotting sequence, particularly Xa and IIa. Small amounts of heparin inhibit Factor Xa, and larger amounts inhibit thrombin (Factor IIa). Heparin also prevents the formation of a stable fibrin clot by inhibiting the activation of the fibrin stabilizing factor. Heparin does not have fibrinolytic activity; therefore, it will not lyse existing clots.

#### Taurolidine

Taurolidine is an antimicrobial drug [*see Microbiology (12.4)*].

### **12.2 Pharmacodynamics**

The pharmacodynamics of DEFENCATH are unknown.

### **12.3 Pharmacokinetics**

No pharmacokinetic studies were conducted because DEFENCATH is intended to be instilled into the catheter lumens at the conclusion of each HD session for patients with kidney failure requiring chronic HD. DEFENCATH is not intended for systemic administration [*see Dosage and Administration (2)*].

### **12.4 Microbiology**

#### Mechanism of Action

The mechanism of action of taurolidine and its metabolites is non-specific, causing damage to microbial cell walls and inhibiting adherence of microorganisms to biological surfaces.

#### Resistance

Time kill studies showed that taurolidine is bactericidal ( $>3 \log_{10}$  CFU reduction) at 1x to 4x minimum inhibitory concentration (MIC). The frequency of resistant mutants was  $<10^{-9}$  at 2x and 4x MIC for two strains each of *Escherichia coli*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Enterococcus faecium* and *Pseudomonas aeruginosa*.<sup>1</sup>

### Antimicrobial Activity

The following *in vitro* data are available but their clinical significance is unknown. Taurolidine has been shown to be active against most isolates of the following microorganisms:

#### Gram-positive

*Staphylococcus aureus* (including methicillin-sensitive *Staphylococcus aureus* (MSSA) and methicillin-resistant *Staphylococcus aureus* (MRSA))

*Staphylococcus epidermidis*

*Enterococcus faecalis*

#### Gram-negative

*Escherichia coli*

*Klebsiella pneumoniae*

*Pseudomonas aeruginosa*

*Serratia marcescens*

#### Fungi

*Candida albicans*

*Candida glabrata*

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

Carcinogenicity studies in animals have not been conducted with taurolidine or heparin.

#### Mutagenesis

Taurolidine was positive *in vitro* in an Ames reverse mutation assay and it increased mutation frequency in L5178Y mouse lymphoma cells. However, taurolidine was negative in an *in vivo* combined micronucleus test and comet assay in rats at the highest dose tested, 600 mg/kg/day administered intravenously.

No studies have been conducted with heparin to determine its mutagenic potential.

#### Impairment of Fertility

Fertility studies have not been conducted with DEFENCATH due to its negligible clinical exposure.

## **14 CLINICAL STUDIES**

The efficacy and safety of DEFENCATH for reducing the incidence of CRBSI in patients with kidney failure receiving chronic HD was evaluated in LOCK-IT-100 (referred to as Trial 1, NCT02651428), a randomized, double-blind, active-controlled, multicenter trial.

A total of 806 patients were randomized in a 1:1 ratio to receive either DEFENCATH or heparin (heparin sodium USP 1,000 units/mL, benzyl alcohol 9.45 mg/mL and sodium chloride 9.0 mg/mL) as a CLS. Enrollment in Trial 1 was not limited to patients with specific types of HD catheters. DEFENCATH or heparin was instilled into central venous HD catheters at the end of all dialysis sessions and was withdrawn prior to the initiation of the next dialysis session. The median age of the study population was 63 years

(range 21-94 years); 58% identified as males and 63% identified as white. The majority of patients (98%) had HD treatment three times per week, and 48% had their catheter implanted within three months prior to randomization.

A clinical adjudication committee (CAC) assessed the cases of CRBSI. The CAC definition for CRBSI included one positive blood culture (other than for coagulase-negative staphylococci, which required a confirmatory culture) from a peripheral site or either the arterial or venous catheter hub or the arterial or venous dialysis blood line and the patient had to have signs and symptoms of infection and no other apparent source of bloodstream infection.

Results of analyses of CAC-adjudicated CRBSI among the DEFENCATH and heparin groups in all randomized patients who received at least one dose of allocated study CLS are presented in Table 3. Patients in the DEFENCATH group had a lower incidence of CRBSI events compared to patients in the control group.

**Table 3. Results of Analyses of CAC-Adjudicated CRBSI in Patients with Kidney Failure Receiving HD in Trial 1**

	<b>DEFENCATH (N=397)</b>	<b>Heparin (N=398)</b>
CAC-adjudicated CRBSI	9 (2.3%)	32 (8.0%)
Event rate per 1000 catheter-days (95% CI)	0.13 (0.07, 0.26)	0.46 (0.33, 0.66)
Risk reduction (95% CI)*	71% (38%, 86%)	
Log-rank test p-value	0.0006	

\* Based on 1 - Hazard Ratio

In total, 5% [18/397] of subjects in the DEFENCATH arm and 5% [21/398] of subjects in the heparin arm died during the trial (difference [DEFENCATH - Heparin]: -0.7% with a 95% CI[-3.7%, 2.3%]).

Among the 9 patients in the DEFENCATH arm with a CAC-adjudicated CRBSI, 1 had a gram-negative organism isolated and 8 had a gram-positive organism isolated; among the 32 patients in the heparin arm with a CAC-adjudicated CRBSI, 14 had a gram-negative organism isolated and 18 had a gram-positive organism isolated.

## **15 REFERENCES**

1. Torres-Viera, C, Thauvin-Eliopoulos C., Souli M, DeGirolamin P, Farris MG Wennerstein CB, Sofia RD and Eliopoulos GM. 2000. Antimicrobial Agents and Chemotherapy 44(6): 1720 - 1724.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

DEFENCATH catheter lock solution is available in:

- 3 mL of catheter lock solution in a single-dose vial containing taurolidine 40.5 mg/3 mL (13.5 mg/mL), and heparin 3,000 USP Units/3 mL (1,000 USP Units/mL) (NDC 72990-103-03)
- 5 mL of catheter lock solution in a single-dose vial containing taurolidine 67.5 mg/5

mL (13.5 mg/mL), and heparin 5,000 USP Units/5 mL (1,000 USP Units/mL) (NDC 72990-105-05)

Each vial contains a sterile, preservative-free, clear aqueous-based solution for instillation in central venous catheters. Each carton contains 10 single-dose vials.

DEFENCATH vials must be stored at a controlled room temperature of 20°C to 25°C (68°F to 77°F); excursions are permitted from 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Do not freeze. DEFENCATH vials must be stored in the commercial carton, prior to the instillation in central venous catheters.

Manufactured for:  
CorMedix Inc.  
300 Connell Drive, Suite 4200  
Berkeley Heights, NJ 07922

## **PRINCIPAL DISPLAY PANEL**

NDC 72990-**103**-10

Rx Only

**DEFENCATH®**

Taurolidine and Heparin Catheter Lock Solution

**Limited  
Population\***

**Taurolidine 40.5 mg/3 mL (13.5 mg/mL) and  
Heparin 3,000 USP Units/3 mL (1,000 USP Units/mL)**

Derived from Porcine Intestinal Mucosa

**3 mL**


**For Instillation in Central Venous Catheters**

**Not** for systemic administration. Do **not** use as a catheter lock flush

\*See prescribing information for DEFENCATH for information about the limited population.

**Contains 10 Single-dose vials.  
Discard unused portion.**

**CorMedix Inc.®**



## PRINCIPAL DISPLAY PANEL

NDC 72990-103-10

Rx Only

**DEFENCATH<sup>®</sup>**

Taurolidine and Heparin Catheter Lock Solution

**Limited  
Population\***

# Taurolidine 40.5 mg/3 mL (13.5 mg/mL) and Heparin 3,000 USP Units/3 mL (1,000 USP Units/mL)

Derived from Porcine Intestinal Mucosa

**3 mL**

**For Instillation in Central Venous Catheters**

**Not** for systemic administration.

Do **not** use as a catheter lock flush

\*See prescribing information for DEFENCATH for information about the limited population.

**Contains 10 Single-dose vials.  
Discard unused portion.**

**CorMedix Inc.®**



## PRINCIPAL DISPLAY PANEL

NDC 72990-103-03

Rx Only

**DEFENCATH®**

Taurolidine and Heparin Catheter Lock Solution

**Limited  
Population\***

**Taurolidine 40.5 mg/3 mL (13.5 mg/mL)  
and Heparin 3,000 USP Units/3 mL  
(1,000 USP Units/mL)**

Derived from Porcine Intestinal Mucosa

**For Instillation in Central Venous  
Catheters**

**Not** for systemic administration  
Do **not** use as a catheter lock flush

Store at 20°C to 25°C  
(68°F to 77°F)

**3 mL Fill Single-Dose Vial  
Discard Unused Portion**

NDC 72990-103-03 Rx Only

**DEFENCATH®**  
Taurolidine and Heparin Catheter Lock Solution | Limited Population\*

**Taurolidine 40.5 mg/3 mL (13.5 mg/mL)  
and Heparin 3,000 USP Units/3 mL  
(1,000 USP Units/mL)**

Derived from Porcine Intestinal Mucosa

**For Instillation in Central Venous  
Catheters**  
**Not** for systemic administration  
Do **not** use as a catheter lock flush

Store at 20°C to 25°C 3 mL Fill Single-Dose Vial  
(68°F to 77°F) Discard Unused Portion

00372990103031

Mfd for: CorMedix Inc.  
300 Connell Drive,  
Suite 4200,  
Berkeley Heights,  
NJ 07922

48423/22/25

## PRINCIPAL DISPLAY PANEL

NDC 72990-103-03

Rx Only

**DEFENCATH®**

Taurolidine and Heparin Catheter Lock Solution

**Limited  
Population\***

**Taurolidine 40.5 mg/3 mL (13.5 mg/mL)  
and Heparin 3,000 USP Units/3 mL  
(1,000 USP Units/mL)**

Derived from Porcine Intestinal Mucosa

**For Instillation in Central Venous  
Catheters**

**Not** for systemic administration  
Do **not** use as a catheter lock flush

Store at 20°C to 25°C  
(68°F to 77°F)

**3 mL Fill Single-Dose Vial  
Discard Unused Portion**

NDC 72990-103-03 Rx Only  
**DEFENCATH**<sup>®</sup> Limited Population\*  
Taurolidine and Heparin Catheter Lock Solution

**Taurolidine 40.5 mg/3 mL (13.5 mg/mL)  
and Heparin 3,000 USP Units/3 mL  
(1,000 USP Units/mL)**  
Derived from Porcine Intestinal Mucosa

**For Instillation in Central Venous  
Catheters**  
**Not** for systemic administration  
Do **not** use as a catheter lock flush

Store at 20°C to 25°C 3 mL Fill Single-Dose Vial  
(68°F to 77°F) Discard Unused Portion

Mfd for: CorMedix Inc.  
300 Connell Drive, Suite 4200,  
Berkeley Heights, NJ 07922

00372990103031

620406

**PRINCIPAL DISPLAY PANEL**

NDC 72990-105-10

Rx Only

**DEFENCATH**<sup>™</sup>

Taurolidine and Heparin Catheter Lock Solution

Limited

Population\*

# Taurolidine 67.5 mg/5mL (13.5 mg/mL) and Heparin 5,000 USP Units/5 mL (1,000 USP Units/mL)

Derived from Porcine Intestinal Mucosa

5 mL

## For Instillation in Central Venous Catheters

Not for systemic administration

Do **not** use as a catheter lock flush

\*See prescribing information for Defencath for information about the limited population.

**Contains 10 Single-dose vials.**  
**Discard unused portion.**

**CorMedix Inc.™**



**PRINCIPAL DISPLAY PANEL**

NDC 72990-105-05

Rx Only

**DEFENCATH™**

Taurolidine and Heparin Catheter Lock Solution

Limited  
Population\*

**Taurolidine 67.5 mg/5mL (13.5 mg/mL)  
and Heparin 5,000 USP Units/5 mL  
(1,000 USP Units/mL)**

Derived from Porcine Intestinal Mucosa

**For Instillation in Central Venous  
Catheters**

**Not** for systemic administration  
Do **not** use as a catheter lock flush

Store at 20°C to 25°C  
(68°F to 77°F)

**5 mL Fill Single-Dose Vial -  
Discard Unused Portion**

NDC 72990-105-05

 **DEFENCATH™**  
Taurolidine and Heparin Catheter Lock Solution

Rx Only

Limited  
Population\*

**Taurolidine 67.5 mg/5mL (13.5 mg/mL)  
and Heparin 5,000 USP Units/5 mL  
(1,000 USP Units/mL)**

Derived from Porcine Intestinal Mucosa

**For Instillation in Central Venous  
Catheters**

**Not** for systemic administration  
Do **not** use as a catheter lock flush

Store at 20°C to 25°C  
(68°F to 77°F)

**5 mL Fill Single-Dose Vial -  
Discard Unused Portion**

Linear Barcode

Mfd for: CorMedix Inc.  
300 Connell Drive, Suite 4200,  
Berkeley Heights, NJ 07922

Lot XXXXXX

Exp YYYY-MM-DD



620281

# DEFENCATH

taurolidine and heparin solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:72990-103
<b>Route of Administration</b>	INTRALUMINAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>TAUROLIDINE</b> (UNII: 8OBZ1M4V3V) (TAUROLIDINE - UNII:8OBZ1M4V3V)	TAUROLIDINE	40.5 mg in 3 mL
<b>HEPARIN</b> (UNII: T2410KM04A) (HEPARIN - UNII:T2410KM04A)	HEPARIN	3000 [USP'U] in 3 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	78.3 mg in 3 mL
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72990-103-10	10 in 1 CARTON	04/09/2024	
1	NDC:72990-103-03	3 mL in 1 VIAL, SINGLE-DOSE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA214520	04/09/2024	

# DEFENCATH

taurolidine and heparin solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:72990-105
<b>Route of Administration</b>	INTRALUMINAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>TAUROLIDINE</b> (UNII: 8OBZ1M4V3V) (TAUROLIDINE - UNII:8OBZ1M4V3V)	TAUROLIDINE	67.5 mg in 5 mL
<b>HEPARIN</b> (UNII: T2410KM04A) (HEPARIN - UNII:T2410KM04A)	HEPARIN	5000 [USP'U] in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	130.5 mg in 5 mL
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72990-105-10	10 in 1 CARTON	04/09/2024	04/09/2024
1	NDC:72990-105-05	5 mL in 1 VIAL, SINGLE-DOSE; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA214520	04/09/2024	04/09/2024

**Labeler** - CorMedix Inc. (849818831)

Revised: 9/2025

CorMedix Inc.