

ISOFLURANE- isoflurane liquid

Piramal Critical Care, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ISOFLURANE USP safely and effectively. See full prescribing information for ISOFLURANE USP.

ISOFLURANE USP liquid for inhalation

Initial U.S. Approval: 1979

INDICATIONS AND USAGE

Isoflurane USP, a general anesthetic, is an inhalation agent indicated for induction and maintenance of general anesthesia. (1)

DOSAGE AND ADMINISTRATION

- Isoflurane USP should be administered only by persons trained in the administration of general anesthesia. Isoflurane USP should only be delivered using a vaporizer specifically designed and designated for use with isoflurane. (2)
- The administration of general anesthesia must be individualized and titrated based on the patient's age and clinical status. (2)

DOSAGE FORMS AND STRENGTHS

Liquid (stable): 100% (3)

CONTRAINDICATIONS

- Patients in whom general anesthesia is contraindicated (4)
- Patients with known sensitivity to Isoflurane USP or other halogenated agents (4)
- Patients with known or suspected genetic susceptibility to malignant hyperthermia (4)
- Patients with a history of confirmed hepatitis due to a halogenated inhalational anesthetic or a history of unexplained moderate to severe hepatic dysfunction (e.g., jaundice associated with fever and/or eosinophilia) after anesthesia with Isoflurane USP or other halogenated inhalational anesthetics (4)

WARNINGS AND PRECAUTIONS

- Malignant Hyperthermia: Malignant hyperthermia may occur, especially in individuals with known or suspected susceptibility based on genetic factors or family history. Discontinue triggering agents, administer intravenous dantrolene sodium, and apply supportive therapies. (5.1)
- Perioperative Hyperkalemia: Perioperative hyperkalemia may occur. Patients with latent or overt neuromuscular disease, particularly with Duchenne muscular dystrophy, appear to be most vulnerable. Early, aggressive intervention is recommended. (5.2)
- Hepatic Reactions: May cause sensitivity hepatitis in patients sensitized by previous exposure to halogenated anesthetics. Approach repeated anesthesia with caution. (5.3)
- Hypersensitivity Reactions: Allergic-type hypersensitivity reactions, including anaphylaxis, have been reported with isoflurane. (5.4)
- Abortions: Increased blood loss comparable to that seen with halothane has been observed in patients undergoing abortions. (5.5)
- QT Prolongation: Carefully monitor cardiac rhythm when administering Isoflurane USP to susceptible patients. (5.6)
- Interactions with Desiccated Carbon Dioxide (CO₂) Absorbents: May react with desiccated CO₂ absorbents to produce carbon monoxide. Replace desiccated CO₂ absorbent before administration of Isoflurane USP. (5.7)
- Pediatric Neurotoxicity: In developing animals, exposures greater than 3 hours cause neurotoxicity. Weigh benefits against potential risks when considering elective procedures in children under 3 years old. (5.8)

ADVERSE REACTIONS

Most common adverse reactions (incidence \geq 5%) are agitation, cough, breath holding, nausea, chills/shivering, vomiting, laryngospasm, delirium. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Piramal Critical Care, Inc. at 1-888-822-8431 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Concomitant use of N₂O and/or opioids reduces the MAC of Isoflurane USP. Adjust dose accordingly. (7.1, 7.2)
- Isoflurane USP decreases the doses of neuromuscular blocking agents required. Adjust dose accordingly. (7.3)

----- **USE IN SPECIFIC POPULATIONS** -----

Geriatric Use: The minimum alveolar concentration (MAC) of Isoflurane USP decreases with increasing patient age. (8.5)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 11/2025

Revised: 11/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information

2.2 Premedication

2.3 Induction

2.4 Maintenance

2.5 Use in Patients with Coronary Artery Disease

3. DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Malignant Hyperthermia

5.2 Perioperative Hyperkalemia

5.3 Hepatic Reactions

5.4 Hypersensitivity Reactions

5.5 Abortions

5.6 QT Prolongation

5.7 Interactions with Desiccated Carbon Dioxide Absorbents

5.8 Pediatric Neurotoxicity

5.9 Laboratory Tests

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Post-Marketing Experience

7 DRUG INTERACTIONS

7.1 Opioids

7.2 Nitrous Oxide

7.3 Neuromuscular Blocking Agents

7.4 Adrenaline

7.5 Calcium Antagonists

7.6 Concomitant use with Beta Blockers

7.7 Concomitant use with MAO Inhibitors

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.2 Pharmacodynamics

12.3 Pharmacokinetics
12.5 Pharmacogenomics
13. NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2 Animal Toxicology and/or Pharmacology
16 HOW SUPPLIED/STORAGE AND HANDLING
16.1 Safety and Handling
16.2 Storage

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Isoflurane USP may be used for induction and maintenance of general anesthesia. Adequate data have not been developed to establish its application in obstetrical anesthesia.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information

Isoflurane should be administered only by persons trained in the administration of general anesthesia. Facilities for maintenance of a patent airway, artificial ventilation, oxygen enrichment, and circulatory resuscitation must be immediately available.

Isoflurane is administered by inhalation. Isoflurane should be delivered from a vaporizer specifically designed for use with isoflurane. Dosage for induction and maintenance must be individualized and titrated to the desired effect according to the patient's age and clinical status. With the exception of neonates, the minimum alveolar concentration (MAC) of isoflurane decreases with increasing patient age.

Nitrous oxide decreases the MAC of isoflurane (see Table 1). Opioids decrease the MAC of isoflurane [see *Drug Interactions (7)*]. Isoflurane potentiates the muscle relaxant effect of all neuromuscular blockers and decreases the required doses of neuromuscular blocking agents [see *Drug Interactions (7)*]. The dose should be adjusted accordingly.

All patients anesthetized with isoflurane should be continually monitored (e.g., monitoring of the electrocardiogram, blood pressure, oxygen saturation, and end tidal CO₂). Isoflurane is a profound respiratory depressant. Excessive respiratory depression may be related to depth of anesthesia and respond to decreasing the inspired concentration of isoflurane. The depressant effect is accentuated by concurrent use of opioids and other respiratory depressants. Respiration should be closely monitored and assisted or controlled ventilation employed when necessary.

2.2 Premedication

Premedication should be selected according to the need of the individual patient, taking into account that secretions are weakly stimulated by Isoflurane USP, and the heart rate tends to be increased.

2.3 Induction

Induction with isoflurane in oxygen or in combination with oxygen-nitrous oxide mixtures may produce coughing, breath holding, laryngospasm and bronchospasm, which increases with the concentration of isoflurane. These difficulties may be avoided by the use of a hypnotic dose of an ultra-short-acting barbiturate. Inspired concentrations of 1.5 to 3% isoflurane usually produce surgical anesthesia in 7 to 10 minutes.

2.4 Maintenance

Isoflurane MAC values according to age are shown below:

Table 1: Effect of Age on Minimum Alveolar Concentration of Isoflurane

Age	Average MAC Value In 100% Oxygen	Average MAC Value In 30% Oxygen and 70% N₂O
Preterm neonates less than 32 weeks gestational age	1.28%	
Preterm neonates 32-37 weeks gestational age	1.41%	
0-1 month	1.60%	
1-6 months	1.87%	
6-12 months	1.80%	
1-5 years	1.60%	
6-10 years	1.45%	
11-18 years	1.38%	
19-30 years	1.28%	0.56%
31-55 years	1.15%	0.50%
55-83 years	1.05%	0.37%

Dosage for induction and maintenance must be individualized and titrated to the desired effect according to the patient's age and clinical status.

Surgical levels of anesthesia may be sustained with a 1 to 2.5% concentration when nitrous oxide is used concomitantly. An additional 0.5 to 1% may be required when isoflurane is given using oxygen alone. If added relaxation is required, supplemental doses of neuromuscular blocking agents may be used.

The level of blood pressure during maintenance is an inverse function of isoflurane concentration in the absence of other complicating problems. Excessive decreases may be due to depth of anesthesia and in such instances may be corrected by lightening anesthesia.

Isoflurane causes a dose-dependent reduction in systemic vascular resistance and

blood pressure. Particular care must be taken when selecting the dosage for patients who are hypovolemic, hypotensive, or otherwise hemodynamically compromised, e.g., due to concomitant medications.

Isoflurane USP markedly increases cerebral blood flow at deeper levels of anesthesia to produce a transient increase in intracranial pressure. In patients with or at risk for elevations of intracranial pressure (ICP), administer isoflurane in conjunction with ICP-reducing strategies, as clinically appropriate.

2.5 Use in Patients with Coronary Artery Disease

Regardless of the anesthetics employed, maintenance of normal hemodynamics is important to the avoidance of myocardial ischemia in patients with coronary artery disease.

Isoflurane can cause dose-dependent coronary vasodilation and has been shown to divert blood from collateral-dependent myocardium to normally perfused areas in an animal model (“coronary steal”). The extent to which coronary steal occurs in patients with steal-prone coronary anatomy is unclear. Monitor for signs of inadequate myocardial perfusion via hemodynamic monitors (e.g., ECG, blood pressure) during isoflurane administration. Consider additional cardiac monitoring in patients with known coronary artery disease, as clinically necessary.

3. DOSAGE FORMS AND STRENGTHS

Isoflurane is a clear, colorless, stable liquid containing no additives or chemical stabilizers, 100% isoflurane.

4 CONTRAINDICATIONS

Isoflurane USP is contraindicated in patients:

- in whom general anesthesia is contraindicated.
- with known sensitivity to Isoflurane USP or to other halogenated agents [*see Warnings and Precautions (5.3)*].
- with known or suspected genetic susceptibility to malignant hyperthermia [*see Warnings and Precautions (5.1), Clinical Pharmacology (12.5)*].
- with a history of confirmed hepatitis due to a halogenated inhalational anesthetic or a history of unexplained moderate to severe hepatic dysfunction (e.g., jaundice associated with fever and/or eosinophilia) after anesthesia with isoflurane or other halogenated inhalational anesthetics.

5 WARNINGS AND PRECAUTIONS

5.1 Malignant Hyperthermia

In susceptible individuals, volatile anesthetic agents, including Isoflurane USP, may trigger malignant hyperthermia, a skeletal muscle hypermetabolic state leading to high oxygen demand. Fatal outcomes of malignant hyperthermia have been reported.

The risk of developing malignant hyperthermia increases with the concomitant administration of succinylcholine and volatile anesthetic agents. Isoflurane USP can induce malignant hyperthermia in patients with known or suspected susceptibility based on genetic factors or family history, including those with certain inherited ryanodine receptor (*RYR1*) or dihydropyridine receptor (*CACNA1S*) variants [see *Contraindications (4)*, *Clinical Pharmacology (12.5)*].

Signs consistent with malignant hyperthermia may include hyperthermia, hypoxia, hypercapnia, muscle rigidity (e.g., jaw muscle spasm), tachycardia (e.g., particularly that unresponsive to deepening anesthesia or analgesic medication administration), tachypnea, cyanosis, arrhythmias, hypovolemia, and hemodynamic instability. Skin mottling, coagulopathies, and renal failure may occur later in the course of the hypermetabolic process.

Successful treatment of malignant hyperthermia depends on early recognition of the clinical signs. If malignant hyperthermia is suspected, discontinue all triggering agents (i.e., volatile anesthetic agents and succinylcholine), administer intravenous dantrolene sodium, and initiate supportive therapies. Consult prescribing information for intravenous dantrolene sodium for additional information on patient management. Supportive therapies include administration of supplemental oxygen and respiratory support based on clinical need, maintenance of hemodynamic stability and adequate urinary output, management of fluid and electrolyte balance, correction of acid base derangements, and institution of measures to control rising temperature.

5.2 Perioperative Hyperkalemia

Use of inhaled anesthetic agents has been associated with rare increases in serum potassium levels that have resulted in cardiac arrhythmias and death in pediatric patients during the postoperative period. Patients with latent as well as overt neuromuscular disease, particularly Duchenne muscular dystrophy, appear to be most vulnerable. Concomitant use of succinylcholine has been associated with most, but not all, of these cases. These patients also experienced significant elevations in serum creatinine kinase levels and, in some cases, changes in urine consistent with myoglobinuria. Despite the similarity in presentation to malignant hyperthermia, none of these patients exhibited signs or symptoms of muscle rigidity or hypermetabolic state. Early and aggressive intervention to treat the hyperkalemia and resistant arrhythmias is recommended, as is subsequent evaluation for latent neuromuscular disease.

5.3 Hepatic Reactions

Cases of mild, moderate and severe postoperative hepatic dysfunction or hepatitis with or without jaundice, including fatal hepatic necrosis and hepatic failure, have been reported with isoflurane.

Such reactions can represent hypersensitivity hepatitis, a known risk of exposure to halogenated anesthetics, including isoflurane. As with other halogenated anesthetic agents, Isoflurane USP may cause sensitivity hepatitis in patients who have been sensitized by previous exposure to halogenated anesthetics [see *Contraindications (4)*].

Clinical judgment should be exercised when isoflurane is used in patients with underlying hepatic conditions or under treatment with drugs known to cause hepatic dysfunction. [see *Contraindications (4)*].

As with all halogenated anesthetics, repeated anesthetics within a short period of time may result in increased effects, particularly in patients with underlying hepatic conditions, or additive effects in patients treated with drugs known to cause hepatic dysfunction. Evaluate the need for repeated exposure in each individual patient and adjust the dose of isoflurane based on signs and symptoms of adequate depth of anesthesia if repeated exposure in a short period of time is clinically indicated.

5.4 Hypersensitivity Reactions

Allergic-type hypersensitivity reactions, including anaphylaxis, have been reported with isoflurane. Manifestations of such reactions have included hypotension, rash, difficulty breathing and cardiovascular collapse.

5.5 Abortions

Increased blood loss comparable to that seen with halothane has been observed in patients undergoing abortions.

5.6 QT Prolongation

QT prolongation, with rare instances of torsade de pointes, have been reported. Monitor QT interval when administering isoflurane to susceptible patients (e.g., patients with congenital Long QT Syndrome or patients taking drugs that can prolong the QT interval).

5.7 Interactions with Desiccated Carbon Dioxide Absorbents

Isoflurane USP, like some other inhalational anesthetics, can react with desiccated carbon dioxide (CO₂) absorbents to produce carbon monoxide, which may result in elevated levels of carboxyhemoglobin in some patients. Barium hydroxide lime and soda lime become desiccated when fresh gases are passed through the CO₂ absorber canister at highflow rates over many hours or days. When a clinician suspects that CO₂ absorbent may be desiccated, it should be replaced before the administration of Isoflurane USP.

The color indicator of most CO₂ absorbents does not necessarily change as a result of desiccation. Therefore, the lack of significant color change should not be taken as assurance of adequate hydration of the CO₂ absorbent material. CO₂ absorbents should be replaced routinely regardless of the state of color indicator following current manufacturer's guidelines for use of anesthesiology equipment.

5.8 Pediatric Neurotoxicity

Published animal studies demonstrate that the administration of anesthetic and sedation drugs that block NMDA receptors and/or potentiate GABA activity increase neuronal apoptosis in the developing brain and result in long-term cognitive deficits when used for longer than 3 hours. The clinical significance of these findings is not clear. However, based on the available data, the window of vulnerability to these changes is believed to correlate with exposures in the third trimester of gestation through the first several months of life, but may extend out to approximately three years of age in humans [see

Use in Specific Populations (8.1, 8.4), Nonclinical Toxicology (13.2)].

Some published studies in children suggest that similar deficits may occur after repeated or prolonged exposures to anesthetic agents early in life and may result in adverse cognitive or behavioral effects. These studies have substantial limitations, and it is not clear if the observed effects are due to the anesthetic/sedation drug administration or other factors such as the surgery or underlying illness.

Anesthetic and sedation drugs are a necessary part of the care of children needing surgery, other procedures, or tests that cannot be delayed, and no specific medications have been shown to be safer than any other. Decisions regarding the timing of any elective procedures requiring anesthesia should take into consideration the benefits of the procedure weighed against the potential risks.

5.9 Laboratory Tests

Transient increases in BSP retention, blood glucose and serum creatinine with decrease in BUN, serum cholesterol and alkaline phosphatase have been observed.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

The following adverse reactions were identified from controlled clinical trials of adult and pediatric subjects exposed to Isoflurane USP. The trials were conducted using a variety of pre-medications, other anesthetics, and surgical procedures of varying lengths.

The most serious reported adverse reactions in alphabetical order are agitation, arrhythmia, breath holding, elevated liver enzyme, hypotension and laryngospasm.

The most frequent adverse reactions (incidence ³ 5%) described in Table 1 are agitation, breath holding, chills/shivering, cough, delirium, laryngospasm, nausea, and vomiting.

Adverse reactions with an incidence between 1% and 5% are provided in Table 2.

Adverse reactions with an incidence less than 1% are provided in Table 3.

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Table 2: Adverse Reactions \geq 5%

System Organ Class (SOC)	Adverse Reaction	Frequency
PSYCHIATRIC DISORDERS	Delirium	6.2% (N=2830)
NERVOUS SYSTEM DISORDERS	Agitation (Excitement)	Induction 51.8% (N=515) ¹
	Breath holding	Induction 23.9% (N=515) ¹

RESPIRATORY, THORACIC, AND MEDIASTINAL DISORDERS	Cough	Induction	28.2% (N=515) ¹
	Laryngospasm	Induction	8.0% (N=515) ¹
GASTROINTESTINAL DISORDERS	Nausea	Recovery	15.4 % (N=2830)
	Vomiting	Recovery	9.5% (N=2830)
GENERAL DISORDERS AND ADMINISTRATIVE SITE CONDITIONS	Chills/shivering		14.0% (N=1691) ²

¹Represents patients not receiving intravenous agents or neuromuscular blocking agents for intubation (i.e., patients receiving inhalation induction).

²Reflects the number of patients with recorded body temperature measurements.

Table 3: Adverse Reactions between 1% and 5%

System Organ Class (SOC)	Adverse Reaction		Frequency
NERVOUS SYSTEM DISORDERS	Movement	Maintenance	1.8% (N=2830)
CARDIAC DISORDERS	Ventricular arrhythmia (Intraoperative)	Induction	2.1% (N=2161)
		Maintenance	2.7% (N=2253)
	Nodal arrhythmia (Intraoperative)	Induction	4.0% (N=2161)
		Maintenance	1.7% (N=2253)
	Atrial arrhythmia (Intraoperative)	Induction	1.6% (N=2161)
		Maintenance	2.2% (N=2253)
Arrhythmia (Postoperative)		1.1% (N=2830)	
RESPIRATORY, THORACIC, AND MEDIASTINAL DISORDERS	Breath holding	Maintenance	1.1% (N=359) ¹
	Cough	Maintenance	4.2 % (N=359) ¹

¹Represents patients not receiving intravenous agents or neuromuscular blocking agents for intubation (i.e., patients receiving inhalation induction).

Table 4: Adverse Reactions less than 1%

System Organ Class (SOC)	Adverse Reaction	Frequency
---------------------------------	-------------------------	------------------

PSYCHIATRIC DISORDERS	Mood changes		0.3% (N=2830)
	Nightmare		0.4% (N=2175) ¹
NERVOUS SYSTEM DISORDERS	Convulsive pattern on electroencephalogram		0.5% (N=200) ²
	Seizure		0.04% (N=2830)
VASCULAR DISORDERS	Hypotension	Postoperative	0.3% (N=2830)
	Hypertension	Postoperative	0.1% (N=2830)
RESPIRATORY, THORACIC, AND MEDIASTINAL DISORDERS	Laryngospasm	Maintenance	0.8% (N=359) ³
	Secretions	Induction	0.2% (N=515) ³
		Maintenance	0.0% (N=359) ³
GASTROINTESTINAL DISORDERS	Vomiting	Induction	0.8% (N=515) ³
	Retching	Induction	1.0% (N=515) ³
		Maintenance	0.8% (N=359) ³
SKIN AND SUBCUTANEOUS TISSUE DISORDERS	Diaphoresis	Induction	0.2% (N=515) ³
		Maintenance	0.0% (N=359) ³

¹Reflects the number of patients interviewed by a physician in the recovery period.

²Reflects the number of recorded electroencephalograms.

³Represents patients not receiving intravenous agents or neuromuscular blocking agents for intubation (i.e., patients receiving inhalation induction).

The following adverse reactions (see Table 5) were observed, but due to limited data, frequency could not be determined.

Table 5: Adverse Reactions with Unknown Frequency

BLOOD AND LYMPHATIC SYSTEM DISORDERS:	White blood cell count increased
METABOLISM AND NUTRITION DISORDERS:	Blood glucose increased
PSYCHIATRIC	

PSYCHIATRIC DISORDERS:	Confused state, Nervousness
NERVOUS SYSTEM DISORDERS:	Ataxia; Dizziness; Drowsiness; Intellectual function decrease
VASCULAR DISORDERS:	Hypotension (Intraoperative); Hypertension (Intraoperative)
HEPATOBIILIARY DISORDERS:	Blood bilirubin increased; Bromsulphthalein clearance decreased; Alanine aminotransferase increased; Aspartate aminotransferase increased; Blood alkaline phosphatase increased; Blood lactate dehydrogenase increased
MUSCULOSKELETAL, CONNECTIVE TISSUE AND BONE DISORDERS:	Myalgia
GENERAL DISORDERS AND ADMINISTRATIVE SITE CONDITIONS:	Asthenia; Fatigue

6.2 Post-Marketing Experience

The following adverse reactions have been identified during post-approval use of Isoflurane USP. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Post-marketing adverse reactions are listed by MedDRA System Organ Class (SOC), then by preferred term in order of decreasing severity.

BLOOD AND LYMPHATIC SYSTEM DISORDERS:Carboxyhemoglobin increased

IMMUNE SYSTEM DISORDERS:Anaphylactic reaction

METABOLISM AND NUTRITION DISORDERS:Hyperkalemia in patients with underlying myopathies

PSYCHIATRIC DISORDERS:Withdrawal syndrome (following multi-day exposure; symptoms include seizure, hallucination, ataxia, agitation, confusion)

NERVOUS SYSTEM DISORDERS:Brain edema, Intracranial pressure increased, Migraine, Myoclonus, Nystagmus, Pupils unequal, Headache

CARDIAC DISORDERS:Cardiac arrest, Ventricular fibrillation, Torsade de pointes, Myocardial infarction, Myocardial ischemia, Atrioventricular block complete, Atrioventricular block second degree, Atrial fibrillation, Electrocardiogram QT prolonged, Atrioventricular block first degree, Ventricular tachycardia, Ventricular extrasystoles, Tachycardia, Bradycardia, Cardiac output decreased

VASCULAR DISORDERS:Flushing

RESPIRATORY, THORACIC, AND MEDIASTINAL DISORDERS:Apnea, Hypoxia, Bronchospasm, Airway obstruction, Respiratory depression, Hypercapnia, Stridor, Hiccough

GASTROINTESTINAL DISORDERS:Pancreatitis

HEPATOBIILIARY DISORDERS:Hepatic necrosis, Hepatic failure, Hepatitis fulminant, Cholestatic hepatitis, Hepatitis, Hepatic steatosis, Jaundice, Gamma- glutamyltransferase increased

SKIN AND SUBCUTANEOUS TISSUE DISORDERS:Rash

MUSCULOSKELETAL, CONNECTIVE TISSUE AND BONE DISORDERS:Rhabdomyolysis

RENAL AND URINARY DISORDERS:Acute renal failure**, Oliguria**

GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS:Malignant hyperthermia, hypothermia

INJURY, POISONING, AND PROCEDURAL COMPLICATIONS*:Unwanted awareness during anesthesia; Dyspnea, Bronchospasm, Stridor, Cough, Dizziness, Paresthesia, Hepatic reactions, Flushing, Rash, Contact dermatitis, Erythema, Periorbital edema, Eye irritation, Conjunctival hyperemia, Headache

*All reactions categorized within this SOC, with the exception of, Unwanted awareness during anesthesia, were from occupational exposure in non-patients.

**Cases of acute renal failure and oliguria have been reported after isoflurane anesthesia. These events may be secondary to hypotension or other effects of isoflurane.

7 DRUG INTERACTIONS

7.1 Opioids

Opioids decrease the Minimum Alveolar Concentration (MAC) of isoflurane. Opioids such as fentanyl and its analogues, when combined with isoflurane, may lead to a synergistic fall in blood pressure and respiratory rate.

7.2 Nitrous Oxide

Nitrous oxide decreases the MAC of isoflurane [*see Dosage and Administration (2.1)*].

7.3 Neuromuscular Blocking Agents

Isoflurane potentiates the muscle relaxant effect of all neuromuscular blocking agents and decreases the required doses of neuromuscular blocking agents. In general, anesthetic concentrations isoflurane at equilibrium reduce the ED₉₅ of succinylcholine, atracurium, pancuronium, rocuronium and vecuronium by approximately 25 to 40% or

more compared to N₂O/opioid anesthesia. If added relaxation is required, supplemental doses of neuromuscular blocking agents may be used.

7.4 Adrenaline

Isoflurane is similar to sevoflurane in the sensitization of the myocardium to arrhythmogenic effect of exogenously administered adrenaline. Doses of adrenaline greater than 5mcg/kg, when administered submucosally may produce multiple ventricular arrhythmias.

7.5 Calcium Antagonists

Isoflurane may lead to marked hypotension in patients treated with calcium antagonists.

7.6 Concomitant use with Beta Blockers

Concomitant use of beta blockers may exaggerate the cardiovascular effects of inhalational anesthetics, including hypotension and negative inotropic effects.

7.7 Concomitant use with MAO Inhibitors

Concomitant use of MAO inhibitors and inhalational anesthetics may increase the risk of hemodynamic instability during surgery or medical procedures.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no adequate and well-controlled studies in pregnant women. In animal reproduction studies, embryofetal toxicity was noted in pregnant mice exposed to 0.075% (increased post implantation losses) and 0.3% isoflurane (increased post implantation losses and decreased livebirth index) during organogenesis.

Published studies in pregnant primates demonstrate that the administration of anesthetic and sedation drugs that block NMDA receptors and/or potentiate GABA activity during the period of peak brain development increases neuronal apoptosis in the developing brain of the offspring when used for longer than 3 hours. There are no data on pregnancy exposures in primates corresponding to periods prior to the third trimester in humans (see Data).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of

major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Data

Animal Data

Pregnant rats were exposed to isoflurane at concentrations of 0%, 0.1%, or 0.4% for two hours per day during organogenesis (Gestational Days 6-15). Isoflurane did not cause malformations or clear maternal toxicity under these conditions.

Pregnant mice exposed to isoflurane at concentrations of 0%, 0.075%, or 0.30% for 2 hours per day during organogenesis (Gestational Days 6-15). Isoflurane increased fetal toxicity (higher post implantation losses at 0.075 and 0.3% groups and significantly lower live-birth index in the 0.3% isoflurane treatment group). Isoflurane did not cause malformations or clear maternal toxicity under these conditions.

Pregnant rats were exposed to concentrations of isoflurane at 0%, 0.1%, or 0.4% for 2 hours per day during late gestation (GD 15-20). Animals appeared slightly sedated during exposure. No adverse effects on the offspring or evidence of maternal toxicity were reported. This study did not evaluate neurobehavioral function including learning and memory in the first generation (F1) of pups.

In a published study in primates, administration of an anesthetic dose of ketamine for 24 hours on Gestation Day 122 increased neuronal apoptosis in the developing brain of the fetus. In other published studies, administration of either isoflurane or propofol for 5 hours on Gestation Day 120 resulted in increased neuronal and oligodendrocyte apoptosis in the developing brain of the offspring. With respect to brain development, this time period corresponds to the third trimester of gestation in the human. The clinical significance of these findings is not clear; however, studies in juvenile animals suggest neuroapoptosis correlates with long-term cognitive deficits [see *Warnings and Precautions (5.8), Nonclinical Toxicology (13.2)*].

8.2 Lactation

Due to insufficient information regarding the excretion of isoflurane in human milk, the potential risks and benefits for each specific patient should be carefully considered before isoflurane is administered to nursing women.

8.4 Pediatric Use

During the induction of anesthesia, saliva flow and tracheobronchial secretion can increase and can be the cause of laryngospasm, particularly in children.

Published juvenile animal studies demonstrate that the administration of anesthetic and sedation drugs, such as Isoflurane USP, that either block NMDA receptors or potentiate the activity of GABA during the period of rapid brain growth or synaptogenesis, results in widespread neuronal and oligodendrocyte cell loss in the developing brain and alterations in synaptic morphology and neurogenesis. Based on comparisons across species, the window of vulnerability to these changes is believed to correlate with exposures in the third trimester of gestation through the first several months of life, but may extend out to approximately 3 years of age in humans.

In primates, exposure to 3 hours of ketamine that produced a light surgical plane of anesthesia did not increase neuronal cell loss, however, treatment regimens of 5 hours or longer of isoflurane increased neuronal cell loss. Data from isoflurane-treated rodents and ketamine-treated primates suggest that the neuronal and oligodendrocyte cell losses are associated with prolonged cognitive deficits in learning and memory. The

clinical significance of these nonclinical findings is not known, and healthcare providers should balance the benefits of appropriate anesthesia in pregnant women, neonates, and young children who require procedures with the potential risks suggested by the nonclinical data [see *Warnings and Precautions (5.8)*, *Nonclinical Toxicology (13.2)*].

8.5 Geriatric Use

The minimum alveolar concentration (MAC) of isoflurane decreases with increasing patient age. The dose should be adjusted accordingly [see *Dosage and Administration (2.4)*].

10 OVERDOSAGE


The symptoms of overdosage of Isoflurane USP can present as a deepening of anesthesia, cardiac and/or respiratory depression in spontaneously breathing patients, and cardiac depression in ventilated patients in whom hypercapnia and hypoxia may occur only at a late stage.

In the event of overdosage, or what may appear to be overdosage, the following action should be taken, as appropriate:

Stop drug administration, establish a clear airway, and initiate assisted or controlled ventilation with pure oxygen. Monitor cardiovascular function and manage signs of poor end-organ perfusion as clinically indicated.

11 DESCRIPTION

Isoflurane USP, a nonflammable liquid administered by vaporizing, is a general inhalation anesthetic drug. It is 1-chloro-2,2,2-trifluoroethyl difluoromethyl ether, and its structural formula is:



Some physical constants are:

Molecular weight		184.5
Boiling point at 760 mm Hg		48.5°C
Refractive index $n_{20/D}$		1.2990- 1.3005
Specific gravity 25°/25°C		1.496
Vapor pressure in mm Hg**	20°C	238
	25°C	295
	30°C	367
	35°C	450

**Equation for vapor pressure calculation:
 $\log_{10} P_{\text{vap}} = A + B/T$ where:

$$A = 8.056$$

$$B = -1664.58$$

$$T = ^\circ\text{C} + 273.16 \text{ (Kelvin)}$$

Partition coefficients at 37°C

Water/gas	0.61
Blood/gas	1.43
Oil/gas	90.8

Partition coefficients at 25°C -rubber and plastic

Conductive rubber/gas	62.0
Butyl rubber/gas	75.0
Polyvinyl chloride/gas	110.0
Polyethylene/gas	~2.0
Polyurethane/gas	~1.4
Polyolefin/gas	~1.1
Butyl acetate/gas	~2.5
Purity by gas chromatography	>99.9%
Lower limit of flammability in oxygen or nitrous oxide at 9 joules/sec. and 23°C	None
Lower limit of flammability in oxygen or nitrous oxide at 900 concentration in joules/sec. and 23°C	Greater than useful concentration in anesthesia.

Isoflurane is a clear, colorless, stable liquid containing no additives or chemical stabilizers. Isoflurane has a mildly pungent, musty, ethereal odor. Samples stored in indirect sunlight in clear, colorless glass for five years, as well as samples directly exposed for 30 hours to a 2 amp, 115 volt, 60 cycle long wave U.V. light were unchanged in composition as determined by gas chromatography. Isoflurane in one normal sodium methoxide-methanol solution, a strong base, for over six months consumed essentially no alkali, indicative of strong base stability. Isoflurane does not decompose in the presence of soda lime (at normal operating temperatures), and does not attack aluminum, tin, brass, iron or copper.

12 CLINICAL PHARMACOLOGY

12.2 Pharmacodynamics

Induction of and recovery from isoflurane anesthesia are rapid. Isoflurane has a mild

pungency which limits the rate of induction, although excessive salivation or tracheobronchial secretions do not appear to be stimulated. Pharyngeal and laryngeal reflexes are readily obtunded. The level of anesthesia may be changed rapidly with isoflurane. Isoflurane is a profound respiratory depressant. As anesthetic dose is increased, tidal volume decreases and respiratory rate is unchanged. This depression is partially reversed by surgical stimulation, even at deeper levels of anesthesia. Isoflurane evokes a sigh response reminiscent of that seen with diethyl ether and enflurane, although the frequency is less than with enflurane.

Blood pressure decreases with induction of anesthesia but returns toward normal with surgical stimulation. Progressive increases in depth of anesthesia produce corresponding decreases in blood pressure. Nitrous oxide diminishes the inspiratory concentration of isoflurane required to reach a desired level of anesthesia and may reduce the arterial hypotension seen with isoflurane alone. Heart rhythm is remarkably stable. With controlled ventilation and normal PaCO₂, cardiac output is maintained despite increasing depth of anesthesia, primarily through an increase in heart rate which compensates for a reduction in stroke volume. The hypercapnia which attends spontaneous ventilation during isoflurane anesthesia further increases heart rate and raises cardiac output above awake levels.

Muscle relaxation is often adequate for intra-abdominal operations at normal levels of anesthesia. Complete muscle paralysis can be attained with small doses of neuromuscular blocking agents. ALL COMMONLY USED NEUROMUSCULAR BLOCKING AGENTS ARE MARKEDLY POTENTIATED WITH ISOFLURANE, THE EFFECT BEING MOST PROFOUND WITH THE NONDEPOLARIZING TYPE. Neostigmine reverses the effect of nondepolarizing neuromuscular blocking agents in the presence of isoflurane. All commonly used neuromuscular blocking agents are compatible with isoflurane.

Isoflurane can produce coronary vasodilation at the arteriolar level in selected animal models; the drug is probably also a coronary dilator in humans. Isoflurane, like some other coronary arteriolar dilators, has been shown to divert blood from collateral dependent myocardium to normally perfused areas in an animal model ("coronary steal"). Clinical trials to date evaluating myocardial ischemia, infarction and death as outcome parameters have not established that the coronary arteriolar dilation property of isoflurane is associated with coronary steal or myocardial ischemia in patients with coronary artery disease.

12.3 Pharmacokinetics

Isoflurane undergoes minimal biotransformation in man. In the postanesthesia period, only 0.17% of the isoflurane taken up can be recovered as urinary metabolites.

12.5 Pharmacogenomics

RYR1 and *CACNA1S* are polymorphic genes, and multiple pathogenic variants have been associated with malignant hyperthermia susceptibility (MHS) in patients receiving volatile anesthetic agents, including Isoflurane USP. Case reports as well as ex-vivo studies have identified multiple variants in *RYR1* and *CACNA1S* associated with MHS. Variant pathogenicity should be assessed based on prior clinical experience, functional studies, prevalence information, or other evidence [see *Contraindications (4)*, *Warnings and Precautions (5.1)*].

13. NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Swiss ICR mice were given isoflurane to determine whether such exposure might induce neoplasia. Isoflurane was given at 1/2, 1/8 and 1/32 MAC for four in-utero exposures and for 24 exposures to the pups during the first nine weeks of life. The mice were killed at 15 months of age. The incidence of tumors in these mice was the same as in untreated control mice, which were given the same background gases, but not the anesthetic.

Mutagenesis

Isoflurane was negative in the *in vivo* mouse micronucleus and *in vitro* human lymphocyte chromosomal aberration assay. In published studies, isoflurane was negative in the *in vitro* bacterial reverse mutation assay (Ames test) in all strains tested (*Salmonella typhimurium* strains TA98, TA100, and TA1535) in the presence or absence of metabolic activation.

Impairment of Fertility

Male and female Sprague-Dawley rats were exposed to isoflurane at concentrations of 0%, 0.15%, and 0.60% (0, 1/8, and 1/2 MAC) 2 hours per day for 14 consecutive days prior to mating. Isoflurane had no effects on either male or female fertility.

13.2 Animal Toxicology and/or Pharmacology

Published studies in animals demonstrate that the use of anesthetic agents during the period of rapid brain growth or synaptogenesis results in widespread neuronal and oligodendrocyte cell loss in the developing brain and alterations in synaptic morphology and neurogenesis. Based on comparisons across species, the window of vulnerability to these changes is believed to correlate with exposures in the third trimester through the first several months of life, but may extend out to approximately 3 years of age in humans.

In primates, exposure to 3 hours of an anesthetic regimen that produced a light surgical plane of anesthesia did not increase neuronal cell loss, however, treatment regimens of 5 hours or longer increased neuronal cell loss. Data in rodents and in primates suggest that the neuronal and oligodendrocyte cell losses are associated with subtle but prolonged cognitive deficits in learning and memory. The clinical significance of these nonclinical findings is not known, and healthcare providers should balance the benefits of appropriate anesthesia in neonates and young children who require procedures against the potential risks suggested by the nonclinical data [*see Warnings and Precautions (5.3), Use in Specific Populations (8.1, 8.4)*].

16 HOW SUPPLIED/STORAGE AND HANDLING

Isoflurane USP, is available in the following presentations:

FILL	CONTAINER	NDC
100 mL	Amber-colored Bottle	66794-020-10
250 mL	Amber-colored Bottle	66794-020-25

Replace the cap securely after each use.

16.1 Safety and Handling

Occupational Caution

The following reactions have been reported following occupational exposure to isoflurane: dyspnea, bronchospasm, stridor, cough, dizziness, paresthesia, hepatic reactions, flushing rash, contact dermatitis, erythema, periorbital edema, eye irritation, conjunctival hyperemia, and headache.

There is no specific work exposure limit established for Isoflurane USP. However, the National Institute for Occupational Safety and Health Administration (NIOSH) recommends that no worker should be exposed at ceiling concentrations greater than 2 ppm of any halogenated anesthetic agent over a sampling period not to exceed one hour.

The predicted effects of acute overexposure by inhalation of Isoflurane USP include headache, dizziness or (in extreme cases) unconsciousness [see *Overdosage (10)*]. There are no documented adverse effects of chronic exposure to halogenated anesthetic vapors (Waste Anesthetic Gases or WAGs) in the workplace. Although results of some epidemiological studies suggest a link between exposure to halogenated anesthetics and increased health problems (particularly spontaneous abortion), the relationship is not conclusive. Since exposure to WAGs is one possible factor in the findings for these studies, operating room personnel, and pregnant women in particular, should minimize exposure. Precautions include adequate general ventilation in the operating room, the use of a well-designed and well-maintained scavenging system, work practices to minimize leaks and spills while the anesthetic agent is in use, and routine equipment maintenance to minimize leaks.

16.2 Storage

Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature]. Preserve in tight containers. Isoflurane contains no additives and has been demonstrated to be stable at room temperature for periods in excess of five years.

17. PATIENT COUNSELING INFORMATION

Anesthesia providers need to obtain the following information from patients prior to administration of anesthesia:

- Medications they are taking, including herbal supplements
- Drug allergies, including allergic reactions to anesthetic agents (including hepatic sensitivity)
- Any history of severe reactions to prior administration of anesthetic

- If the patient or a member of the patient's family has a history of malignant hyperthermia or if the patient has a history of Duchenne muscular dystrophy or other latent neuromuscular disease.

Anesthesia providers should inform patients of the following risks associated with Isoflurane USP:

- Post-operative nausea and vomiting and respiratory adverse effects including coughing.

There is no information of the effects of Isoflurane USP following anesthesia on the ability to operate an automobile or other heavy machinery. However, patients should be advised that the ability to perform such tasks may be impaired after receiving anesthetic agents. Patients should not undertake hazardous tasks, such as driving, for at least 24 hours following administration of a general anesthetic.

Anesthesia providers should inform parents and caregivers of pediatric patients that emergence from anesthesia in children may evoke a brief state of agitation that may hinder cooperation.

Anesthesia providers should inform patients that Isoflurane USP, as well as other general anesthetics, may cause a slight decrease in intellectual function for 2 or 3 days following anesthesia. As with other anesthetics, small changes in moods and symptoms may persist for up to 6 days after administration.

Effect of anesthetic and sedation drugs on early brain development

Studies conducted in young animals and children suggest repeated or prolonged use of general anesthetic or sedation drugs in children younger than 3 years may have negative effects on their developing brains. Discuss with parents and caregivers the benefits, risks, and timing and duration of surgery or procedures requiring anesthetic and sedation drugs [see *Warnings and Precautions (5.8)*].

Manufactured by:

Piramal Critical Care, Inc.

3950 Schelden Circle

Bethlehem, PA 18017, USA

(888) 822-84

Revised: November 2025

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL-100ml

NDC 66794-020-10

ISOFLURANE, USP

LIQUID FOR INHALATION

Rx Only

100 mL

Exp. Date:
 Mfg. Date:
 LOT No.:

Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature]. Preserve in tight containers. Replace the cap securely after each use.

IMPORTANT: Read accompanying product information for directions pertaining to use of Isoflurane, USP.

Contains: Isoflurane (1-chloro-2,2,2-trifluoroethyl difluoromethyl ether). A nonflammable, nonexplosive inhalation anesthetic.

NDC 66794-020-10 **Rx Only**

Isoflurane, USP
Liquid for Inhalation

100 mL


 3 6 6 7 9 4 0 2 0 1 0 6

Piramal
 Critical Care
 Manufactured by:
Piramal Critical Care, Inc.
 3950 Schelden Circle
 Bethlehem, PA 18017, USA
 (888) 822-8431

20688647
 34040603

Non Varnish Coding Area
 30 x 50.8 mm
 Prefixes are preprinted
 Batch info to be printed on line

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL-250ml

NDC 66794-020-25

ISOFLURANE, USP

LIQUID FOR INHALATION

Rx Only

250 mL

Exp. Date:
 Mfg. Date:
 LOT No.:

Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature]. Preserve in tight containers. Replace the cap securely after each use.

IMPORTANT: Read accompanying product information for directions pertaining to use of Isoflurane, USP.

Contains: Isoflurane (1-chloro-2,2,2-trifluoroethyl difluoromethyl ether). A nonflammable, nonexplosive inhalation anesthetic.

NDC 66794-020-25 **Rx Only**

Isoflurane, USP
Liquid for Inhalation

250 mL


 3 6 6 7 9 4 0 2 0 2 5 0

Piramal
 Critical Care
 Manufactured by:
Piramal Critical Care, Inc.
 3950 Schelden Circle
 Bethlehem, PA 18017, USA
 (888) 822-8431

20688646
 34041103

Non Varnish Coding Area
 76 X 40 mm
 Prefixes are preprinted
 Batch info to be printed on line

ISOFLURANE

isoflurane liquid

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66794-020
Route of Administration	RESPIRATORY (INHALATION)		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOFLURANE (UNII: CYS9AKD70P) (ISOFLURANE - UNII:CYS9AKD70P)	ISOFLURANE	1 mL in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66794-020-10	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/13/2013	
2	NDC:66794-020-25	250 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	11/13/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074416	11/13/2013	

Labeler - Piramal Critical Care, Inc. (805600439)

Revised: 11/2025

Piramal Critical Care, Inc.