

RITE AID MAXIMUM STRENGTH- benzocaine gel, dentifrice
Rite Aid

5820630 Rite Aid Max Str Oral Pain Relief Gel (8015316)

Drug Facts

Active ingredient

Benzocaine 20%

Purpose

Oral anesthetic

Use

temporarily relieves pain associated with mouth and gum irritations

Warnings

Methemoglobinemia warning: use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

Do not use

- more than directed
- for more than 7 days unless told to do so by a dentist or doctor
- for teething
- in children under 2 years of age

Stop use and ask a doctor if

- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- symptoms do not improve in 7 days
- allergic reaction occurs

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to open tube, cut tip of tube on score mark
- adults and children 2 years of age and older: apply to affected area up to 4 times

daily or as directed by dentist/doctor

- children under 12 years of age: adult supervision should be given in the use of this product
- children under 2 years of age: do not use
- for denture irritation apply a thin layer to the affected area; do not reinsert dental work until irritation/pain is relieved; rinse mouth well after reinserting

Other information

store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients Blue 1, Carbomer, Flavor, Glycerin, Polyethylene Glycol, Propylene Glycol, Red 40, Sodium Saccharin, Sorbic Acid, Water, Yellow 10

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE

CAMP HILL, PA 17011

RITE AID® PHARMACY

MAXIMUM STRENGTH

oral pain relief

benzocaine 20%

anesthetic gel

instant pain relief for:

toothaches

gum pain

canker sores

denture pain

NET WT 0.33 OZ (9 g)



RITE AID MAXIMUM STRENGTH

benzocaine gel, dentifrice

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0316
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	20 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B6978945GQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBIC ACID (UNII: X045WJ989B)	
GLYCERIN (UNII: PDC6A3C0OX)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

WATER (UNII: 059QF0KO0R)

**CARBOMER HOMOPOLYMER TYPE B (ALLYL PENTAERYTHRITOL OR ALLYL SUCROSE
CROSSLINKED)** (UNII: K6MOM3T5YL)

Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT (Characteristic Mint)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0316-4	1 in 1 CARTON	03/14/2024	
1		9 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	01/20/2011	

Labeler - Rite Aid (014578892)

Registrant - Lornamead Inc. (078584069)

Establishment

Name	Address	ID/FEI	Business Operations
Lornamead Inc.		080046418	manufacture(11822-0316)

Revised: 1/2025

Rite Aid