

FAMOTIDINE- famotidine tablet
Bi-Mart

Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert:

Do not use if you are allergic to famotidine or other acid reducers.

Do not use

- with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent wheezing, particularly with heartburn
- frequent **chest pain**
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens

- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **adults and children 12 years and over:**
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- **children under 12 years:** ask a doctor

Other information

- read the directions and warnings before use
- store at 20° - 25°C (68° - 77°F)
- protect from moisture

Inactive ingredients

Hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sodium starch glycolate, polyethylene glycol, talc, titanium dioxide, triacetin.

Questions?

call toll-free **1-844-912-4012**

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL

2 65116 25517 0

BI-MART

NDC 37835-165-50

Compare to the active ingredient in Maximum Strength PEPICID AC**

ACID REDUCER

FAMOTIDINE 20 MG

MAXIMUM STRENGTH

PREVENTS & RELIEVES HEARTBURN DUE TO ACID INDIGESTION

ACID REDUCER 50 TABLETS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each tablet)

Purpose

Famotidine 20 mg.....

Acid reducer

Uses

■ relieves heartburn associated with acid indigestion and sour stomach

■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.

Do not use ■ with other acid reducers

■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness

■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

■ frequent wheezing, particularly with heartburn ■ frequent chest pain

■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ kidney disease

PEEL FOR DIRECTIONS 165-50-BM 02/25

LIFT HERE

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

■ pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ adults and children 12 years and over: ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.

■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn

■ do not use more than 2 tablets in 24 hours

■ children under 12 years: ask a doctor

Other information

■ read the directions and warnings before use

■ store at 20°-25°C (68°-77°F)

■ protect from moisture

Inactive ingredients

Hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark Maximum Strength PEPICID AC®

Distributed by:

BI-MART

Eugene, OR 97402

BI-MART UNCOMMITTAL MONEY BACK GUARANTEE: If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

HINGE

FAMOTIDINE			
famotidine tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-165
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)		FAMOTIDINE	20 mg
Inactive Ingredients			
Ingredient Name			Strength
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
STARCH, CORN (UNII: O8232NY3SJ)			
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)			

TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
TRIACETIN (UNII: XHX3C3X673)				
Product Characteristics				
Color	white (white to off-white)		Score	no score
Shape	ROUND		Size	6mm
Flavor			Imprint Code	V;15
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-165-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA215822		06/01/2025	

Labeler - Bi-Mart (027630078)

Registrant - Bi-Mart (027630078)

Establishment			
Name	Address	ID/FEI	Business Operations
VKT Pharma Private Limited		871408062	manufacture(37835-165)

Revised: 6/2025

Bi-Mart