

HEALTH MART ARTHRITIS PAIN RELIEF- acetaminophen tablet, film coated, extended release

Strategic Sourcing Services LLC

McKesson Arthritis Pain Relief Drug Facts

Active ingredient (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,
ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults	<ul style="list-style-type: none">• take 2 caplets every 8 hours with water• swallow whole; do not crush, chew, split or dissolve• do not take more than 6 caplets in 24 hours• do not use for more than 10 days unless directed by a doctor
under 18 years of age	<ul style="list-style-type: none">• ask a doctor

Other information

- store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F).

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Tylenol® 8HR Arthritis Pain active ingredient

650 mg

Arthritis Pain Relief

Acetaminophen Extended-Release Tablets 650 mg

Pain Reliever/Fever Reducer

For the temporary relief of minor arthritis pain

Gluten Free

SEE NEW WARNING

50 Caplets*-650 mg Each *Capsule-Shaped Tablets

1	NDC:62011-0338-1	1 in 1 CARTON	05/30/2017	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA075077		05/30/2017	

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 8/2019

Strategic Sourcing Services LLC