

XT-1 PRE-POST PART B- sodium chlorite solution solution
AgroChem Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

XT-1 Pre-Post Part B

XT-1 PRE POST |PART B

DANGER

CAUSES SEVERE SKIN AND EYE IRRITATIONS

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

IF ON SKIN(or hair): Take off immediately all contaminated clothing. Rinse skin with water/shower. IF INHALED: Remove person to fresh air and keep

comfortable for breathing. Immediately call a POISON

CENTER/doctor. IF IN EYES: Rinse cautiously with water for several

minutes. Remove contact lenses, if present and easy to do. Continue

rinsing. Immediately call a POISON CENTER/doctor. Wash

contaminated clothing before reuse.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

INGREDIENTS:

Sodium Chlorite - 0.45%

XT-1 PRE POST is a two-part Chlorine Dioxide teat dip or spray and can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

Directions:

NOTE: MUST BE MIXED PRIOR TO USE. MIX PART A & PART Bin equal parts.

CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing , and will continue to generate minimum Chlorine Dioxide of 1000 ppm for up to 5-7 days.

Directions:

General Use

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

XT-1 PRE POST | PART B

TEAT DIP

DANGER

CAUSES SEVERE SKIN AND EYE IRRITATIONS



**FOR FARM AND INDUSTRIAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
KEEP OUT OF DIRECT SUNLIGHT.
AVOID FREEZING**

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water/shower. **IF INHALED:** Remove person to fresh air and keep comfortable for breathing. Immediately call a POISON CENTER/doctor.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER/doctor. Wash contaminated clothing before reuse.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS:

Sodium Chlorite - 0.45%

NET CONTENTS:

LOT#:

EXP:

XT-1 PRE POST is a two-part Chlorine Dioxide teat spray that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

Directions:

NOTE: MUST BE MIXED PRIOR TO USE. MIX PART A & PART B in equal parts.

CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing, and will continue to generate minimum Chlorine Dioxide of 1000ppm for up to 5-7 days.

General Use

Teat Sprays can also be foamed or sprayed using recommended equipment.

If using a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Pre Dipping Use

Perform for stripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, the full length of each teat with the product.

Allow product to stand for 30 seconds. Wipe teats dry after application, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip, foam, or spray, the full length of each teat. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the product should be applied to the teats, once a day for 3-4 days after the last milking.

Applications should resume at least twice daily one week before calving.



AgroChem INC.
FARM & DAIRY PRODUCTS

26 Freedom Way • Saratoga Springs, NY 12866 • (518) 226-4850

XT-1 PRE-POST PART B

sodium chlorite solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:17307-3480
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII: Z63H374SB6)	SODIUM CHLORITE	0.014526 in 1 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3480-1	19 kg in 1 CONTAINER		
2	NDC:17307-3480-2	57 kg in 1 PACKAGE		
3	NDC:17307-3480-3	208 kg in 1 DRUM		
4	NDC:17307-3480-4	1040 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		07/02/2018	

Labeler - AgroChem Inc (805427700)

Registrant - AgroChem Inc (805427700)

Establishment

Name	Address	ID/FEI	Business Operations
AgroChem Inc		805427700	api manufacture, manufacture

Revised: 7/2018

AgroChem Inc