

**NIGHTTIME HBP COLD AND FLU RELIEF- acetaminophen, dextromethorphan
hbr, doxylamine succinate solution
CVS PHARMACY**

CVS 44-013 HBP

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 30 mg
Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - runny nose and sneezing
 - sore throat
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- liver disease
- glaucoma
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years: do not use

Other information

- **each 30 mL contains:** sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, high fructose corn syrup, polyethylene glycol 400, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

♥CVS

Health®

Compare to the active ingredients
in Vicks® NyQuil® HBP Cold & Flu*

Decongestant Free

Multi-Symptom

Nighttime

High Blood

Pressure

COLD & FLU

ACETAMINOPHEN

Pain reliever, Fever reducer

DEXTROMETHORPHAN HBr

Cough suppressant

DOXYLAMINE SUCCINATE

Antihistamine

For people with high blood pressure

Relieves: Aches & fever, Runny nose,
Sneezing, Cough

Cherry

Flavor

Ages 12 Years & Over

12 FL OZ (355 mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING**

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® HBP Cold & Flu.

Distributed by : CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

© 2024 CVS/pharmacy

CVS.com®

1-800-SHOP CVS

V-19849

50844

ORG032301302

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

CVS Health Compare to the active ingredients in Vicks® NyQuil® HBP Cold & Flu*
Decongestant Free

**Multi-Symptom
Nighttime
High Blood
Pressure**

COLD & FLU

ACETAMINOPHEN
Pain reliever, Fever reducer
DEXTROMETHORPHAN HBR
Cough suppressant
DOXYLAMINE SUCCINATE
Antihistamine
For people with high blood pressure
Relieves: Aches & fever, Runny nose, Sneezing, Cough

Cherry Flavor
Ages 12 Years & Over
12 FL OZ (355 mL)

Drug Facts PEEL BACK FOR COMPLETE DRUG FACTS
TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

Active ingredients (in each 30 mL) Purpose
Acetaminophen 650 mg Pain reliever/fever reducer
Dextromethorphan Hbr 30 mg Cough suppressant
Doxylamine succinate 12.5 mg Antihistamine

Uses temporarily relieves common cold and flu symptoms:
■ headache ■ minor aches and pains
■ sore throat ■ runny nose and sneezing
■ cough due to minor throat and bronchial irritation
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® HBP Cold & Flu.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2024 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-19849

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

#442927
FPO 80%
UPC# 050428082423
X XXXXXX XXXXXX X

no print / no varnish area
lot no. & exp. date

Drug Facts (continued)
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have
■ difficulty in urination due to enlargement of the prostate gland ■ liver disease ■ glaucoma
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus)
Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers
When using this product
■ marked drowsiness may occur
■ excitability may occur, especially in children
■ avoid alcoholic beverages
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ use caution when driving a motor vehicle or operating machinery
Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days

Drug Facts (continued)
■ redness or swelling is present
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
■ do not take more than directed
■ do not exceed 4 doses per 24 hours
■ mL = milliliter
■ only use the dose cup provided
■ adults and children 12 years and over: 30 mL every 6 hours
■ children under 12 years: do not use
Other information
■ each 30 mL contains: sodium 13 mg
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package
Inactive ingredients
anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, high fructose corn syrup, polyethylene glycol 400, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sucralose
Questions or comments?
1-800-426-9391

CVS 44-013

NIGHTTIME HBP COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-086
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	red (maroon)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-086-02	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/04/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/04/2020	

Labeler - CVS PHARMACY (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(69842-086) , pack(69842-086)

Revised: 4/2025

CVS PHARMACY