

MUCINEX FAST-MAX SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH-
dextromethorphan hydrobromide, guaifenesin capsule, liquid filled
RB Health (US) LLC

Mucinex Fast-Max Severe Congestion & Cough

Drug Facts

Active ingredients (in each liquid gel)

Dextromethorphan HBr 10 mg
Guaifenesin 200 mg

Purposes

Cough suppressant
Expectorant

Keep out of reach of children.

In case of overdose, get medical help or contact a
Poison Control Center right away.

Dextromethorphan HBr 10 mg....Cough suppressant
Guaifenesin 200mg.....Expectorant

Uses

Uses:

■ helps loosen phlegm (mucus) and thin bronchial
secretions to rid the bronchial passageways of
bothersome mucus and make coughs more
productive

Temporarily relieves:

- cough due to minor throat and bronchial
irritation as may occur with the common cold or
inhaled irritants
- the intensity of coughing
- the impulse to cough to help you get to sleep

Do not use:

■ if you are now taking a prescription monoamine
oxidase inhibitor (MAOI) (certain drugs for
depression, psychiatric, or emotional conditions,
or Parkinson's disease), or for 2 weeks after
stopping the MAOI drug. If you do not know if your
prescription drug contains an MAOI, ask a doctor
or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with
smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if

■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 12 capsules in any 24-hour period
- adults and children 12 years of age and over: take 2 capsules every 4 hours
- children under 12 years of age: do not use
- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

D&C yellow no.10, FD&C blue no. 1, gelatin, glycerin, hypromellose, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

Fast Dissolving Liquid Gels!

NDC 72854-153

MAXIMUM STRENGTH

MUCINEX®

FAST-MAX®

MAXIMUM STRENGTH **NEW**

Mucinex[®]

FAST-MAX[®]

SEVERE CONGESTION & COUGH

| Drug Facts | Purposes |
|--|----------|
| Active ingredients (in each capsule) Dextromethorphan HBr 10 mg...Cough suppressant Guaifenesin 200 mg.....Expectorant | |
| Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily relieves: ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ■ the intensity of coughing ■ the impulse to cough to help you get to sleep | |
| Warnings Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus) When using this product do not use more than directed Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. | |

Drug Facts (continued)
if pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ do not take more than 12 capsules in any 24-hour period
■ adults and children 12 years of age and over: take 2 capsules every 4 hours
■ children under 12 years of age: do not use

Other information
■ store at 20-25°C (68-77°F)
■ avoid excessive heat

Inactive ingredients
D&C yellow no.10, FD&C blue no. 1, gelatin, glycerin, hypromellose, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

reckitt
Patents: www.reckitt.com/patents
Dist. by: RB Health (US)
Parsippany, NJ 07054-0224 081424 3311321
©2024 RB Health

FAST DISSOLVING LIQUID GELS **40% SMALLER*** ACTUAL SIZE

MAXIMUM STRENGTH **NEW**

Mucinex[®]

FAST-MAX[®]

SEVERE CONGESTION & COUGH

Dextromethorphan HBr » Cough Suppressant
Guaifenesin » Expectorant

✓ Controls Cough
✓ Relieves Chest Congestion
✓ Thins & Loosens Mucus

16 LIQUID GELS (Liquid Filled Capsules) **40% SMALLER*** **AGES 12+**

*Compared to Maximum Strength Mucinex Fast-Max Cold & Flu Liquid Gels
Scan for FAQs and instructions on proper disposal of medications or visit www.mucinex.com/FAQs
Please visit our website www.mucinex.com

Do not take more than a total of 12 capsules in any 24-hour period.
Take only as directed.
Keep carton for full information.

NDC 72854-153-16

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

PARENTS:
Learn about teen medicine abuse
www.SupplyMedicineAbuse.org

how2recycle.info
TRAY
MULTI-WATER
PAPER
BOX

638249983918

MUCINEX FAST-MAX SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH

dextromethorphan hydrobromide, guaifenesin capsule, liquid filled

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:72854-153 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------------------|----------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| POVIDONE (UNII: FZ989GH94E) | |
| WATER (UNII: 059QF0KO0R) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |

| | |
|---|--|
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| LECITHIN, SOYBEAN (UNII: 1DI56QDM62) | |
| LIGHT MINERAL OIL (UNII: N6K5787QVP) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| GELATIN (UNII: 2G86QN327L) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| SORBITOL SOLUTION (UNII: 8KW3E207O2) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | green | Score | no score |
| Shape | OVAL | Size | 16mm |
| Flavor | | Imprint Code | AR08 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:72854-153-16 | 2 in 1 CARTON | 06/01/2024 | |
| 1 | | 8 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 06/01/2024 | |

Labeler - RB Health (US) LLC (081049410)