

ASPIRIN- aspirin tablet, coated
Marc Glassman, Inc.

1018 - MAR - 2018-1206

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID ¹)

¹ nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves:
 - headache
 - muscle pain
 - toothache
 - menstrual pain
 - pain of colds
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose

- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, polyethylene glycol, pregelatinized starch, starch, titanium dioxide

PRINCIPAL DISPLAY PANEL

Marc's®

NDC 68998-018-11

†Compare to the active ingredient in Bayer® Aspirin Tablets

Regular Strength

Aspirin

Pain Reliever/Fever Reducer (NSAID)

Acutal Size

100 COATED TABLETS 325 MG EACH

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer Aspirin Tablets.

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin, 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves:
 ■ headache ■ menstrual pain
 ■ muscle pain ■ pain and fever of colds
 ■ toothache ■ minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction which may include:
 ■ hives
 ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product.
 ■ take more or for a longer time than directed
 Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.
Ask a doctor before use if
 ■ stomach bleeding warning applies to you

Drug Facts (continued)

■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you are taking a diuretic
 ■ you have asthma
Ask a doctor or pharmacist before use if you are
 taking a prescription drug for:
 ■ diabetes ■ gout ■ arthritis
 ■ blood thinners ■ heart medicine
Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts for more than 10 days
 ■ redness or swelling is present
 ■ fever gets worse or lasts for more than 3 days
 ■ new symptoms occur
 ■ ringing in the ears or loss of hearing occurs
 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Directions ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours; not to exceed 12 tablets in 24 hours
 ■ children under 12 years: consult a doctor

Drug Facts (continued)
Other information (68-77°F) in a dry place
 ■ store between 20-25°C
 ■ rebin caps for complete product information and warnings
Inactive ingredients
 corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide
 Distributed by:
 Marc Classman, Inc.
 West 130th Street
 Cleveland, OH 44130

Marc's NDC 68998-018-11
 †Compare to the active ingredient in Bayer® Aspirin Tablets

Regular Strength
Aspirin
 PAIN RELIEVER / FEVER REDUCER (NSAID)
 Actual Size
 100 COATED TABLETS - 325 MG EACH

Marc's NDC 68998-018-11
 †Compare to the active ingredient in Bayer® Aspirin Tablets

Regular Strength
Aspirin
 PAIN RELIEVER / FEVER REDUCER (NSAID)
 Actual Size
 100 COATED TABLETS - 325 MG EACH

COATING FREE AREA

ASPIRIN
 aspirin tablet, coated

| Product Information | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68998-018 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 325 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| MINERAL OIL (UNII: T5L8T28FGP) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|----------|-------|--------------|----------|
| Color | white | Score | no score |
| Shape | ROUND | Size | 10mm |
| Flavor | | Imprint Code | TCL;011 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:68998-018-11 | 1 in 1 CARTON | 11/01/2009 | |
| 1 | | 100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| 2 | NDC:68998-018-14 | 1 in 1 CARTON | 11/01/2009 | 10/30/2027 |
| 2 | | 300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013 | 11/01/2009 | |

Labeler - Marc Glassman, Inc. (094487477)