

ASPIRIN 81MG ADULT LOW DOSE- aspirin tablet, delayed release TopCare

TCR - 1080B - 2019-1006

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for this product

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

PRINCIPAL DISPLAY PANEL

2 120CT BOTTLES INSIDE

TWIN PACK

TopCare® health

NDC 36800-880-08

COMPARE TO BAYER® LOW DOSE ASPIRIN ACTIVE INGREDIENT†

LOW DOSE

ASPIRIN REGIMEN**

Aspirin 81mg

PAIN RELIEVER (NSAID)

Safety Coated

OUR PHARMACISTS RECOMMEND

240 TOTAL ENTERIC COATED TABLETS

actual size

<p>DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING</p> <p>Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate</p> <p>Drug Facts (continued) ■ retain carton for complete product information and warnings</p>	<p>Other information ■ store between 20°-25°C (68°-77°F) in a dry place</p> <p>Directions ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued) ■ take a blood thinning (anticoagulant) or steroid drug ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ shock ■ asthma (wheezing) ■ facial swelling</p> <p>early sign of Raye's syndrome, a rare but serious illness. Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.</p>
<p>240 TOTAL ENTERIC COATED TABLETS</p> <p>Aspirin 81mg</p> <p>LOW DOSE ASPIRIN REGIMEN**</p> <p>PAIN RELIEVER (NSAID)</p> <p>TopCare® health</p>	<p>Drug Facts ■ retain carton for complete product information and warnings</p> <p>Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate</p> <p>Other information ■ store between 20°-25°C (68°-77°F) in a dry place</p> <p>Directions ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued) ■ take a blood thinning (anticoagulant) or steroid drug ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ shock ■ asthma (wheezing) ■ facial swelling</p> <p>early sign of Raye's syndrome, a rare but serious illness. Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.</p>
<p>240 TOTAL ENTERIC COATED TABLETS</p> <p>Aspirin 81mg</p> <p>LOW DOSE ASPIRIN REGIMEN**</p> <p>PAIN RELIEVER (NSAID)</p> <p>TopCare® health</p>	<p>Drug Facts ■ retain carton for complete product information and warnings</p> <p>Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate</p> <p>Other information ■ store between 20°-25°C (68°-77°F) in a dry place</p> <p>Directions ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Drug Facts (continued) ■ take a blood thinning (anticoagulant) or steroid drug ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ shock ■ asthma (wheezing) ■ facial swelling</p> <p>early sign of Raye's syndrome, a rare but serious illness. Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.</p>

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
© TOPCO AP00201 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin. Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
BROWN IRON OXIDE (UNII: 1N032N7MFO)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	heart
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-880-08	2 in 1 CARTON	10/03/2016	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:36800-880-18	2 in 1 CARTON	03/21/2019	
2		365 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/03/2016	

Labeler - TopCare (006935977)

Revised: 12/2024

TopCare