

**TRIPLE ANTIBIOTIC PLUS LIDOCAINE- bacitracin zinc, neomycin sulfate ,
polymyxin b sulfate, lidocaine ointment
TRIFECTA PHARMACEUTICALS USA LLC**

Triple Antibiotic Ointment + Lidocaine

Active ingredients (each gram contains)

Bacitracin zinc 500 units
Neomycin sulfate 3.5 mg
Polymyxin B sulfate 10,000 units

Purpose

First aid antibiotic

Active Ingredient

Lidocaine 40mg

Purpose

External Analgesic

Uses

Uses first aid to help prevent infection and for the temporary relief of pain or discomfort in minor:● cuts ● scrapes ● burns

Keep out of reach of children

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Warnings

For external use only.

Ask Doctor Before use if you have

Ask Doctor before use if you have

- Deep or puncture wounds
- Animal bites
- serious burns

Do Not Use

Do Not Use

- If you are allergic to any of the ingredients
- In the eyes
- Over large areas of the body
- In large quantities, particularly over raw surfaces or blistered areas

Stop Use and Ask a doctor if

Stop Use and Ask a doctor if

- You need to use longer than 1 week
- Condition persists or gets worse
- Symptoms persist for more than 1 week or clear up and occur again within a few days
- Rash or other allergic reaction develops

Directions

- Adults and Children 2 years of age and older
- Clean the affected area
- Apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- May be covered with a sterile bandage
- Children under 2 years of age, Ask a doctor

Questions

Questions? Call 1-888-296-9067

Inactive Ingredient

Petrolatum

Other information

- store at 20° to 25°C (68° to 77°F)

Distributed By

Distributed by:

Trifecta Pharmaceuticals USA®

101 NE Third Avenue, Suite 1500

Ft. Lauderdale, FL 33301, USA

1-888-296-9067

* This product is not manufactured or distributed by Johnson & Johnson Corporation,

owner of the registered trademark Neosporin® plus Lidocaine
Lidocaine known to numb in 30 to 60 minutes.

Package Label

OUTSIDE BOX



Triple Antibiotic Ointment
+ LIDOCAINE

4

PACK

*COMPARE TO THE ACTIVE INGREDIENTS IN NEOSPORIN® PLUS LIDOCAINE



NDC: 69396-154-44

4

PACK

Triple Antibiotic Ointment
+ LIDOCAINE

BACITRACIN ZINC • LIDOCAINE
NEOMYCIN SULFATE • POLYMYXIN B SULFATE

With Fast-Acting Lidocaine for Pain Relief*
Numbs Away the Pain

Net Wt. 4 oz. (113.6 g)

FIRST AID + LIDOCAINE PAIN RELIEF

LOT:
EXP:

With Fast-Acting Lidocaine for Pain Relief*
Numbs Away the Pain
First Aid Antibiotic/Pain Relieving Ointment





X0049L8467
Globe (4-Pack) Lidocaine Fl...y Use & Infection Protection
New

Drug Facts

Each gram contains 500 units Lidocaine, 40 mg Neomycin B Sulfate, 3 mg Polymyxin B Sulfate, 10,000 units First aid antibiotic First aid antibiotic

Purpose

External analgesic First aid antibiotic

Warnings

For external use only.

Do not use • if you are allergic to any of the ingredients • in the eyes • over large areas of the body

• in large quantities, particularly over raw surfaces or blistered areas

Ask a doctor before use if you have • deep or puncture wounds • animal bites • serious burns

Stop use and ask a doctor if • you need to use longer than 1 week • condition persists or gets worse

• symptoms persist for more than 1 week, or clear up and occur again within a few days • a rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 2 years of age and older:

- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily

Children under 2 years of age: ask a doctor

Other information Store at 20° to 25°C (68° to 77°F).

Distributed by:

Tritech Pharmaceuticals USA
100 NE Third Avenue, Suite 1500
Fort Lauderdale, FL 33301 USA
www.tritech-pharma.com
SKU: 408104-FBA

* This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Neosporin Plus Lidocaine.

† Lidocaine known to numb in 30 to 60 minutes.

Questions? Call 1-888-296-9067

Label

OUTSIDE BOX

Fast-Acting Lidocaine for Pain Relief*
Numbs Away the Pain
First Aid Antibiotic/Pain Relieving Ointment

COMPARE TO THE ACTIVE INGREDIENTS IN NEOSPORIN® PLUS LIDOCAINE

NDC: 69396-154-22

2 PACK

Globe®

Triple Antibiotic Ointment
 BACITRACIN ZINC • LIDOCAINE • NEOMYCIN SULFATE • POLYMYXIN B SULFATE
First Aid Antibiotic/Pain Relieving Ointment

Net Wt. 2 oz. (56.8 g)

+ LIDOCAINE

LOT: _____
 EXP: _____

Distributed by:
Trifecta Pharmaceuticals USA*
 101 NE Third Avenue, Suite 1500
 Ft. Lauderdale, FL 33301 USA
www.trifecta-pharma.com
 SKU: 408102

Globe®
 A symbol of pharmacy skill and professionalism.

X004906SMR

Globe (2 Pack) First Aid Ant...ay The Pain, Bacitracin Zinc, New

Drug Facts (continued)

Active ingredients (each gram contains)

Bacitracin Zinc 500 units
 Lidocaine 40 mg
 Neomycin Sulfate 3.5 mg
 Polymyxin B Sulfate 10,000 units

Uses

First aid to help prevent infection and for temporary relief of pain or discomfort in minor:
 ■ Cuts ■ Scrapes ■ Burns

Warnings

Do not use ■ if you are allergic to any of the ingredients ■ in the eyes ■ Over large areas of the body ■ in large quantities, particularly over raw surfaces or blistered areas ■ Ask a doctor before use if you have ■ Deep or puncture wounds ■ Animal bites ■ Serious burns

For external use only.

Directions

■ Adults and children 2 years of age and older: ■ Clean the affected area of a scrape or burn with soap and water. ■ Apply a small amount of the product (an amount equal to the surface area of the lip or a finger) on the area 1 to 3 times daily. ■ May be covered with a sterile bandage.

■ Children under 2 years of age: Ask a doctor.

Other information Store at 20° to 25°C (68° to 77°F).

Inactive ingredients Petroleum.

Questions? Call 1-888-296-9067

Drug Facts

Stop use and ask a doctor if ■ You need to use longer than 1 week ■ Condition persists or gets worse ■ Symptoms persist for more than 1 week, or clear up and occur again within a few days ■ A rash or other allergic reaction develops

Purpose

First aid antibiotic

Keep out of reach of children.

External analgesic

First aid antibiotic

Uses

First aid antibiotic

First aid antibiotic

If swallowed, get medical help or contact a Poison Control Center right away.


* This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Neosporin® Plus Lidocaine. Lidocaine known to numb in 30 to 60 minutes.

PACKAGE LABEL

OUTSIDE BOX



INNER TUBE



TRIPLE ANTIBIOTIC PLUS LIDOCAINE

bacitracin zinc, neomycin sulfate , polymyxin b sulfate, lidocaine ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69396-154
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [USP'U] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	40 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69396-154-01	1 in 1 BOX	03/31/2024	
1		28.4 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:69396-154-22	2 in 1 BOX	06/26/2024	
2		28.4 g in 1 TUBE; Type 0: Not a Combination Product		
3	NDC:69396-154-44	4 in 1 BOX	03/31/2024	
3		28.4 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M004	03/31/2024	

Labeler - TRIFECTA PHARMACEUTICALS USA LLC (079424163)

Registrant - Trifecta Pharmaceuticals USA (079424163)

Revised: 4/2025

TRIFECTA PHARMACEUTICALS USA LLC