

BENZINUM- benzene pellet
Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Benzinum 200CK HPUS

Benzinum 200CK HPUS

Active ingredient**: See product name on front panel (**contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Relieves fatigue with paleness *

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

lactose, sucrose

BoironUSA.com Info@boiron.com

1-800-BOIRON-1 (1-800-264-7661)

Distributed by Boiron, Inc.

Newtown Square, PA 19073

Do not use if pellet dispenser seal is broken.


Contains approx.80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

**C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

Benzinum 200ck
 Relieves fatigue with paleness *



Made in France HOMEOPATHIC MEDICINE
 HPUS NDC 0220-0817-41

Do not use if pellet dispenser seal is broken

LOT	EXP
3 06960	81741 3

Contains approx. 80 pellets. US
 Peel for Drugs Facts and instructions for use.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

Benzinum 200 ck

Drug Facts

Active ingredient:** See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions: ■ Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Drug Facts (continued)

Inactive ingredients: lactose, sucrose


Questions or comments?

BoironUSA.com Info@Boiron.com
 1-800-BOIRON-1 (1-800-264-7661)

**C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

Distributed by Boiron Inc.
 Newtown Square, PA 19073

How to dispense pellets?



1 Turn tube upside down.
 2 Twist until 5 pellets are dispensed into cap.
 3 Carefully remove cap and use it to pour pellets under the tongue.

BENZINUM

benzene pellet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-0817
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZENE (UNII: J64922108F) (BENZENE - UNII:J64922108F)	BENZENE	200 [kp_C] in 200 [kp_C]

Inactive Ingredients

Ingredient Name	Strength
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	white	Score	
Shape	ROUND	Size	4mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0220-0817-41	200 [kp_C] in 1 TUBE; Type 0: Not a Combination Product	01/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		01/01/2024	

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment

Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-0817)

Revised: 4/2024

Boiron