## LOW DOSE ASPIRIN- aspirin tablet, coated Walmart Inc.

-----

#### Low Dose Aspirin 81 mg Delayed-Release Tablets

#### **Drug Facts**

#### Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain reliever

#### Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

#### Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

#### Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur ringing in the ears or a loss of hearing occurs

#### If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

#### Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package
- avoid excessive heat above 40°C (104°F)

#### Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

#### Questions or comments?

Call 1-888-287-1915



# **Drug Facts** (continued)

take more or for a longer time than directed

**Do not use** ■ if you are allergic to aspirin or any other pain eaction to this product or any of its ingredients eliever/fever reducer ■ if you have ever had an allergic

diuretic ■ you have asthma Ask a doctor before use if stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a

Stop use and ask a doctor it prescription drug for ■ gout ■ diabetes ■ arthritis Ask a doctor or pharmacist before use if you are taking a

■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days an allergic reaction occurs. Seek medical help right away.
 you experience any of the following signs of stomach bleeding:
 feel faint vomit blood have bloody or black stools

redness or swelling is present

ringing in the ears or a loss of hearing occurs

new symptoms occur

so by a doctor because it may cause problems in the unborr use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do If pregnant or breast-feeding, ask a health professional before child or complications during delivery

**keep out of reach of children.** In case of overdose, get nedical help or contact a Poison Control Center right away

# Directions

drink a full glass of water with each dose

■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

■ children under 12 years: consult a doctor

Other information

 use by expiration date on package
 avoid excessive heat above 40°C (104°F) ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)

ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium laury #10 aluminum lake, iron oxide ochre, methacrylic acid and Inactive ingredients anhydrous lactose, carnauba wax colloidal silicon dioxide, croscarmellose sodium, D&C yellow sulfate, starch, talc, titanium dioxide, triethyl citrate

NDC 79903-481-50

**Actual Size** 

equate

Aspirin

**Safety Coated** 

Aspirin Regimen\*

81 mg Delayed-

**Release Tablets** Pain Reliever (NSAID)

500 TABLETS

**Questions or comments?** Call 1-888-287-1915

\*\*Talk to your doctor or other healthcare provider before using this product for your heart

Orua Facts

Purpose

ain reliever

pirin 81 mg

Active ingredient (in each tablet)

**US&S** ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this** product will not provide fast relief of headaches or other symptoms needing immediate relief

ask your doctor about other uses for enteric coated 81 mg Aspirin.

recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with symptoms could be an early sign of Reye's syndrome, a rare but nausea and vomiting occur, consult a doctor because these Reye's syndrome: Children and teenagers who have or are

Allergy alert: Aspirin may cause a severe allergic reaction which serious illness may include:

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ hives ■ facial swelling ■ asthma (wheezing)

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription take a blood thinning (anticoagulant) or steroid drug NSAIDs (aspirin, ibuprofen, naproxen, or others)


have 3 or more alcoholic drinks every day while using this product

*Drug Facts* (continued under l Bentonville, AR 72716
This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® DISTRIBUTED BY: Walmart Inc., Low Dose Aspirin. 481R 0524

. E EXP.:



801350



#### **LOW DOSE ASPIRIN**

aspirin tablet, coated

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:79903-481

Route of Administration ORAL

#### **Active Ingredient/Active Moiety**

Ingredient Name Basis of Strength Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)

ASPIRIN (UNII: R16CO5Y76E) ASPIRIN 81 mg

## Inactive Ingredients Ingredient Name Strength

SODIUM HYDROXIDE (UNII: 55X04QC32I)

CARNAUBA WAX (UNII: R12CBM0EIZ)

BROWN IRON OXIDE (UNII: 1N032N7MFO)

POLYSORBATE 80 (UNII: 60ZP39ZG8H)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

CROSCARMELLOSE SODIUM (UNII: M280L1HH48)

ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	

Product Characteristics			
Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	heart
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:79903-481- 11	2 in 1 CARTON	07/30/2024		
1		250 in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:79903-481- 50	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/30/2024		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/30/2024	

### Labeler - Walmart Inc. (051957769)

## Registrant - TIME CAP LABORATORIES INC (037052099)

Establishment				
Name	Address	ID/FEI	<b>Business Operations</b>	
TIME CAP LABORATORIES INC		037052099	manufacture(79903-481)	

Revised: 7/2024 Walmart Inc.