

**MUCINEX CHILDRENS FEVER SORE THROAT AND COUGH- acetaminophen,  
dextromethorphan hydrobromide solution  
RB Health (US) LLC**

-----  
**Mucinex Children's Fever, Sore Throat & Cough**

**Drug Facts**

**Active ingredients / Purpose  
(in each 10 mL)**

Acetaminophen 325 mg..Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg....Cough suppressant

Acetaminophen 325 mg..Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg....Cough suppressant

**Uses**

- temporarily relieves:
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help your child get to sleep
- minor aches and pains
- sore throat
- headache
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists

for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

### **Ask a doctor before use if the child has**

- liver disease
- chronic cough that lasts such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if the child is taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms

### **Directions**

#### **Directions**

- this product does not contain directions or complete warnings for adult use

- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

**Other information**

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

**Inactive ingredients:** anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions? 1-866-682-4639

NDC 72854-064

Children's  
Mucinex®

Honey  
Flavor Liquid

4 FL OZ  
(118 mL)

**Drug Facts**

**Active ingredients (in each 10 mL)** **Purposes**  
**Acetaminophen 325 mg...Pain reliever/fever reducer**  
 Dextromethorphan HBr 10 mg....Cough suppressant

**Uses**

- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help your child get to sleep
  - minor aches and pains
  - sore throat
  - headache
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if the child has**

- liver disease
- chronic cough that lasts such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if the child is taking the blood thinning drug warfarin**

**When using this product do not use more than directed (see Overdose warning)**

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

**Other information**

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

**Inactive ingredients** anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

**Questions? 1-866-682-4639**

'as an OTC Brand in the Children's Cough/Cold Category (AlphaImpactRxProVoice Survey)

**PARENTS:**  
 Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Keep carton for full information.

**Pediatrician Recommended** NDC 72854-064-04

**Mucinex Children's**

**FEVER, SORE THROAT & COUGH**

**Acetaminophen 325 mg - Pain Reliever/Fever Reducer**  
 Dextromethorphan HBr 10 mg - Cough Suppressant

- ✓ Reduces Fever
- ✓ Relieves Sore Throat
- ✓ Controls Cough

**Ages 6+ years**

**Honey Flavor Liquid**

**Each dose of Honey flavored Mucinex Children's Fever, Sore Throat & Cough Liquid:**

- ✓ Reduces Fever
- ✓ Relieves Sore Throat
- ✓ Controls Cough

**Dosing Cup Included**

Questions or comments?  
 1-866-MUCINEX (1-866-682-4639)  
 or [www.mucinex.com](http://www.mucinex.com)

Dist. by: RB Health (US)  
 Parsippany, NJ 07054-0224  
 ©2024 RB Health  
 Made in Mexico  
 120123 3288606 US patents: [www.reckitt.com/patents](http://www.reckitt.com/patents)

**reckitt**

4 FL OZ (118 mL)

## MUCINEX CHILDRENS FEVER SORE THROAT AND COUGH

acetaminophen, dextromethorphan hydrobromide solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72854-064
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 10 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	

<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	HONEY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-064-04	1 in 1 CARTON	05/01/2024	
1		118 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

**Labeler** - RB Health (US) LLC (081049410)

Revised: 4/2024

RB Health (US) LLC