

**GUAIFENESIN AND DEXTROMETHORPHAN HYDROBROMIDE- guaifenesin and dextromethorphan hydrobromide syrup**  
**Method Pharmaceuticals**

-----

**Guaifenesin and Dextromethorphan Hydrobromide Syrup**

**Active ingredients**

Dextromethorphan HBr, USP 10 mg  
Guaifenesin, USP 100 mg

**Purpose**

Cough suppressant  
Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- persistent cough or chronic cough such as occurs with smoking, asthma, chronic bronchitis, emphysema
- cough accompanied by excessive phlegm (mucus)

**Stop use and ask a doctor if**

- cough lasts more than 7 days or occurs with fever, rash, or headaches that lasts. This could be a sign of a serious condition
- hypersensitive to any ingredients

**If pregnant or breast-feeding,**

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- do not take more than 6 doses in any 24-hour period. **This adult product is not intended for use in children under 12 years of age**

adults and children 12 years and over	2 teaspoonfuls (TSP) every 4 hours
children 6 to 12 years of age	1 teaspoonful (TSP) every 4 hours
children under 6 years	<b>DO NOT USE</b>

## TAMPER-EVIDENT

Do not use if seal cap is broken or missing.

## Inactive ingredients

Anhydrous citric acid, dextrose, FD&C Red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate

## Other information

- store at room temperature 20°-25°C (68°-77°F)
- each 5 mL contains:** sodium 3 mg

## Questions

1-877-250-3427

## Principal Display Panel

Method Pharmaceuticals

NDC 58657-504-08

Guaifenesin and Dextromethorphan Hydrobromide Syrup

100 mg/10 mg/5 mL

Expectorant

Cough Suppressant

Grape Flavor

**Non Drowsy • Alcohol Free**

FOR AGES 12 AND UP

**8 fl. oz. (237 mL)**

Manufactured For:  
 Method Pharmaceuticals, LLC  
 Southlake, TX 76092 Rev. 02/2024



NDC 58657-504-08

## Guaifenesin and Dextromethorphan Hydrobromide Syrup

**Expectorant  
Cough Suppressant**

DO NOT USE IF TAMPER EVIDENT SEAL  
UNDER CAP IS BROKEN OR MISSING

8 fl. oz. (237mL)

<b>Drug Facts</b>	
<b>Active ingredient (in each 5 mL or TSP)</b> Dextromethorphan Hydrobromide, USP 10 mg ..... Cough Suppressant Guaifenesin, USP 100 mg ..... Expectorant	<b>Purpose</b>
<b>Uses</b> <ul style="list-style-type: none"> <li>■ temporarily relieves cough due to minor throat and bronchial irritation</li> <li>■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive</li> </ul>	
<b>Warnings</b> Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
<b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>■ persistent cough or chronic cough such as occurs with smoking, asthma, chronic bronchitis, emphysema</li> <li>■ cough accompanied by excessive phlegm (mucus)</li> </ul>	
<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>■ cough lasts more than 7 days or occurs with fever, rash, or headaches that lasts. This could be a sign of a serious condition</li> <li>■ hypersensitive to any ingredients</li> </ul>	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	

<b>Drug Facts (continued)</b>	
<b>Directions</b> ■ do not take more than 6 doses in any 24-hour period. <b>This adult product is not intended for use in children under 12 years of age</b>	
adults and children 12 years and over	2 teaspoonfuls (TSP) every 4 hours
children 6 to 12 years of age:	1 teaspoonful (TSP) every 4 hours
children under 6 years	<b>DO NOT USE</b>
<b>TAMPER-EVIDENT:</b> Do not use if seal under cap is broken or missing.	
<b>Inactive ingredients</b> Anhydrous citric acid, dextrose, FD&C Red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate	
<b>Other information</b> ■ store at room temperature 20°-25°C (68°-77°F) ■ each 5 mL contains: sodium 3 mg	
<b>Questions</b> 1-877-250-3427	
Manufactured For: <b>Method Pharmaceuticals, LLC</b> <span style="float: right;">LR-206</span> Southlake, TX 76092 <span style="float: right;">Rev. 02/2024</span>	
 N 3 58657 50408 3	

## GUAIFENESIN AND DEXTROMETHORPHAN HYDROBROMIDE

guaifenesin and dextromethorphan hydrobromide syrup

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58657-504
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>DEXTROSE</b> (UNII: IY9XDZ35W2)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-504-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/01/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/01/2024	

**Labeler** - Method Pharmaceuticals (060216698)

Revised: 10/2025

Method Pharmaceuticals