

**HEADACHE RELIEF- acetaminophen, aspirin, caffeine tablet**  
**Walmart Inc.**

-----  
**Walmart Acetaminophen 250 mg, Aspirin 250 mg, Caffeine 65 mg Tablets**

***Drug Facts***

***Active ingredients (in each caplet)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

***Purposes***

Pain reliever

Pain reliever

Pain reliever aid

***Uses***

- temporarily relieves minor aches and pains due to:

-headache

-a cold

-arthritis

-muscular aches

-toothache

-premenstrual and menstrual cramps

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

**Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**



NDC 79903-370-07

**equate™**

EXTRA STRENGTH

**Headache Relief**  
**Acetaminophen 250 mg**  
**Aspirin 250 mg (NSAID)**  
**and Caffeine 65 mg**

Pain Reliever /  
 Pain Reliever Aid

100  
 Caplets\*\*

(\*\*Capsule-Shaped Tablets)

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL**  
**OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active Ingredients (in each caplet)**

Acetaminophen 250 mg.....Pain reliever  
 Aspirin 250 mg (NSAID)\*.....Pain reliever  
 Caffeine 65 mg.....Pain reliever aid

\*nonsteroidal anti-inflammatory drug  
**Uses:** temporarily relieves minor aches and pains due to: headache, a cold, arthritis, muscular aches, toothache, premenstrual and menstrual cramps

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

**Keep out of reach of children.**  
 If pregnant or breast-feeding, ask a health professional before use.

**Directions** ■ Do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

**Other information** ■ store at 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

**Questions or comments?** Call 1-888-287-1915

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

803550

LOT: 370R 0524

EXP: Vamish Omit Area

**Drug Facts (continued)**

cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

**Do not use**  
 ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**  
 ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are**  
 ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug or are under a doctor's care for any serious condition

**Stop use and ask a doctor if**  
 ■ an allergic reaction occurs. Seek medical help right away.  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stools  
 ■ have stomach pain that does not get better  
 ■ ringing in the ears or loss of hearing occurs  
 ■ painful area is red or swollen  
 ■ pain gets worse or lasts for more than 10 days  
 ■ fever gets worse or lasts for more than 3 days  
 ■ any new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**  
 In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not use more than directed

■ drink a full glass of water with each dose

■ adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours

■ children under 12 years: ask a doctor

**Other information**

■ store at 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

**Inactive ingredients:** carnuba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call 1-888-287-1915

NDC 79903-370-20

**equate™**

EXTRA STRENGTH

**Headache Relief**  
**Acetaminophen 250 mg**  
**Aspirin 250 mg (NSAID)**  
**and Caffeine 65 mg**  
**Pain Reliever /**  
**Pain Reliever Aid**

200  
 CAPLETS\*\*

(\*\*Capsule-Shaped Tablets)

Actual Size

Compare to  
 Excedrin® Extra  
 Strength active  
 ingredients!

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg.....Pain reliever  
 Aspirin 250 mg (NSAID)\*.....Pain reliever  
 Caffeine 65 mg.....Pain reliever aid

\*nonsteroidal anti-inflammatory drug

**Uses**

■ temporarily relieves minor aches and pains due to:  
 ■ headache ■ a cold ■ arthritis ■ muscular aches  
 ■ toothache ■ premenstrual and menstrual cramps

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

**Keep out of reach of children.**  
 If pregnant or breast-feeding, ask a health professional before use.

**Directions** ■ Do not use more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

**Other information** ■ store at 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

**Questions or comments?** Call 1-888-287-1915

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

803550

LOT: 370R 0524

EXP: Vamish Omit Area

# HEADACHE RELIEF

acetaminophen, aspirin, caffeine tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79903-370
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (Caplet)	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL370
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-370-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	07/30/2024	
2	NDC:79903-370-21	2 in 1 CARTON	07/30/2024	
2	NDC:79903-370-01	100 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/30/2024	

**Labeler** - Walmart Inc. (051957769)

**Registrant** - TIME CAP LABORATORIES INC (037052099)

## Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES INC		037052099	manufacture(79903-370)

## Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(79903-370)

Revised: 11/2025

Walmart Inc.