

HEADACHE RELIEF- acetaminophen, aspirin, caffeine tablet
Walmart Inc.

Walmart Acetaminophen 250 mg, Aspirin 250 mg, Caffeine 65 mg Tablets

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:

-headache

-a cold

-arthritis

-muscular aches

-toothache

-premenstrual and menstrual cramps

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20°- 25°C (68°- 77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?* Call **1-888-287-1915*

equate™

EXTRA STRENGTH

Headache Relief
Acetaminophen 250 mg
Aspirin 250 mg (NSAID)
Pain Reliever /
Pain Reliever Aid

200
CAPLETS**

(**Capsule-Shaped Tablets)

Actual Size

NDC 79903-370-20

Compare to
Excedrin® Extra
Strength active
ingredients!

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID)* Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis ■ muscular aches
■ toothache ■ premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 8 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may

Drug Facts (continued under label)

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

DISTRIBUTED BY: Wal-Mart Inc.,
Bentonville, AR 72716
This product is not manufactured or
distributed by GSK Consumer Healthcare,
owner of the registered trademark,
Coaxinal Extra Strength. 370R 0524



LOT:

EXP:



equate™

EXTRA STRENGTH

Headache Relief
Acetaminophen 250 mg
Aspirin 250 mg (NSAID)
and Caffeine 65 mg

Pain Reliever /
Pain Reliever Aid

100
Caplets**

(**Capsule-Shaped Tablets)

NDC 79903-370-01

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL
OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID)* Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains due to: headache, a cold,
arthritis, muscular aches, toothache, premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from
chicken pox or flu-like symptoms should not use this product. When using this
product, if changes in behavior with nausea and vomiting occur, consult a
doctor because these symptoms could be an early sign of Reye's syndrome, a
rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may
include:
■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may
occur if you take:
■ more than 8 caplets in 24 hours, which is the maximum
daily amount ■ with other drugs containing acetaminophen
■ 3 or more
alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause
severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems ■ take a blood thinning
(anticoagulant) or steroid drug ■ take other drugs containing prescription or
nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or
more alcoholic drinks every day while using this product ■ take more or for a
longer time than directed

Caffeine warning: The recommended dose of this product contains about as
much caffeine as a cup of coffee. Limit the use of caffeine-containing
medications, foods, or beverages while taking this product because too much
caffeine may cause nervousness, irritability, sleeplessness, and, occasionally,
rapid heartbeat.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Directions ■ do not use more than directed ■ drink a full glass of water
with each dose ■ adults and children 12 years of age and over: take 2 caplets
every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years:
ask a doctor

Other information ■ store at 20° - 25°C (68° - 77°F); excursions permitted
between 15° - 30°C (59° - 86°F) ■ use by expiration date on package

Questions or comments? Call 1-888-287-1915

Satisfaction guaranteed - For questions or comments
please call 1-888-287-1915.

DISTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716

370R 0524 801350

LOT:

EXP:

Drug Facts (continued)

cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs
(aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about
as much caffeine as a cup of coffee. Limit the use of caffeine-containing
medications, foods, or beverages while taking this product because too
much caffeine may cause nervousness, irritability, sleeplessness, and,
occasionally, rapid heartbeat.

Do not use
■ If you have ever had an allergic reaction to acetaminophen, aspirin or
any other pain reliever/fever reducer
■ with any other drug containing acetaminophen (prescription or
nonprescription). If you are not sure whether a drug contains
acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if
■ you have liver disease ■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney
disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are
■ taking a prescription drug for diabetes, gout, or arthritis
■ taking any other drug or are under a doctor's care for any serious
condition

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ ringing in the ears or loss of hearing occurs
■ painful area is red or swollen
■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ any new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is
especially important not to use aspirin at 20 weeks or later in pregnancy
unless specifically directed to do so by a doctor because it may cause
problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center
(1-800-222-1222) right away. Quick medical attention is critical for adults
as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed
■ drink a full glass of water with each dose
■ adults and children 12 years of age and over: take 2 caplets every 6
hours; not more than 8 caplets in 24 hours
■ children under 12 years: ask a doctor

Other information
■ store at 20° - 25°C (68° - 77°F); excursions permitted between 15° - 30°C
(59° - 86°F) ■ use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide,
hypromellose, microcrystalline cellulose, mineral oil, povidone,
pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-888-287-1915



HEADACHE RELIEF

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79903-370
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	

MINERAL OIL (UNII: T5L8T28FGP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Caplet)	Size	18mm
Flavor		Imprint Code	TCL370
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-370-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	07/30/2024	
2	NDC:79903-370-21	2 in 1 CARTON	07/30/2024	
2	NDC:79903-370-01	100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/30/2024	

Labeler - Walmart Inc. (051957769)

Registrant - TIME CAP LABORATORIES INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES INC		037052099	manufacture(79903-370)

Revised: 8/2024

Walmart Inc.