

**NIGHTTIME COLD AND FLU READYIN CASE- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid**  
**P & L Development, LLC**

-----  
**Drug Facts**

**Active ingredients (in each 30 mL)**

**Acetaminophen 650 mg**

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purposes**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves these common cold/flu symptoms:
- minor aches and pains
- headache
- sore throat
- fever
- runny nose and sneezing
- cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is

accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
  
- **When using other Day Time or Night Time products, carefully read each label to insure correct dosing**

## Other information

- **each 30 mL contains;** potassium 5 mg
- **each 30 mL contain** sodium 19 mg
- store between 20-25°C (68-77°F). Do not refrigerate

## Inactive ingredients

acesulfame potassium, alcohol, citric acid, D&C yellow 10, FD&C green 3, FD&C yellow 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

## Questions or Comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## Principal Display Panel

**\*Compare to active ingredients in Vicks® NyQuil®\* Cold & Flu\***

### Night Time

#### Cold & Flu relief

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

pain reliever/fever reducer

cough suppressant

antihistamine

relieves:

- fever,
- headache

- sore throat
- cough
- minor aches and pains
- runny nose & sneezing

for ages 12 years & over

alcohol 10%

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company, Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

Manufactured by:

PL Developments

11865 S. Alameda St

Lynwood, CA 90262

**Product Label**

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu<sup>®</sup> NDC 49580-0346-4

**ready in case**

**\$1.25**

**night time cold & flu relief**

**Acetaminophen 650 mg**

**Dextromethorphan HBr 30 mg**

**Doxylamine Succinate 12.5 mg**

pain reliever/fever reducer  
cough suppressant  
antihistamine

relieves:

- fever
- headache
- sore throat
- cough
- minor aches and pains
- runny nose and sneezing

for ages 12 years and over  
alcohol 10%

4 fl oz (118 mL)

**Drug Facts**

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg.....Pain reliever/fever reducer

Dextromethorphan HBr 30 mg.....Cough suppressant

Doxylamine succinate 12.5 mg.....Antihistamine

**Uses** ■ temporarily relieves common cold/flu symptoms ■ headache  
■ minor aches and pains ■ sore throat ■ runny nose and sneezing  
■ fever ■ cough due to minor throat and bronchial irritation

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if: ■ adults take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount of this product ■ child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

Manufactured by: **PL Developments**  
11865 S. Alameda St  
Lynwood, CA 90262

PLD-94078 1.0009646

**PARENTS:**  
Look about for another dose  
www.StupidCousin.com

**Drug Facts (continued)**

acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers

**Drug Facts (continued under label)** **PEEL HERE** →



**Drug Facts (continued)**

■ taking the blood thinning drug warfarin

**When using this product** ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions** ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device.  
■ mL = milliliter ■ keep dosing cup with product  
■ adults and children 12 years and over: 30 mL every 6 hours ■ children under 12 years of age: do not use  
■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information** ■ each 30 mL contains: potassium 5 mg ■ each 30 mL contains: sodium 19 mg ■ store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** acesulfame potassium, alcohol, citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or comments?** Call 1-877-753-3935  
Monday-Friday 9AM-5PM EST

**READYinCASE Night time Cold & Flu Relief**

<b>NIGHTTIME COLD AND FLU READYINCASE</b>			
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49580-0346
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)		DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49580-0346-8	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/30/2014	
2	NDC:49580-0346-2	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/30/2014	
3	NDC:49580-0346-4	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/30/2014	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/30/2014	

**Labeler** - P & L Development, LLC (101896231)

Revised: 2/2024

P & L Development, LLC