

SHISEIDO CLEAR SUNSCREEN SPF 50 PLUS- avobenzone, homosalate, octisalate, and octocrylene stick

SHISEIDO AMERICAS CORPORATION

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

SHISEIDO CLEAR SUNSCREEN STICK SPF 50+

Drug Facts

<i>Active ingredients</i>	<i>Purpose</i>
AVOBENZONE 2.5%	Sunscreen
HOMOSALATE 10.0%	Sunscreen
OCTISALATE 5.0%	Sunscreen
OCTOCRYLENE 10.0%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every two hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
 - children under 6 months: Ask a doctor

Inactive Ingredients

DIPHENYLSILOXY PHENYL TRIMETHICONE•MINERAL OIL•HYDROXYSTEARIC

ACID▪TRIETHYLHEXANOIN▪PEG/PPG-9/2 DIMETHYL ETHER▪METHYL METHACRYLATE
CROSSPOLYMER▪DIBUTYL LAUROYL GLUTAMIDE▪C12-15 ALKYL
BENZOATE▪POLYAMIDE-8▪SILICA DIMETHYL SILYLATE▪TOCOPHERYL
ACETATE▪PEG/PPG-14/7 DIMETHYL ETHER▪LECITHIN▪GLYCYRRHIZA GLABRA
(LICORICE) ROOT EXTRACT▪PPG-17▪CASTOR OIL/IPDI COPOLYMER▪CAPRYLIC/CAPRIC
TRIGLYCERIDE▪BHT▪TOCOPHEROL▪PEG-6▪PENTAERYTHRITYL TETRA-DI-T-BUTYL
HYDROXYHYDROCINNAMATE▪SILICA▪FRAGRANCE▪

Other information

Protect this product in this container from excessive heat and direct sun.

Questions or comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 20 g Stick Pouch Label

SHISEIDO
GINZA TOKYO

Clear Sunscreen Stick
50+

SynchroShield
WetForce × HeatForce

BROAD SPECTRUM
SPF 50+
For Face/Body
WATER RESISTANT
(80 MINUTES)
SUNSCREEN

BIO

RE
USE

20g NET WT. .7 OZ.

SHISEIDO

GINZA TOKYO

Clear Sunscreen Stick

50+



SynchroShield

WetForce x HeatForce

BROAD SPECTRUM

SPF 50+

For Face/Body

WATER RESISTANT

(80 MINUTES)

SUNSCREEN



20g NET WT. .7 OZ.



SHISEIDO
CLEAR SUNSCREEN STICK
BROAD SPECTRUM SPF 50+

A clear stick sunscreen that goes on smoothly without stickiness or a mess on your hands. Featuring SynchroShield™, its protective veil becomes even stronger when it detects heat, water or perspiration. This multi-protective stick provides a seamless defense against UVA/UVB rays and dryness while providing a skin emollient effect. Convenient for use anytime, anywhere.

- Containing the botanical ingredient complex Profense CL, it helps to prevent the appearance of aging signs such as wrinkles and uneven skin tone.
- Anti-pollution formula that helps to keep pollutants from adhering to the skin, and contains also an anti-oxidant to help to protect against damage that can be caused by them.

NON-COMEDOGENIC. DERMATOLOGIST AND OPHTHALMOLOGIST-TESTED.

Drug Facts

Active ingredients

Active ingredients	Purpose
AVOBENZONE 2.5%	Sunscreen
HOMOSALATE 10.0%	Sunscreen
OCTISALATE 5.0%	Sunscreen
OCTOCRYLENE 10.0%	Sunscreen

Uses ■ helps prevent sunburn ■ if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions For sunscreen use: ■ apply liberally 15 minutes before sun exposure ■ reapply: · after 80 minutes of swimming or sweating · immediately after towel drying · at least every 2 hours ■ Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including: · limit time in the sun, especially from 10 a.m. – 2 p.m. · wear long-sleeve shirts, pants, hats, and sunglasses ■ children under 6 months: Ask a doctor

Inactive ingredients DIPHENYLSILOXY PHENYL TRIMETHICONE · MINERAL OIL · HYDROXYSTEARIC ACID · TRIETHYLHEXANOIN · PEG/PPG-9/2 DIMETHYL ETHER · METHYL METHACRYLATE CROSSPOLYMER · DIBUTYL LAUROYL GLUTAMIDE · C12-15 ALKYL BENZOATE · POLYAMIDE-8 · SILICA DIMETHYL SILYLATE · TOCOPHERYL ACETATE · PEG/PPG-14/7 DIMETHYL ETHER · LECITHIN · GLYCYRRHIZA GLABRA (LICORICE) ROOT EXTRACT · PPG-17 · CASTOR OIL/POI COPOLYMER · CAPRYLIC/CAPRIC TRIGLYCERIDE · BHT · TOCOPHEROL · PEG-6 · PENTAERYTHRITYL TETRA-DI-T-BUTYL HYDROXYHYDROCINNAMATE · SILICA · FRAGRANCE ·

Other Information

■ protect this product in this container from excessive heat and direct sun.

Questions or Comments? Call toll free 1-800-906-7503



This very water resistant veil contributes to less elution when you enter the sea, compared to our non-water resistant sunscreens.



This bag is made from 72% certified biodegradable materials.



Please repurpose this nomadic pouch before discarding it with your trash.

Specially formulated by
Shiseido Laboratories, Japan.

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.

GLO. 16980

www.shiseido.com

<http://s1872.com/16980AA>

See our website for more information.



7 30852 16980 7

SHISEIDO CLEAR SUNSCREEN SPF 50 PLUS

avobenzone, homosalate, octisalate, and octocrylene stick

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-616
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	500 mg in 20 g
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	2000 mg in 20 g
OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	1000 mg in 20 g
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	2000 mg in 20 g

Inactive Ingredients

Ingredient Name	Strength
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
MINERAL OIL (UNII: T5L8T28FGP)	
12-HYDROXYSTEARIC ACID (UNII: 933ANU3H2S)	
TRIETHYLHEXANOIN (UNII: 7K3W1BIU6K)	
PEG/PPG-9/2 DIMETHYL ETHER (UNII: LM6Q66468X)	
METHYL METHACRYLATE/GLYCOL DIMETHACRYLATE CROSSPOLYMER (UNII: EG97988M5Q)	
ALKYL (C12-15) BENZOATE (UNII: A9EJ3J61HQ)	
DIBUTYL LAUROYL GLUTAMIDE (UNII: 3V7K3IA58X)	
POLYAMIDE-8 (4500 MW) (UNII: 77723GV81A)	
SILICA DIMETHYL SILYLATE (UNII: EU2PSPOG0W)	
PPG-17 (UNII: OV0Q322E0U)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
.ALPHA.-TOCOPHEROL ACETATE, DL- (UNII: WR1WPI7EW8)	
POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ)	
PENTAERYTHRITOL TETRAKIS(3-(3,5-DI-TERT-BUTYL-4-HYDROXYPHENYL)PROPIONATE) (UNII: 255PIF62MS)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
GLYCYRRHIZA GLABRA (UNII: 2788Z9758H)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-616-20	1 in 1 POUCH	01/01/2021	
1		20 g in 1 CARTRIDGE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part352	01/01/2021	

Labeler - SHISEIDO AMERICAS CORPORATION (193691821)

Revised: 11/2020

SHISEIDO AMERICAS CORPORATION